OPEN ACCESS

Mental health in elite athletes: International Olympic Committee consensus statement (2026)

Claudia L Reardon ,[1,2] Vincent Gouttebarge ,[3,4] Emily Kroshus-Havril ,[5,6] Cindy Miller Aron ,[1] Roald Bahr,[7,8] Cheri Blauwet ,[9] João Mauricio Castaldelli-Maia ,[10,11] Camille Cheng,[12] Alan Currie,[13] Jeffrey Lee Derevensky ,[14] Carla Edwards ,[15] Sarah Fussek,[16] Paul Gorczynski,[17] Michael Grandner,[18] Doug Hyun Han ,[19] Mary E Hitchcock ,[20] Frank Lu,[21] Andrew Massey ,[22] David McDuff ,[23] Margo Mountjoy,[24] Rosemary Purcell ,[25,26] Margot Putukian ,[27] Simon M Rice ,[25,28] Scott Sloan ,[7] Torbjørn Soligard ,[7] Jorunn Kaiander Sundgot-Borgen ,[29] Leslie Swartz ,[30] Jane S Thornton ,[7,31] Tshepang Tshube,[32] Brian Hainline [33]

► Additional supplemental material is published online only. To view, please visit the journal online (https://doi.org/10.1136/bjsports-2025-111514).

For numbered affiliations see end of article.

Correspondence to
Dr Claudia L Reardon; clreardon@wisc.edu

Accepted 1 June 2026
Published Online First 11 August 2026

ABSTRACT

This consensus statement represents an update to *Mental Health in Elite Athletes: International Olympic Committee Consensus Statement (2019)*. To advance a more standardised, evidence-based approach to mental health in elite athletes, an International Olympic Committee (IOC) Consensus Group critically evaluated the current science via a systematic literature review, sought consensus using Delphi methodology and provided recommendations relevant to this topic. Key themes include that mental health symptoms and disorders are (1) common among elite athletes; (2) may have sport-related aetiologies and manifestations; and (3) can impact performance. Routine screening for and monitoring of mental health symptoms and disorders in elite athletes using validated instruments is recommended. Management strategies should promote mental well-being in addition to preventing and treating mental health symptoms and disorders. These strategies should address all contributors to mental health symptoms, including factors within the environments in which athletes train and compete. Treatment may include psychotherapy, medication and other culturally or contextually appropriate modalities. With appropriate support and treatment, many athletes experiencing mental health symptoms and disorders can continue to train and compete safely, but sometimes modification of sport participation may be necessary. Elite athletes who are transitioning out of sport, elite Para athletes and elite young (adolescent) athletes are relatively understudied and may have distinct mental health concerns, in addition to those impacting elite athletes in general.

Check for updates

© Author(s) (or their employer(s)) 2026. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ Group.

To cite: Reardon CL, Gouttebarge V, Kroshus-Havril E, *et al*. *Br J Sports Med* 2026;**60**:1083–1130.

INTRODUCTION

In 2018, the International Olympic Committee (IOC) Medical and Scientific Commission established a Mental Health in Elite Athletes Consensus Group comprised of a variety of relevant disciplines. The group reviewed the best evidence to date in order to provide a consensus statement for clinical practice and individual and systemic interventions to improve mental health among elite athletes.[1] Since then, the IOC has undertaken several activities to implement their consensus statement findings in the 'real world' of elite sport and to respond to the field's unaddressed needs (box 1).

In the interim, there has been significant research productivity, confirming that mental health symptoms are commonly reported by current and former elite athletes as well as by their entourage members, including coaches and medical professionals.[2–10] The latest research also increasingly recognises two significant themes in how mental health in elite athletes is understood.

First, elite athlete mental health exists on a continuum.[11–14] The IOC defines the mental health continuum as a model that describes the range of mental health experiences an individual can have from optimal mental well-being to mental health symptoms and disorders.[11 12] Athletes' experiences on the continuum can and often do change over time.[13 15] While recognition of overt mental health disorders in elite athletes remains essential, situating athlete well-being within a dynamic continuum offers an additional framework for understanding the spectrum of psychological functioning in this population. For example, elite athletes may experience transient stress reactions related to competitions, to various temporary setbacks that are a normal and common part of elite athlete life, or to intense training periods.[13 16 17] Thus, there are important mental health considerations even if an athlete does not have a diagnosable disorder.[13]

Second, factors impacting elite athlete mental health are not confined to attributes of the individual athlete.[13 18–21] While this ecological approach to understanding determinants of mental health symptoms and disorders has long been part of discourse outside of sport,[22] this perspective has only recently become central to the understanding of mental health in sport. For example, the adaptation of an ecological systems model to elite athlete mental health by Purcell *et al*[18] describes the multiple layers of sport-related factors that influence elite athlete mental health (figure 1).

Taking these factors into account, the IOC convened a second consensus meeting on 10-12

Br J Sports Med: first published as 10.1136/bjsports-2025-111514 on 11 August 2026. Downloaded from http://bjsm.bmj.com/ on August 12, 2026 at Universitetsbiblioteket i Tromsø . Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies.

KEY POINTS

⇒ Mental health support for elite athletes should extend beyond individual management and should include assuring that the training environments, leadership behaviours, safeguarding systems and organisational culture are also supporting elite athlete mental health and well-being.
⇒ Clinicians and sport organisations should proactively address stigma, confidentiality concerns, fear of deselection and low mental health literacy as part of routine elite athlete care.
⇒ Sport organisations should implement routine screening and surveillance of elite athlete mental health at predefined time points and following significant events.
⇒ All elite sport settings should establish, rehearse and regularly review mental health emergency action plans with defined roles and escalation pathways.
⇒ Management of mental health symptoms and disorders should be individualised, evidence-based and culturally responsive, integrating psychotherapy, pharmacotherapy and other contextually appropriate modalities as indicated.
⇒ Psychological responses to injury should be routinely assessed, and return-to-sport for elite athletes with mental health symptoms and disorders should incorporate athlete safety, stability and function as guiding principles.
⇒ Athletes should receive targeted mental health screening and support before, during and after major transitions in sport.
⇒ Mental health assessment and care for elite Para athletes must account for disability-specific stressors, accessibility barriers, classification systems and heightened transition risks.
⇒ Mental health screening and support for elite adolescent athletes should reflect developmental stage, educational demands, family dynamics and the early onset of many common mental health disorders.
⇒ Coaches, support staff and health professionals for elite athletes should receive role-appropriate mental health education to support early recognition, safe referral and psychologically informed practice.

June 2025 in Lausanne, Switzerland, at which experts and athlete representatives reviewed the scientific literature addressing mental health in elite athletes. The Consensus Group defined elite athletes in the same way as the first consensus statement: those competing at professional, Olympic or collegiate (higher-education/tertiary) levels. The Consensus Group was charged with the following:

- To review the literature describing prevalence, screening and monitoring, diagnosis, impact on performance and management considerations regarding specific mental health symptoms and disorders in elite athletes.
- To review the literature describing stressors and environmental factors across ecological systems that impact elite athletes along the mental health continuum.
- To make actionable recommendations for health care providers of elite athletes using a consensus approach.
- To disseminate the above information in a consensus statement intended to serve as a resource for health care providers of elite athletes, other elite athlete entourage members and sports organisations.

An outline of this manuscript is provided as a guide in box 2.

Box 1 Major International Olympic Committee (IOC) work products addressing mental health in elite athletes since the organisation's 2019 consensus statement on mental health in elite athletes

⇒ Formation of the IOC Mental Health Working Group (standing working group of the IOC)
⇒ Development of a comprehensive set of mental health screening tools (IOC Sport Mental Health Assessment Tool 1 (SMHAT-1), IOC Sport Mental Health Recognition Tool 1 (SMHRT-1), IOC Sport Mental Health Assessment Tool 2 (SMHAT2) and IOC Sport Mental Health Recognition Tool 2 (SMHRT2)) for use in elite athletes[28 29]
⇒ Expansion of Athlete365 e-learning modules[783]
⇒ Creation of the IOC Mental Health in Elite Athletes Toolkit[11]
⇒ Publication of a mental health surveillance supplement to the IOC consensus statement on injury and illness surveillance[40]
⇒ Establishment of IOC Diploma and Certificate programmes on mental health in elite sport[784 785]
⇒ Creation of the IOC Mental Health Action Plan[12] as the IOC's strategic roadmap to promote mental well-being across the Olympic movement
⇒ Dissemination of IOC Mental Health Guidelines for Major Sporting Events[701 786]

METHODS

Consensus approaches are appropriate for aggregating expert opinion when there is a limited evidence base or no gold standard guidance that is appropriate across populations and settings.[23] For this project, consensus was sought using (1) literature reviews that informed candidate actionable guidance statements; (2) an in-person meeting to discuss the reviews and statements; (3) Delphi methodology to vote on and revise the statements; and (4) finalisation of the statements and consensus manuscript.[24] The consensus process (figure 2) was finalised by the consensus meeting co-chairs (VG, BH, CLR) and Delphi expert (EK-H) and is consistent with the *British Journal of Sports Medicine* author guidelines for consensus statements.[25]

Panel selection

The meeting co-chairs and IOC leadership from the Health, Medicine and Science Department (organising work group) identified members of a panel, ultimately comprised of 31 individuals with lived, clinical and/or scientific expertise in the mental health of elite athletes. Clinical or scientific expertise was determined by a literature search of peer-reviewed academic publications on the planned consensus sub-topics, as published during the previous 5 years. The panel also included an elite athlete representative (CC), selected from among the IOC's Mental Health Ambassadors, and two International Sports Federation representatives (SF and AM), selected based on their engagement in IOC mental health initiatives.

Equity, diversity and inclusion statement

Considerations for panel selection across all roles included inviting participants from different countries and regions of the world and from equity-deserving groups. To that end, panellists completed a brief demographic intake form (see online supplemental file 1) querying their race and ethnicity, gender identity, sexual orientation and disability status. These demographic data were reviewed by the organising work group as panel membership was finalised to ensure that equity-deserving groups were

Br J Sports Med: first published as 10.1136/bjsports-2025-111514 on 11 August 2026. Downloaded from http://bjsm.bmj.com/ on August 12, 2026 at Universitetsbiblioteket i Tromsø . Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies.




Figure 1 An ecological systems model for describing the multiple layers of sport-related factors that contribute to elite athlete mental health. Adapted and reprinted with permission from Purcell *et al*.[18]

represented. Countries represented on the panel were Australia, Botswana, Brazil, Canada, China, Hong Kong China, Norway, South Africa, South Korea, Switzerland, UK and USA.

Evidence review

The organising work group initially identified 27 focal topic areas, each with lead author(s) determined based on scientific and clinical expertise. An experienced librarian (MH) conducted a systematic search of each topic area spanning database inception to May 2024 using PubMed (Medline), SportDiscus (Ebsco), PsycINFO (Ebsco), Scopus (Elsevier) and any additional literature databases felt relevant by individual topic leaders, for each topic. Leaders of topics without a recent published systematic review were invited to write separate, more detailed, subspecialty papers on their topics; these will be published separately. Each topic team was asked to draft a review that summarised the key information from their literature search and to propose actionable guidance statements for clinicians based on their review. Written reviews and draft guidance statements were shared with panellists prior to the in-person meeting.

In-person meeting and discussion

The organising work group and other panellists met in person from 10 June to 12 June 2025 in Lausanne, Switzerland. The purpose of the meeting was to discuss the quality and level of evidence from the literature reviews, clinical and lived experiences relevant to the focal topics and any literature gaps. For each focal topic, panellists reviewed and critiqued the candidate actionable guidance statements for clinicians. Panellists were also encouraged to consider appropriateness and feasibility of candidate statements across populations and contexts. At the conclusion of the in-person meeting, participants were provided with an anonymous open-ended exit form to share any further suggestions about the candidate statements.

Determination of guidance statements for clinicians

Meeting discussion notes and output of the anonymous open-ended exit feedback form were reviewed by the organising work group and used to make edits to the candidate actionable guidance statements for clinicians. These statements were then emailed to panellists via an anonymous web-based survey. After each statement, participants were asked to rate the extent to which they agreed with including this guidance, with response options ranging from 1 (strongly disagree) to 9 (strongly agree). Participants who rated a guidance statement as <7 were asked to explain the reason for their score. The minimal threshold for a guidance statement to be included in the consensus output was set a priori as requiring two conditions to be met: (1) 70% or more of participants scoring as 7–9, and (2) no participants scoring as 1–3. The first condition (percent agreement) is among the most common approaches to operationalising consensus using the Delphi method.[26] The second condition was included to ensure that the perspectives of participants with specialised and unique skillsets or lived experiences were not overlooked because the group average for a given statement was high. There is precedent for using this type of more stringent approach to operationalising consensus in Delphi studies, pending the topic/nature of the guidance and the participant group.[26]

Open-ended feedback for statements not meeting the minimal consensus thresholds was discussed by the consensus meeting co-chairs, Delphi expert and relevant topic expert(s). They determined whether there were modifications that would address the concerns raised while keeping

Br J Sports Med: first published as 10.1136/bjsports-2025-111514 on 11 August 2026. Downloaded from http://bjsm.bmj.com/ on August 12, 2026 at Universitetsbiblioteket i Tromsø . Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies.

Box 2 Outline of *Mental Health in Elite Athletes: International Olympic Committee Consensus Statement (2026)*

⇒ Screening and surveillance of mental health symptoms and disorders in elite athletes
⇒ General approaches to management of mental health symptoms and disorders in elite athletes
  Psychotherapy
  Pharmacological treatment
  Mental health considerations for removal from and return to sport
⇒ Specific mental health symptoms and disorders in elite athletes
  Sleep disorders and sleep concerns
  Major depressive disorder and depression symptoms
  Suicide
  Anxiety disorders and obsessive-compulsive and related disorders
  Trauma- and stressor-related disorders
  Eating disorders and Relative Energy Deficiency in Sport (REDs)
  Attention-deficit/hyperactivity disorder
  Bipolar and psychotic disorders
  Personality disorders
  Substance misuse and use disorders
  Gambling disorder and other behavioural addictions
  Neuropsychiatric aspects of sport-related concussion
⇒ Stressors and environmental factors that impact elite athlete mental health
  Creating an environment that supports mental well-being and resilience in elite athletes
  Cultural considerations
  Barriers to mental health care seeking
  Social media and mental health
  How injury, performance and mental health intersect
  Mental health emergencies
  Transitions in elite sport
⇒ Elite Para athletes
⇒ Elite young (adolescent) athletes
⇒ Sport as a subset of physical activity to promote mental health
⇒ Final considerations
  Actionable recommendations
  Dissenting viewpoints
⇒ Discussion
⇒ Conclusion

the guidance consistent with the underlying evidence base. Participants were provided aggregated quantitative results and comments from each round of ratings and were asked to re-rate each statement that was modified, following the same process as outlined above. An a priori limit of no more than three rounds of ratings was established.[26] All guidance statements meeting the consensus threshold at the end of this process were included in the final consensus output. Any guidance statement included in this final output that had one or more ratings of ≤5 was highlighted as having dissenting opinions, with a synthesis of the open-ended feedback about the statement presented in the consensus output.

Final manuscript

The initial draft topic summaries were edited based on discussion at the meeting. The systematic literature searches were reconducted to identify materials published between 1 April 2024 and 30 June 2025, with duplicates removed by Endnote X20, using the same search strategies as initially used. The edited final document was circulated for review and further editing by the panel.

RESULTS

The results of the systematic literature reviews are summarised as follows.

Screening and surveillance of mental health symptoms and disorders in elite athletes

Routine screening and surveillance for mental health symptoms and disorders at regularly scheduled time points and at the time of significant occurrences/events remain uncommon in elite sports.[27] Mental health *screening* involves the use of tools to identify individual athletes who may be experiencing mental health symptoms or disorders in order to facilitate appropriate support. Mental health *surveillance* involves the systematic collection, analysis and interpretation of athlete mental health data for either clinical or research purposes. Clinically, surveillance can be used to identify and monitor athletes who need mental health support and, from a research perspective, surveillance at a cohort level can identify mental health trends and risk factors and inform interventions.

Screening

Few mental health screening tools have been validated in elite athletes. The IOC Sport Mental Health Assessment Tool 1 (SMHAT-1) was developed for sports medicine physicians and other licensed/registered health professionals to identify elite athletes (aged ≥16 years) potentially experiencing mental health symptoms and disorders to facilitate timely referral of those in need of support and/or treatment.[28] Based on evaluations of the accuracy and utility of the SMHAT-1, this measure was revised and the IOC Sport Mental Health Assessment Tool 2 (SMHAT2) was developed to provide a more comprehensive three-step approach including: universal screening (five validated disorder-specific screening questionnaires) as step 1, conditional screening (two validated disorder-specific screening questionnaires) as step 2 and an aid to clinical assessment and management as step 3.[28–30] Importantly, steps 1 and 2 of the SMHAT-1 and SMHAT2 are symptom screeners but not diagnostic tools, with step 3 of these tools focusing on clinical assessment and diagnosis that should be conducted by clinicians.

Surveillance

The IOC published a consensus statement in 2019 on injury and illness surveillance,[31] which defined guideline procedures for monitoring athlete health in a variety of sporting contexts, such as over the course of a sports season or at major sports events. Subsequently, sport-specific statements with recommendations tailored for several specific sports[32–37] and for Para sport[38] were published. Supplemental manuscripts to address female athlete nuance[39] and specifically athlete mental health symptoms and disorders[40] were also subsequently published.

Br J Sports Med: first published as 10.1136/bjsports-2025-111514 on 11 August 2026. Downloaded from http://bjsm.bmj.com/ on August 12, 2026 at Universitetsbiblioteket i Tromsoe . Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies.




Figure 2 Methodology for the International Olympic Committee consensus meeting on mental health in elite athletes.

The supplemental manuscript on athlete mental health symptom and disorder monitoring defines the principles of athlete mental health surveillance including terminology, classification coding, mode of onset, aetiology, severity and tracking of multiple and/or recurrent episodes. Practice guidelines to implement athlete mental health surveillance (when, who and how) are discussed. Finally, this publication provides four examples of mental health surveillance tools that can be used depending on the resources available, the timing and location of the surveillance, the study design and/or the clinical scenario. Table 1 summarises the recommended surveillance tools.

General approaches to management of mental health symptoms and disorders in elite athletes

Management of mental health symptoms and disorders in elite athletes should consider all the psychological, general medical, sociocultural and ecological factors that may contribute to their presentation. Clinicians (licensed health care professionals) should consider the most appropriate consensus or evidence-based recommendations from multidisciplinary sources, while recognising cultural diversity as well as differences in access to and acceptability of certain treatments.[1]

Br J Sports Med: first published as 10.1136/bjsports-2025-111514 on 11 August 2026. Downloaded from http://bjsm.bmj.com/ on August 12, 2026 at Universitetsbiblioteket i Tromsø . Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies.

Table 1 Summary of the types of mental health surveillance tools, timing of data collection for each type, the recommended tools within each type and the roles of the athlete and the health professional

Type of surveillance	Timing of data collection	Tool	Athlete role	Health professional role
Screening	At specific or episodic time points (eg, pre-season, post major event, significant injury)	IOC SMHAT2	Data entry: completion of questionnaire(s)	To analyse and respond to athlete reporting
Longitudinal athlete-based surveillance	At regular predetermined time intervals throughout the season (eg, weekly, bi-weekly, monthly)	MH OSTRC-H (online supplemental file 2)	Data entry: completion of questionnaire	To analyse and respond to athlete reporting
Longitudinal surveillance by a health professional	Throughout the entire season as MH problem(s) occur	Modified IOC Season Surveillance Tool (online supplemental file 3)	Reporting of MH symptoms to the entourage health care provider	Data entry: to record cases that receive medical attention; to analyse and respond to athlete reporting
Sport event-related surveillance	At regular predetermined time intervals during sporting events (eg, daily reporting during Olympic Games, Tournaments, World Championships)	Modified IOC (Multi-) Sport Event Surveillance Tool (online supplemental file 4)	Reporting of MH symptoms to the entourage health care provider	Data entry: to record the cases that receive medical attention; to analyse and respond to athlete reporting

Table (modified) and supplemental files reprinted with permission from Mountjoy *et al*.[40]
IOC, International Olympic Committee; MH, mental health; OSTRC-H, Oslo Sport Trauma Research Center-Health; SMHAT2, Sport Mental Health Assessment Tool 2.

Psychotherapy

Psychotherapy is defined as the application of psychological modalities for the treatment of mental health symptoms and disorders, problems of living and/or facilitation of personal growth.[41 42] It is generally regarded as first-line treatment, where it is available, for individuals with mild to moderate mental health symptoms or disorders, either as monotherapy or combined with other non-pharmacological and pharmacological strategies. Principles of psychotherapy form an important foundational component of the management of mental health symptoms and disorders in elite athletes.[43] Psychotherapy is a form of treatment delivered by psychotherapists (licensed providers trained in specific psychotherapeutic modalities), distinguishing it from the many other types of support, guidance and mental health promotion that can be delivered outside of a treatment setting by individuals such as faith leaders, performance coaches, advisors, mentors and other members of the athlete support group.

Multiple psychotherapeutic modalities have shown empirical effectiveness across a range of mental health symptoms and disorders, particularly within general populations (table 2). Psychoeducation is an evidence-based foundational feature across psychotherapeutic modalities that provides information for patients and their supports to better understand and cope with mental health symptoms and disorders. Incorporating psychoeducational factors into psychotherapy can reduce stigma, normalise responses, clarify purposes of therapeutic interventions and create a shared language for athletes and their support networks, all of which support long-term change.[44] Furthermore, it provides a proactive approach to assist athletes in developing skills and competencies to cope with sporting career challenges and challenging environments.[45] From a performance standpoint, psychotherapy can bring together tools from mental health treatment and from performance psychology to help elite athletes achieve optimal performance. Examples include well-being frameworks grounded in positive psychology and performance science, which emphasise the optimisation of mental states, the cultivation of resilience and the capacity to function effectively under pressure.[46 47]

Essential for psychotherapy treatment planning is understanding the multifactorial elements leading to the athlete's struggles and contributing to ongoing challenges, balanced with the athlete's inherent strengths.[48] A biopsychosocial formulation is a valuable tool that includes cultural, ethnic and socioeconomic considerations.[49] When engaging elite athletes in psychotherapy, the psychotherapist must be aware of how the athlete's personal factors (including personality, beliefs and misgivings) and sport-related factors (including schedules, demands and expectations) may impact treatment. Incorporating an appreciation of the ecological systems (eg, family, coaches, teammates, national governing bodies) in which the athlete functions is important.[43]

Reluctance to engage in psychotherapy can be due to stigma, minimisation of symptoms, scheduling conflicts that prioritise training and fear. Mindfulness-based therapies can sometimes be accessible first interventions in the face of these barriers, particularly given their efficacy regarding overall health and performance.[50] Flexibility in the type of medium through which psychotherapy is delivered may also help overcome barriers. Although there are limited data on the use of online delivery of psychotherapy with elite athletes, there is evidence of its efficacy and efficiency.[51] It is essential to consider safety, stability and function when determining the most appropriate method and location of evaluation and treatment.[52] Elite Para athletes can have additional barriers to engaging in psychotherapy. Factors that may impact accessibility for this group include stigma, sensory impairment and intellectual disabilities.[53] Providing access to diverse treatment providers may facilitate engagement and inclusion in psychotherapy across all groups of elite athletes.

The transnational mobility of elite athletes and coaches, coupled with the complexity and diversity of those who seek mental health support, necessitates a culturally responsive and inclusive approach to psychotherapy.[54 55] Psychotherapists must ground themselves in the cultural contexts of their clients. Effective psychotherapists accommodate the language and culture of the client. A significant number of elite athletes from indigenous communities and low- and middle-income countries do not have access to psychotherapists, and psychotherapists may not speak the most common languages in these areas. In addition, some elite athletes and coaches find benefit in the use of approaches to wellness that may lack a robust evidence base,[55 56] but that are important in their culture. For example, athletes in sub-Saharan Africa have used verbal incarnations, juju and psyching verses as a source of intrinsic motivation and coping mechanisms for dealing with adversity.[56 57] Consequently, providers across cultures must collaborate and share knowledge to provide contextual support and bridge any language/cultural gaps.

Br J Sports Med: first published as 10.1136/bjsports-2025-111514 on 11 August 2026. Downloaded from http://bjsm.bmj.com/ on August 12, 2026 at Universitetsbiblioteket i Tromsø . Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies.

Table 2 Select types of psychotherapy for targeted mental health symptoms and disorders

Psychotherapy type	Targeted symptoms and disorders
Acceptance and commitment therapy[787]	► Transdiagnostic
Cognitive and behavioural therapies*[788]	► Insomnia (CBT-I) ► Depression ► Anxiety disorders ► Obsessive-compulsive disorder ► Post-traumatic stress disorder ► Eating disorders ► Attention-deficit/hyperactivity disorder ► Bipolar disorder ► Schizophrenia ► Personality disorders ► Substance misuse and use disorders ► Behavioural addictions ► Relationship distress ► Anger ► Chronic pain
Dialectical behaviour therapy[402 789–791]	► Depression ► Anxiety disorders ► Post-traumatic stress disorder ► Bulimia and binge-eating disorder ► Borderline personality disorder ► Substance misuse and use disorders ► Self-injurious behaviour ► Suicidal thoughts and behaviours
Eye movement desensitisation and reprocessing (EMDR)[792]	► Post-traumatic stress disorder
Interpersonal psychotherapy[793]	► Interpersonal deficits ► Grief and bereavement ► Role dispute ► Role transition
Mindfulness-based therapies*[794–796]	► Depression ► Anxiety disorders ► Post-traumatic stress disorder ► Substance misuse and use disorders ► Chronic pain
Motivational enhancement therapy*[470 797]	► Substance misuse and use disorders ► Behavioural addictions
Psychodynamic psychotherapy[798]	► Depression ► Anxiety disorders ► Eating disorders ► Personality disorders ► Chronic pain ► Relationship distress
Social rhythm therapy[799]	► Bipolar disorder
Solution-focused therapy[800]	► Transdiagnostic
Supportive psychotherapy[801]	► Transdiagnostic

*Evidence exists for elite athletes.
CBT-I, cognitive behavioural therapy for insomnia.

Pharmacological treatment

Pharmacological treatment is generally considered when mental health symptoms or disorders are moderate to severe and/or they cause significant dysfunction.[1] Typically, a combination of medication and psychotherapy is regarded as superior to medication alone.[58] Clinicians should consider the degree of distress or functional impairment, diagnosis and athlete willingness to take medication.[48] Pharmacological treatment of mental health disorders in elite athletes should take into account established treatment algorithms and clinical practice guidelines, such as those from national psychiatric associations, which have been shown to improve outcomes in the general population.[48 59–62] However, additional considerations for elite athletes are necessary. Specifically, health care providers should understand: (1) any potential negative impacts of the medication on sport performance; (2) any safety risks of the medication when used by elite athletes; and (3) whether the medication is prohibited by any sport governing body, such as the World Anti-Doping Agency (WADA).[1]

Athletes and their providers should be aware of and monitor for potential side effects that may interfere with performance, without forgoing effective treatment out of fear of side effects, which can often be managed. In a 2024 international survey of sports psychiatrists, the side effects that most commonly impact their choice of psychiatric medications for athletes were: (1) sedation (which can be both a performance and a safety issue); (2) weight gain; and (3) general reduction in performance from any cause.[63]

Medications may improve sports performance by virtue of improvement in the disorder they were designed to treat. For example, if an athlete's depression improves with use of an antidepressant medication, this may be associated with improved sports performance as they return to their baseline interest in and motivation for sport.[1] However, if a medication could produce a performance advantage beyond that attributable to improvement in the condition (eg, stimulants in athletes without attention-deficit/hyperactivity disorder), then it may be considered ergogenic and would likely be prohibited by WADA and/or other governing bodies.[64] If an elite athlete who is subject to such rules has a medical condition that requires a prohibited medication, then they must follow the rules of their governing body(s), which often involves application for a Therapeutic Use Exemption (TUE) (or similar application) from the athlete's relevant anti-doping organisation.[65]

Table 3 shows the broad categories of psychiatric medications, symptoms/disorders commonly treated by each, relevant side effects/safety concerns for elite athletes, potential performance impact and prohibitions in elite sport. There are several caveats for the information included in this table. First, data on the use of psychiatric medications in elite athletes are limited, with frequently underpowered sample sizes (often n<10), typically single-dose designs (rather than real-world chronic use at typical dosages) and few female subjects. Additionally, participants in these studies typically have not had the mental health

Br J Sports Med: first published as 10.1136/bjsports-2025-111514 on 11 August 2026. Downloaded from http://bjsm.bmj.com/ on August 12, 2026 at Universitetsbiblioteket i Tromsø . Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies.

Table 3 Different categories of psychiatric medications and features of each category relevant to elite athletes

Medication category	Symptoms/disorders most commonly treated	Relevant side effects/safety concerns	Performance impact	Prohibitions
Sedative-hypnotics	▶ Insomnia and other sleep disorders ▶ Anxiety disorders	▶ Sedation including hangover effect[1 66] ▶ Benzodiazepines (especially those with short half-lives)[802] may cause dependence[803–805]	▶ Inconsistent impact of benzodiazepines and imidazopyridines (eg, zolpidem) per two systematic reviews, with possible greater negative impact from medications in these classes with longer half-lives (n=9 and n=10 studies, respectively)[66 806] ▶ Several studies[807 808] and a systematic review (n=15 studies)[161] have demonstrated no consistent negative impact of melatonin	▶ Melatonin and magnesium are non-regulated supplements in some countries, and thus as with all such products could unknowingly contain prohibited substances[63 366]
Antidepressants	▶ Depression ▶ Anxiety disorders and obsessive-compulsive and related disorders ▶ Trauma- and stressor-related disorders ▶ Sleep disorders (sedating antidepressants) ▶ Eating disorders (especially fluoxetine) ▶ Pain	▶ Serotonin-enhancing medications may cause gastrointestinal upset (often temporary) ▶ Bupropion may increase risk of heat illness[809] and, in athletes with eating disorders, of seizures[810] ▶ TCAs and mirtazapine are relatively more likely than other antidepressants to cause sedation and weight gain[366] ▶ TCAs may cause orthostatic dizziness, cardiac side effects (eg, tachycardia, supraventricular and ventricular arrhythmias, chronotropic incompetence)[811 812] and theoretically blood levels could fluctuate with exercise[813] ▶ Some antidepressants may cause emotional blunting[814]	▶ Single-dose bupropion at high dosing (600 mg) may be associated with performance enhancement in warm[809 815] but not temperate[816] conditions ▶ Inconsistent findings (neutral or negative impact on performance) from six studies on the SSRI paroxetine[817–822] ▶ Two studies on the most activating SSRI, fluoxetine, demonstrated no impact on performance after acute[319 823] or continuous dosing[823] ▶ Several studies on single-dose reboxetine (NRI) demonstrated a negative impact on performance per systematic review (n=5 studies)[66]	▶ Bupropion is currently included in WADA's *in competition* monitoring programme (being monitored for patterns of misuse in sport but not prohibited)[824]
Other anxiolytics	▶ Anxiety disorders	▶ Buspirone may cause dizziness[825] ▶ Beta-blockers (used for performance anxiety in non-athletes) may cause decreased pulse and blood pressure[826]	▶ Single 45 mg dose of buspirone demonstrated negative impact on performance in one study[827] ▶ Beta-blockers may problematically decrease cardiopulmonary capacity[828]	▶ Beta-blockers are prohibited *in competition* by WADA in automobile, billiards, darts, golf, mini-golf, and some skiing/snowboarding sports and both in and out of competition in archery, shooting, and underwater sports[64]
Stimulants and other ADHD medications	▶ ADHD ▶ Binge-eating disorder ▶ Narcolepsy and related sleep disorders	▶ Stimulants may cause weight loss, insomnia, anxiety, cardiac side effects (eg, tachycardia), and increased risk of heat illness[829] ▶ Atomoxetine and viloxazine may cause gastrointestinal discomfort and sedation or insomnia[343 830] ▶ Guanfacine and clonidine may cause sedation and cardiac side effects (eg, hypotension, bradycardia, QTc prolongation)[831 832]	▶ Stimulants may cause performance enhancement, especially in the heat, per several studies[829 833–835] and a systematic review (n=4 studies)[66]	▶ Stimulants are prohibited *in competition* by WADA,[64] the NCAA,[836] and many professional sports leagues
Mood stabilisers	▶ Bipolar disorder	▶ Sedation ▶ Weight gain ▶ Tremor ▶ Lithium blood levels may fluctuate with exercise[837 838] and increase with concomitant NSAID use[839 840] ▶ Polyuria (lithium)[839]	▶ Unknown	–

Continued

Br J Sports Med: first published as 10.1136/bjsports-2025-111514 on 11 August 2026. Downloaded from http://bjsm.bmj.com/ on August 12, 2026 at Universitetsbiblioteket i Tromsø . Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies.

Table 3 Continued

Medication category	Symptoms/disorders most commonly treated	Relevant side effects/safety concerns	Performance impact	Prohibitions
Antipsychotics	► Psychotic disorders ► Bipolar disorder ► Antidepressant augmentation	► Sedation ► Weight gain ► Tremor and other movement disorder side effects[366] ► QTc prolongation (especially ziprasidone and typical antipsychotics)[841] ► Increased risk of heat illness[842 843]	► Impairing effects per single studies each of oxypertine,[844] olanzapine,[845] risperidone,[845] and clozapine[846]	–
Medications for substance use disorders	► Withdrawal and cravings in substance use disorders	► Varenicline may cause nausea, insomnia, and vivid dreams[847] ► Naltrexone may cause gastrointestinal upset ► Nicotine vaping devices (used by some as a smoking cessation intervention) may have a negative impact on lung physiology and immune function[848]	► Unknown	► Opioid agonists (eg, buprenorphine, methadone) prohibited *in competition* by WADA[64]

ADHD, attention-deficit/hyperactivity disorder ; NCAA, National Collegiate Athletic Association; NRI, norepinephrine reuptake inhibitor ; NSAID, non-steroidal anti-inflammatory drug; SSRI, selective serotonin reuptake inhibitor; TCA, tricyclic antidepressant; WADA, World Anti-Doping Agency.

disorders the medications are designed to treat,[66] making the results difficult to interpret. Second, the availability and usage of psychiatric medications varies across countries. Relatedly, elite athletes who are travelling for sport and their prescribing clinicians should be aware that they may not be allowed to bring particular medications, especially controlled substances such as stimulants, opioids and benzodiazepines, across geographic borders into certain countries.[67] Third, non-traditional, experimental, controversial and/or emerging pharmacotherapies such as psychedelics (eg, psilocybin) are not included in this table, given the limited evidence base.[68] An important concern with such substances is that purity may not be guaranteed, which risks adverse analytical findings on drug testing. Fourth, for elite Para athletes there may be complex medical backgrounds impacting medication choices.[69 70] Finally, while this table highlights potential issues of relevance to elite athletes, it is not meant to dissuade consideration of medication treatment, which can be helpful and necessary, for this population.

Mental health considerations for removal from and return to sport

Elite athletes may experience a continuum of mental health experiences ranging from optimal mental well-being to the presence of mental health symptoms and disorders,[71] and can often continue to meet the demands of training and competition.[72] When mental health symptoms or disorders are experienced on an intense or disruptive level, there may be occasions when the athlete's health care team and/or sport team must decide if the athlete should be removed from the sport environment. While protocols for modifications and interruptions related to injury or general medical illness are relatively well defined in the sports medicine literature, protocols related to mental health symptoms and disorders (outside of eating disorders) are not well defined. This is likely due, in part, to the complex influences that various types of mental health symptoms have on cognition, behaviour, function and performance, in addition to the fact that individuals who experience mental health symptoms can have preservation of function and stability much of the time. However, individuals who experience mental health symptoms or disorders may experience impairment in their sport participation and performance.[72]

Removal from sport (RFS) involves the intentional exclusion of the athlete from training and/or competition. RFS may be partial, with defined restrictions and monitoring, or complete, with no form of training or competition permitted. Having a mental health diagnosis by itself should not preclude sport participation; however, it is important to recognise how certain mental health symptoms or disorders can potentially influence sport participation. Symptoms and disorders that can sometimes compromise safety, stability and function (table 4) may result in the athlete either having to modify sport participation with medical monitoring or having to be removed from sport participation until such a time that they are deemed suitable to return.[72]

As a result of these complexities, an athlete's suitability to participate in sport should be assessed within a framework that is anchored in the key elements of safety, stability and function.[72] Specifically, safety must be considered from medical and psychological perspectives. Medical safety involves the influence of one's medical state on overall morbidity and mortality, including cardiovascular, respiratory, neurological, neuropsychiatric, metabolic and musculoskeletal elements. Psychological safety involves the influence of cognitive, emotional and behavioural elements on the risk of physical or psychological harm to self or others. Stability must also be considered from medical and psychological perspectives, both of which should be further defined according to a specified time period or exposure to varying conditions as appropriate. Medical stability entails the ability to sustain medical wellness, while psychological stability involves the ability to sustain cognitive, emotional and behavioural wellness. Function reflects the person's ability to attend to their personal needs, navigate fundamental skills required to independently care for themselves (ie, activities of daily living), fulfil responsibilities and engage in meaningful relationships.

Return to sport (RTS) often entails clearance of an athlete by a licensed clinician before training or competition resumes. Clearance may be graduated (with adjustments guided by monitoring and benchmarks), partial or complete. When an athlete has been

Br J Sports Med: first published as 10.1136/bjsports-2025-111514 on 11 August 2026. Downloaded from http://bjsm.bmj.com/ on August 12, 2026 at Universitetsbiblioteket i Tromsø . Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies.

Table 4 Examples of impairment caused by select mental health symptoms and disorders that can interfere with continued sport participation

Mental health symptoms or disorders*	Potential negative impact on sport
Depression[176]	▶ Lack of engagement ▶ Amotivation ▶ Insomnia or other factors leading to low energy ▶ Negative thinking ▶ Negative body language ▶ Distractibility ▶ Social isolation ▶ Emotional dysregulation ▶ Low appetite leading to inadequate intake
Anxiety, obsessive-compulsive and trauma/stressor-related disorders[176]	▶ Distractibility ▶ Insomnia or other factors leading to low energy ▶ Emotional dysregulation ▶ Poor confidence ▶ Low appetite or gastrointestinal upset leading to inadequate intake ▶ Inability to execute tasks (may be due to 'fight, flight or freeze' response, fear of negative judgment or failure, lack of confidence or negative self-talk) ▶ Panic attacks during sport (which affect breathing, cognition, heart rate and physical and mental composure) ▶ Interference with function via the need to perform rituals or compulsions ▶ Avoidance of sport circumstances reminiscent of past traumas
Eating disorders or disordered eating[176 849 850]	▶ Low energy (may be associated with Relative Energy Deficiency in Sport or insomnia associated with eating disorders) ▶ Electrolyte imbalances and cardiac risks may be amplified with exertion ▶ Cognitive fogginess, decreased explosivity and reduced endurance during sporting activities
Bipolar mania[176 349]	▶ Intrusive and/or disruptive behaviour ▶ Distractibility ▶ Emotional dysregulation ▶ Aggression ▶ Inflated self-esteem or self-appraisal ▶ Sexual disinhibition ▶ Impaired insight and judgment ▶ Intensity/stimulating environment of sport could perpetuate manic symptoms
Psychosis[176 349 851 852]	▶ Disruption in the sport environment due to: – Delusions (which may appear as confrontations or accusations against teammates and staff) – Bizarre behaviour and unusual thought patterns – Impaired cognitive/executive function – Negative symptoms (including restricted emotional expression and avolition)
Personality disorders[176]	▶ Sanctions, legal difficulties, anti-doping violations, injuries and conflict with team
Substance misuse or use disorders[176 418]	▶ Sport performance while under the influence of (or experiencing withdrawal from) substances may be negatively impacted by: – Impaired judgment, concentration, cognitive processing, reaction time, motor coordination, and cognitive awareness – Emotional dysregulation – Insomnia or hypersomnia ▶ Relationships with teammates and staff may be strained if substance use and its effects interfere with: – Team environment – Individual behaviour – Performance ▶ Substance use may lead to positive drug tests, which can compromise the career of the individual and have effects on the team

*Symptoms may either fulfil clinical criteria for diagnosis or may not fulfil full criteria yet result in clinically significant impairment in safety, stability and/or function.

away from the sport environment due to mental health symptoms or disorders, the athlete's health care team should be deliberate and collaborative in guiding their RTS.[72] It is important to consider the elements of the athlete's sport ecosystem when planning for RTS. Specific elements that are bidirectionally impacted by an athlete's RTS include teammates, coaches, staff, family, organisations and other aspects of their sport environment. Consideration needs to be given to the potential risks associated with training and competing (both at home and on the road). These elements and the athlete's overall symptom load can be considered under the same three themes of safety, stability and function.

Treatment contracts or agreements may be helpful to establish conditions by which the reintegration into sport can be facilitated and supported with ongoing monitoring. This can be particularly helpful in cases involving eating disorders,[72 73] behavioural disturbances and severe mental health disorders that require adherence to treatment protocols.

Specific guidelines have been established for exclusion or withdrawal from and return to sport for individuals with anorexia nervosa and bulimia nervosa,[74–76] but not for other mental health symptoms and disorders. In any case, RFS and RTS decision-making and planning should involve multidisciplinary collaboration and monitoring and, to the extent it is safe to do so, shared decision-making with the athlete.[77–79]

Specific mental health symptoms and disorders in elite athletes

Sleep disorders and sleep concerns

Prevalence

The prevalence of diagnosed sleep disorders among elite athletes is not well established, primarily because these conditions are

Br J Sports Med: first published as 10.1136/bjsports-2025-111514 on 11 August 2026. Downloaded from http://bjsm.bmj.com/ on August 12, 2026 at Universitetsbiblioteket i Tromsoe . Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies.

under-diagnosed.[80 81] However, data from several sources estimate that average sleep duration among elite athletes is below the recommended 7 hours, with typical sleep duration being approximately 6.5 hours[81–84] and athletes reporting that they do not get sufficient sleep an average of 3.8 days per week.[85] Regarding poor sleep quality, studies report prevalence estimates of about 20%, and up to 50% depending on the sport and setting.[85–91] For example, among a cohort of Olympic athletes, 50% demonstrated poor sleep quality.[86] Among elite Paralympic athletes, disturbances in sleep amount and quality have been reported across South African,[92] Israeli,[92] Brazilian[93 94] and Swedish[95] cohorts; across cohorts, approximately one-third reported excessive sleepiness, one-third reported habitual sleep duration of ≤6 hours and >50% reported poor sleep quality.

Regarding other sleep disorders, insomnia disorder appears prevalent among elite athletes. Of collegiate athletes, 22% reported persistent difficulty falling asleep at least 3 nights per week and, relatedly, 61% reported excessive daytime tiredness at least 3 days per week.[85] Other studies in elite athlete cohorts estimate the prevalence of likely insomnia disorder to be >25%.[96 97] The prevalence of sleep apnoea is not well-established, but it is estimated that 8% of collegiate American football players are at high risk of sleep apnoea,[98] while ice hockey players had a 10% prevalence[99] and 24% of a sample of elite rugby players had been diagnosed with sleep apnoea.[100] The prevalence of restless legs syndrome is not well-known but is estimated to be approximately 13% among elite runners.[101] Circadian rhythm sleep-wake disorders may be common among elite athletes.[102] Specifically, delayed sleep-wake phase disorder is especially common among younger adults[103–106] due to normal delays in circadian rhythms that occur in adolescence and young adulthood. This has been observed among students and others who require early awakening for morning activities.[102 107] Finally, other disorders also disproportionately common among adolescents and young adults (eg, non-REM parasomnias, narcolepsy) may be relatively more common among elite athletes who are in these age groups,[108–111] but prevalence rates compared with general populations are unknown.

Diagnosis

Insomnia disorder is diagnosed based on patient report or prospective sleep diary and is defined as a persistent difficulty initiating or maintaining sleep at least 3 nights per week, lasting at least 3 months despite adequate opportunity for sleep and resulting in some kind of daytime impairment.[112] Typically, taking at least 30 minutes to fall asleep, being awake for at least 30 total minutes trying to resume sleep during the night and/or awakening at least 30 minutes before desired awakening time are criteria for likely diagnosis.[113] This is differentiated from insomnia symptoms, such as more transient, infrequent, situational and/or less severe difficulties falling and/or staying asleep. Sleep apnoea is defined as the presence of at least five respiratory events (eg, apnoeas, hypopnoeas) per hour during sleep as observed using diagnostic equipment at home or in-laboratory. Sleep apnoea can be obstructive (most events are due to an obstruction of the airway), central (most events are due to a reduced respiratory drive, absent physical obstruction) or mixed (a combination of both types).[112] Restless legs syndrome is a neuromuscular disorder that is diagnosed by patient report and includes an uncomfortable (non-painful) sensation in the legs that occurs predominantly at night or when lying down, creates an urge to move and is relieved by movement.[112] Diagnosis of the latter may also be supplemented by serum ferritin levels, as levels ≤50 µg/L indicate risk.[112 114 115] Differential diagnosis should consider nocturnal leg cramps (which can be painful and related to cardiovascular and/or musculoskeletal issues)[116] and periodic limb movement disorder (a neuromuscular condition that involves involuntary movements during sleep).[117] Circadian rhythm sleep-wake disorders are typically diagnosed by self-report and arise when the available sleep period is misaligned relative to the endogenous circadian rhythms.[118 119] For example, delayed sleep-wake phase disorder occurs when an individual's natural sleep phase is later in the night, but the preferred sleep phase (eg, per schedule demands) is earlier. Non-REM parasomnias are also typically diagnosed by patient report and include incomplete arousals from sleep in the form of sleep-talking, sleep-walking, night terrors or other similar wake-like behaviours that are occurring in non-REM sleep.[110 112] Hypersomnia disorders such as narcolepsy and idiopathic hypersomnia are diagnosed using a combination of patient report and in-laboratory observation.[112 120 121] Sleep specialists (sleep medicine or behavioural sleep medicine) are important referrals for assuring proper diagnostic evaluation, although access to specialised clinicians and/or proper diagnostic tools and technology may be a barrier to diagnosis and care.

Impact

Shorter sleep duration among athletes has been associated with worse physical and mental performance, increased injury risk,[122] impaired recovery and increased risk for a range of mental health symptoms and disorders, including suicidal ideation.[1 80 81 85 122–126] Irregularity of sleep schedules has been associated with worse physical and mental performance and impaired general health.[107 127–129] Unhealthy timing of sleep (sleeping at a time inconsistent with the body's internal sleep-wake rhythms) has been associated with depressed mood, increased anxiety and worse performance.[130 131]

Insomnia disorder is associated with impaired athletic performance and increased risk for adverse mental health outcomes among athletes. Prospective data indicate that baseline insomnia disorder and daytime sleepiness increase the likelihood of sustaining a sport-related concussion within the subsequent year.[132] Furthermore, meta-analyses show that insomnia is a reliable predictor of depressive disorders, anxiety disorders and substance use disorders. Insomnia is also a well-established suicide risk factor, with symptoms that persist for extended periods of time (eg, >3 months) and/or insomnia disorder associated with an approximate tripling of suicidal ideation, attempts and deaths.[125 133–139] Untreated sleep apnoea, regardless of aetiology, increases fatigue, elevates risk for cardiometabolic disease (eg, obesity, dyslipidaemia, inflammation and hypertension),[140–142] is associated with depression and other mental health symptoms[143 144] and impairs athletic performance.[145–147] Repetitive head impact exposure in contact sports can also be causally implicated in REM Behaviour Disorder.[148]

Management

To improve sub-clinical sleep issues and sleep and circadian health in general (eg, sleep duration, sleep timing, sleep quality, daytime sleepiness), current guidelines include a combination of regular screening and education.[80 81 102 149] Additional components may include tracking (eg, wearables),[150–152] schedule modification,[149 153 154] environmental manipulations[149 155] and instituting programmes around sleep banking and adapting to travel.[81 102]

Br J Sports Med: first published as 10.1136/bjsports-2025-111514 on 11 August 2026. Downloaded from http://bjsm.bmj.com/ on August 12, 2026 at Universitetsbiblioteket i Tromsø . Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies.

Non-pharmacological/behavioural strategies are often recommended for sleep disorders in elite athletes because they are effective and because many medications are associated with increased injury risk, unacceptable side effects and/or performance impairment. The recommended treatment for insomnia disorder is cognitive behavioural therapy for insomnia (CBT-I), a multicomponent intervention that differs from other CBT approaches in that it includes stimulus control therapy, sleep restriction therapy and other techniques specific to insomnia.[156–158] Interventions based on CBT-I (eg, sleep coaching) may also be effective at improving sleep health.[159]

While behavioural interventions remain first-line, pharmacological and nutraceutical approaches are sometimes considered when symptoms persist or when short-term support is needed. An international survey of sports psychiatrists showed that melatonin and magnesium were top choices of medications/supplements for insomnia in athletes.[63 160] Melatonin is the best-studied pharmacological sleep aid in athletes and does not appear to adversely impact performance.[161–165] It often improves sleep quality, but it may have limited utility for those with more severe symptoms.[166] Melatonin and magnesium are often purchased as non-prescription supplements, which may contain impurities—including prohibited substances—and unknown quantities of the labelled ingredients. Imidazopyridines (eg, zolpidem and zopiclone) and benzodiazepines are effective for insomnia disorder but frequently cause next-day performance impairments and may cause tolerance and dependence.[167] Trazodone, gabapentin, hydroxyzine, mirtazapine and many other medications are sometimes used as sleep aids but have not been studied in elite athletes.

For sleep apnoea, the recommended treatment is usually positive airway pressure therapy.[168] Alternatively, oral appliances are recommended for some individuals, especially those with mild to moderate symptoms.[169 170] For circadian rhythm disturbances, melatonin and timed light exposure are the treatments of choice.[118 119 171] For hypersomnia disorders, recommended treatments include stimulants as well as hypersomnia-specific agents (eg, modafinil, armodafinil, solriamfetol, pitolisant); in the case of narcolepsy, oxybate-based medications for sleep consolidation are also frequently used.[172–174] A TUE is required for all of these medications when prescribed for the management of hypersomnia disorders.

Major depressive disorder and depression symptoms

Major depressive disorder (MDD) is characterised by low mood and/or loss of interest or pleasure in activities. Burnout and overtraining syndrome (OTS) share overlapping features with depression, although each has distinct characteristics,[175] and burnout and OTS are not included in the Diagnostic and Statistical Manual of Mental Disorders, Fifth Edition, Text Revision (DSM-5-TR).[176]

Prevalence

Most research in elite athletes has examined depressive symptoms rather than a formal diagnosis of MDD and has been cross-sectional in nature, with few longitudinal studies conducted.[177–179] Depressive symptoms have been reported as highest in the mid-season compared with the off- and pre-season.[180] Reports have shown depressive symptoms within elite athletes range between 1% and 68%, depending on variability in sampling techniques, instrumentation and data collection methods,[1 181–184] with higher rates of symptoms reported when anonymity is preserved.[183] Overall, depressive symptom rates are similar between elite athletes and non-athletes.[185] Comparisons between current and retired athletes have shown a lower prevalence of symptoms in retired athletes.[186] Regarding the probable prevalence of overt depressive episodes, one meta-analysis provided a pooled prevalence of 16.2% of elite athletes experiencing at least moderate levels of depression, although there was much heterogeneity among the included studies.[183]

High levels of burnout have been reported in adolescent elite athletes, with 21.8% reporting medium levels and 9.1% severe levels.[187] Although overall prevalence results for different elite sports are lacking, cross-temporal meta-analytical results between 1997 and 2019 indicate that burnout symptoms for athletes have worsened with time.[188] Specifically, athletes have developed a reduced sense of athletic accomplishments and an increased sense of sport devaluation over time, which may contribute to higher levels of burnout. The prevalence of OTS is difficult to estimate as it depends on the definition of a syndrome that lacks clear diagnostic boundaries.[189] Nonetheless, prevalence of OTS in elite athletes has been reported to be between 7% and 18%.[190]

Diagnosis

To diagnose MDD, symptoms must be both persistent (minimum 2 weeks) and pervasive across activities.[176] Distinguishing among burnout, OTS and MDD can be challenging (table 5), but is important for providing the most effective treatment. Burnout in sports is commonly defined by physical and emotional exhaustion, reduced sense of accomplishment and a devaluation of the sport itself.[191 192] OTS is characterised by physical exhaustion, objectively impaired sporting performance and mood disturbances.[193] In MDD, the symptoms and impairments are pervasive, while symptoms of burnout and OTS are largely context specific (confined to sport).[175] The situation is further complicated as problematic low energy availability (LEA) and Relative Energy Deficiency in Sport (REDs) are also associated with depressive symptoms and fatigue,[194] although LEA and REDs are not independent risk factors for MDD in the absence of disordered eating (DE) or an eating disorder (ED).[195]

The ecological systems model can be used to stratify sport-specific risk factors for MDD and depressive symptoms.[7 10 196] Individual athlete risk factors include female sex, concussion or other injury, surgery, adverse life events, osteoarthritis, chronic pain, being under 25 years of age, enduring stress, burnout, perfectionism, competitive anxiety, poor sleep, negative attribution after failure, negative coping strategies, high athletic identity and sudden transition out of sport. Microsystem (interpersonal) factors include low social support, coach conflicts and

Table 5 Comparison of key features of burnout, overtraining syndrome and major depressive disorder

	Mood	Anhedonia	Motivation	Energy	Considerations
Burnout	Decreased, context specific	Present, context specific	Decreased, context specific	Decreased, context specific	Improves with rest and other environmental changes
OTS	Decreased, context specific	Present, context specific	May remain high	Decreased, pervasive	Improves with rest and other environmental changes
MDD	Decreased, pervasive	Present, pervasive	Decreased, pervasive	Decreased, pervasive	Worsens with rest

MDD, major depressive disorder; OTS, overtraining syndrome.

Br J Sports Med: first published as 10.1136/bjsports-2025-111514 on 11 August 2026. Downloaded from http://bjsm.bmj.com/ on August 12, 2026 at Universitetsbiblioteket i Tromsø . Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies.

family issues such as separation. Exosystem (athlete's sport) and macrosystem factors include those that pertain to the sporting institution such as physical settings, cultural norms, regulatory policies, individual versus team sports (higher rates of depression in individual sports), certain positions, levels of play and exposure to non-accidental violence. Athletes also experience a similar range of biological/genetic, social/environmental and psychological risk factors as their non-athletic counterparts.[1 177]

Impact

The consequences of depressive symptoms can be physical (eg, decreased performance, injury risk), cognitive (eg, loss of interest, diminished motivation), behavioural (eg, increased risk of leaving sport, difficult adhering to injury rehabilitation programmes), relational (eg, strained team and family relationships), financial (eg, loss of sponsorship) and psychological.[1 175 196] Athletes with a depressive disorder may be at increased risk of developing other mental health conditions including gambling disorder,[197] substance misuse,[198] exercise addiction, REDs or EDs.[194 199]

Management

Treatment for MDD typically involves psychotherapy, with or without medication.[177] CBT is suitable for many athletes due to its structured self-directed approach.[200] Medications are typically considered in moderate to severe depression.[1] In a 2024 international survey of practising sports psychiatrists, bupropion was a top choice for use in athletes with depression without an anxiety component, while selective serotonin reuptake inhibitors (SSRIs) (escitalopram, fluoxetine and sertraline, respectively) were the top choices for depression with an anxiety component.[63] Note that, while commonly prescribed in some countries, bupropion is not licensed as an antidepressant in some regions, potentially limiting its use. Additionally, athletes with OTS may exhibit impaired immune function and depressive symptoms, for which SSRIs may be less effective.[201–203]

Suicide

Prevalence

Studies on suicidal ideation in professional athletes show prevalence rates between 6.9% and 18%.[204] An analysis of deaths by suicide within the National Collegiate Athletic Association (NCAA) between 2002 and 2022 revealed increases for both males and females over time, with higher rates of suicide among Division I and II (larger college) athletes compared with Division III athletes.[205] Over this 20-year period, suicide rose to the second most common cause of death for collegiate athletes after accidents. Comparative studies suggest that elite athletes generally face a lower risk of death by suicide compared with non-athletes,[204] with identified risk factors including male sex,[206 207] non-white race[206 208] and depression.[209 210]

Management

Although elite athletes often avoid seeking help due to the stigma surrounding mental health, improving mental health literacy through access to information about symptoms and support services can be beneficial. Effective prevention and management of mental health symptoms, particularly MDD, are crucial.[204 205 211 212] Other prevention strategies include identifying risk factors, targeted support, management of social media, which may impact feelings of self-worth, and structured safety plans. Responsible media reporting practices are critical to reduce suicide contagion risk, as recommended by major health organisations.[213–215] Postvention strategies that provide support to individuals who have lost someone to suicide have been shown to be effective in non-sport and should be considered.[216]

Anxiety disorders and obsessive-compulsive and related disorders

Anxiety disorders include generalised anxiety disorder, panic disorder, agoraphobia, social anxiety disorder, specific phobia and separation anxiety disorder. Anxiety disorders are among the most common and most studied mental health disorders affecting elite athletes.[9] Among elite athletes (n=221) seeking mental health treatment at outpatient clinics in Sweden, 69% met the diagnostic criteria for an anxiety disorder.[9] In elite Para athletes, symptoms of mixed anxiety/depression range from 34%[95] to 76%[217] and may be severely exacerbated during the Paralympic games compared with pre- and post-competition periods.[218] Obsessive-compulsive and related disorders include obsessive-compulsive disorder (OCD), body dysmorphic disorder, hoarding disorder, trichotillomania and excoriation disorder, all of which have generally been less well studied in elite athletes.

Prevalence and diagnosis of specific disorders

Generalised anxiety symptoms and disorder (GAD): GAD is characterised by excessive and persistent worry about multiple events or activities. The GAD-7 is the most utilised measure of anxiety in elite sport, implemented in at least 40 studies to date. The percentage of elite athletes meeting 'moderate to severe' generalised anxiety based on the GAD-7 ranges from 1.2%[219] to 29.5%.[220] Consistent with the general population, female elite athletes report higher GAD-7 scores than males,[14 219–227] and gender-fluid or gender-nonconforming athletes report higher anxiety than female athletes.[228] Generalised anxiety in elite athletes is associated with comorbid depression,[226 229–234] DE,[226 233] REDs[194] and LEA,[235] post-traumatic stress disorder (PTSD),[236] attention-deficit/hyperactivity disorder (ADHD),[237] sleep disturbance[124 234] and psychological distress.[238 239] Sport-related factors associated with increased generalised anxiety include injury,[221 223 237] post-concussion symptoms,[240] higher training load[233] and sport burnout.[238] Conversely, more match experience,[230] satisfaction in sport[237] and team sports[223] are associated with lower anxiety. Personal factors associated with reduced anxiety in elite athletes include self-esteem,[233 239] coping skills,[239] basic psychological needs satisfaction,[238] mental toughness,[237] resilience[241] and older age.[230]

Panic disorder and agoraphobia: Panic disorder is characterised by intense and unexpected panic attacks and the anticipatory fear of experiencing another attack or losing control. Agoraphobia is characterised by a marked fear or anxiety about situations in which escape might be difficult or assistance unavailable should panic-like symptoms or other incapacitating or potentially embarrassing symptoms occur. The prevalence of panic disorder in elite athletes ranges from 2.8%[242] to 4.5%,[221] while agoraphobia ranges from 1.8%[242] to 7%.[243] Rates are higher in elite female athletes than elite male athletes.[221 242] Agoraphobia symptoms in a sample of male elite Para athletes were lower compared with non-athletic peers with a disability.[244]

Social anxiety disorder: Social anxiety disorder is characterised by the fear of interaction with and negative evaluation by others. The estimated prevalence of social anxiety disorder diagnosis or caseness in elite athletes ranges from 1.3%[242] to 14.7%.[221] Most studies report no sex differences,[221 242 245] with the exception of higher rates of clinically significant social anxiety symptoms in collegiate female athletes (37%) compared with male collegiate athletes (22%).[246] Social anxiety disorder

Br J Sports Med: first published as 10.1136/bjsports-2025-111514 on 11 August 2026. Downloaded from http://bjsm.bmj.com/ on August 12, 2026 at Universitetsbiblioteket i Tromsø . Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies.

in elite athletes may manifest as avoidance of team meetings or other group activities (eg, meals, training or rehabilitation sessions).[247]

Social physique anxiety (SPA): SPA is not an official DSM-5-TR diagnosis but generally refers to anxiety associated with being observed or judged on one's appearance. Several studies have focused on SPA in female cohorts of adolescent and collegiate athletes.[248–253] None have reported the estimated prevalence of SPA, but key correlates of increased SPA in adolescent female elite athletes include higher BMI[248 251] or overweight perceptions,[249] participation in individual aesthetic sports,[248] perfectionism,[252 254] body dissatisfaction,[252] bulimia nervosa,[252] DE attitudes[253] and female sex.[254 255] Elite Para athletes report lower SPA in comparison with non-Para athletes,[256] and elite athletes exhibit lower SPA compared with non-athletes.[257] SPA may be reduced by focused attention to sport-specific tasks.[253]

Competitive performance anxiety: Competitive performance anxiety is not an official DSM-5-TR diagnosis, but it is one of the most researched forms of anxiety in elite athletes. It refers to a state or trait-like response to competition or other stressful sport situations in which the athlete fears that they will be unable to perform as desired or to manage stress, coupled with anxious cognitive appraisals, behavioural responses and/or physiological arousal.[258] Competitive performance anxiety can overlap with other anxiety disorders (eg, GAD or social anxiety) and should be carefully distinguished, taking into account the pattern of symptom onset (usually proximal to competition), duration and intensity.

Obsessive-compulsive disorder (OCD): OCD is characterised by the experience of intrusive and distressing thoughts, images or impulses (obsessions) and/or repetitive behaviours or rituals (compulsions) that are usually performed to negate intrusive thoughts. The estimated prevalence of OCD in elite athletes varies from 1.7%[242] to 5–8%[9 259] through to 18.9% for combined 'OCD and related disorders' (eg, body dysmorphia, hoarding, trichotillomania).[260] Reported rates are higher among female athletes in some[242 260] but not all studies.[9] These rates are higher than the estimated 1.3% lifetime prevalence observed in community samples.[261] OCD *symptoms* are also commonly reported among elite athletes.[259] Idiosyncratic rituals and magical thinking, which are not pathological if they do not cause significant distress or dysfunction, are also common in this population.[262] Comprehensive assessment is necessary to differentiate OCD from more commonplace superstitions among elite athletes.[263]

Body and muscle dysmorphia: Body dysmorphic disorder refers to a preoccupation with perceived imperfections or flaws in appearance that cause distress and/or impairment. To date, only one study has examined rates of body dysmorphia in elite athletes, with 7.8% of female NCAA athletes reporting increased symptoms.[264] Research has instead focused on muscle dysmorphia (MD), a subtype of body dysmorphic disorder that refers to preoccupation with the idea that one's body build is too small or insufficiently muscular. Most studies have been conducted with male cohorts of weightlifters or body builders.[265 266] Rates of MD at or above threshold in weightlifters—particularly in relation to body size and symmetry—range from 6.3% to 8%.[267 268] Rates of MD at or above threshold in mixed-sport athletes are reportedly 13%.[269] In mixed sex samples, male athletes report higher body size and symmetry concerns than female athletes.[270] Higher rates of LEA are associated with higher MD symptoms in female elite and sub-elite athletes.[269] No epidemiological studies were found on other anxiety disorders or OCD-related disorders in elite athletes.

Impact

Anxiety disorders and anxiety sensitivity[271] (the fear of experiencing anxiety-related sensations including physiological, cognitive and social symptoms based on the belief that these symptoms have harmful consequences) are often associated with comorbid mental health symptoms or disorders in elite athletes.[226 229–234] Anxiety, OCD and related disorders can be debilitating and lead to impaired sport training or performance.[272–274] Competitive performance anxiety may also negatively impact sport performance and concomitantly be associated with other mental health symptoms or disorders such as depression or suicidal ideation.[272 273]

Management

There is a paucity of intervention studies for anxiety, OCD and related disorders in elite athletes relative to the number of epidemiological studies. Sport-specific contributing factors should be assessed and considered when tailoring a management plan to the individual athlete. Cognitive and behavioural therapies are regarded as first-line treatments for mild to moderate anxiety disorders,[42 48] with anxiolytic medications considered for more severe presentations or for athletes for whom psychotherapy has been ineffective.[247] According to a 2024 survey of sports psychiatrists, top medication choices across these disorders are the SSRIs escitalopram, fluoxetine and sertraline (in unspecified order).[63] Given the prevalence of anxiety, OCD and related symptoms in elite athletes and the potential for such conditions to become persistent and debilitative, it is critical to prioritise prevention and early detection and intervention.

Trauma- and stressor-related disorders

Exposure to traumatic or stressful event(s) is a precondition for a diagnosis of trauma- and stressor-related disorders. Among these disorders, athlete research has focused mostly on PTSD, which is characterised by at least 1 month of fear-based re-experiencing (eg, nightmares), avoidance and hyperarousal symptoms, and negative cognitions/mood.[176] Acute stress disorder is distinguished from PTSD as the duration of the former is 3 days to 1 month.[176] Adjustment disorder is the development of emotional or behavioural symptoms within 3 months of an identifiable stressor, resolving within 6 months after the stressor or its consequences have terminated.[176] Adjustment disorders are frequently co-morbid with other conditions. Prolonged grief disorder is a trauma-related disorder that can be diagnosed after 12 months of clinically significant distress or impairment following the death of a loved one.[176]

Prevalence

Accurate prevalence estimates of trauma- and stressor-related disorders in elite athletes are yet to be established across settings, but the numbers are likely significant. For example, stressor-related disorders (including acute stress reactions, adjustment disorder and PTSD) were recently observed in 25% of Swedish elite athletes and 72% of high performance coaches[9] among mental health treatment-seeking cohorts, although these numbers were derived from clinician diagnosis rather than structured interview. Prior research indicates 13.3% of professional athletes across sports who had been exposed to at least one traumatic event reported symptoms consistent with a PTSD diagnosis.[275] Limited research is available on the prevalence of PTSD symptoms in elite Para athletes. However, the risk may be as high or higher than in general populations of elite athletes, given

Br J Sports Med: first published as 10.1136/bjsports-2025-111514 on 11 August 2026. Downloaded from http://bjsm.bmj.com/ on August 12, 2026 at Universitetsbiblioteket i Tromsoe . Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies.

that a substantial proportion of elite Para athletes have acquired disabilities directly resulting from trauma.[53]

Diagnosis

The diagnosis of PTSD and other stressor-related disorders can be complicated given the complexities of expression in the elite athlete population.[53 276] Athletes' high tolerance for physical and psychological distress coupled with avoiding situations or mood states that the athlete perceives might interfere with training and competition can contribute to mental health symptoms being masked by perfectionism, compartmentalisation and dissociation.[277] Masking strategies may serve as temporarily adaptive coping techniques common to sport, enabling athletes to sustain engagement in both training and competition. However, they can simultaneously be maladaptive by creating a barrier to diagnosis of mental health disorders and by inhibiting long-term symptom improvement.[277]

For elite athletes, precipitating traumatic or stressful events may include organisational and institutional maltreatment,[276 278] interpersonal violence, concussion[279] and musculoskeletal injury.[280 281] Secondary or vicarious trauma can occur within sport teams following the serious injury or death of a teammate.[282] Elite athletes also may have experienced trauma or stressors outside of and/or prior to their sport participation. For example, 4.7% of male elite athletes and 15.1% of female elite athletes reported childhood sexual abuse. In female athletes this is associated with increased suicidal ideation.[283] Additionally, Para athletes are at a higher risk of experiencing interpersonal and sexual violence compared with their elite athlete peers without a disability.[278]

Cultural factors should be considered in assessments, with attention to 'idioms of distress' (ie, culturally influenced ways of coping).[276 284] For example, among African Americans in the USA, PTSD appears to be more prevalent yet less often diagnosed due to intersecting factors including stigma surrounding mental illness, reluctance to seek help beyond family networks, concerns about racial bias in health care and limited access to culturally competent care.[285] Consideration must be given to guilt, shame and stigma associated with traumatic experiences, as they complicate disclosure and therefore treatment. Lack of help seeking can be particularly common among athletes in environments that minimise pain, discourage emotional expression and/or promote hyper-masculine ideals.[286–288]

Acute and persistent symptoms, both emotional and physical, should be viewed through a trauma- and violence-informed lens.[289] This involves attention to individual developmental factors, cultural marginalisation, racialised trauma and transgenerational trauma.[276] Attending to the integrated effects from a biopsychosocial perspective can increase the potential for diagnostic accuracy, which then can inform treatment approaches.[276]

Early identification and treatment of athletes experiencing trauma-related symptoms including those of acute stress disorder may prevent progression to PTSD, while treatment of athletes already meeting criteria for PTSD may improve individual and sports performance outcomes.[276 289]

Impact

Symptoms can appear in the form of somatic or performance breakdowns/complaints.[277] Specifically, athletes may exhibit mood dysregulation, behavioural difficulties, problems concentrating and social withdrawal, all of which can lead to inconsistent performance. Trauma- and stressor-related disorders may predispose to injury as well. For example, athletes who undergo orthopaedic surgery may have had more cumulative (eg, childhood) trauma.[280] Without appropriate identification and intervention, trauma-related disorders can lead to ongoing impairment in social, occupational and sport function.[276]

Management

In general populations with PTSD, there is strong support for the use of structured and time-limited psychotherapy including prolonged exposure therapy, cognitive processing therapy and trauma-focused cognitive behavioural therapy,[290 291] with increasing support for eye movement desensitisation and reprocessing therapy.[276 292] Relaxation (eg, mindfulness) and somatic (eg, yoga) therapies may support symptom management.[276] Therapeutic alliance is critical for effective treatment given its impact on dropout, treatment adherence and outcomes.[276] First-line pharmacological treatment for PTSD includes SSRIs or SNRIs.[276] A 2024 international survey of sports psychiatrists showed that sertraline, escitalopram and fluoxetine were the top choices for medication management of PTSD in athletes.[63]

Additionally, team-based approaches with the athletes and staff directly or indirectly affected by secondary/vicarious trauma can provide effective intervention. Where indicated, a licensed mental health professional can provide psychological first aid for teams, reducing the stigma of seeking help in the case of vicarious trauma.[276] Promotion of a trauma- and violence-informed, non-judgmental, destigmatising culture may mitigate the complications of shared trauma as well as increase the likelihood of athletes receiving early intervention.[276 290]

Eating disorders and Relative Energy Deficiency in Sport (REDs)

Anorexia nervosa, bulimia nervosa, binge-eating disorder and other specified feeding or eating disorder are among the eating disorders (EDs) with defined criteria as outlined in the DSM-5-TR.[176] Disordered eating (DE) consists of abnormal eating behaviours not meeting criteria for an ED.[75] REDs is a syndrome marked by impaired physiological and psychological functioning caused by exposure to problematic (severe and prolonged) LEA.[293] REDs is caused by either under-fuelling or by over-expenditure of energy through high volumes of exercise. Specifically, under-fuelling can be either intentional (eg, via dieting or EDs) or unintentional (eg, via lack of knowledge or food insecurity). In both cases, there is a mismatch between energy intake through food and energy requirements of the body to maintain exercise and bodily function.[293]

Prevalence

Estimated ED prevalence is higher among elite female and male athletes (up to 42% and 33%, respectively)[294–298] compared with non-athletes.[74 295 297] In elite Para athletes, elevated Eating Disorder Examination Questionnaire (EDE-Q) scores have been reported in 36.7% and 26.4% of females and males, respectively.[299] The estimated prevalence of REDs is 22–80%[90 300–306] and 28–80% in elite female and male athletes, respectively, depending on the sport.[303 304 307–309]

Diagnosis

A multidisciplinary health care team (sports medicine physicians, dietitians and licensed mental health professionals) should monitor for signs of EDs and REDs during periodic health evaluations or when medically indicated.[41 293] Screening athletes for EDs poses unique challenges including under-reporting and difficulties in distinguishing between normative and pathological behaviours in an elite sport context.[297] Specifically, the

Br J Sports Med: first published as 10.1136/bjsports-2025-111514 on 11 August 2026. Downloaded from http://bjsm.bmj.com/ on August 12, 2026 at Universitetsbiblioteket i Tromsø . Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies.

sport-specific diet, training volume and the drive to optimise body composition/weight can obscure ED symptoms and result in diagnostic challenges.[41] The EDE-Q[310] is a sensitive and specific validated screening tool to identify elite athletes at risk for EDs.[298 311 312] However, clinical interviews are needed to diagnose an ED.[297] The 2023 International Olympic Committee's REDs Clinical Assessment Tool (IOC REDs CAT2) is a validated instrument that facilitates clinical diagnosis and risk stratification for REDs.[311]

Elite athletes experience sport-specific risks for EDs such as early sport specialisation, performance pressures, problematic body composition measurement practices, frequent weight regulation, injury and erroneous coaching behaviours.[75 295] Risk factors for REDs include high volumes of exercise, problematic practices to improve power-to-weight ratio, attempts to achieve excessive leanness and sport-specific physique demands.[293]

Impact

The impacts of EDs on health are complicated by medical comorbidities across the neuroendocrine, skeletal, nutritional, gastrointestinal, dental and reproductive systems.[313] The most prevalent mental health comorbidities for EDs in general are anxiety, mood, substance use and trauma- and stressor-related disorders.[313]

The impact of REDs on health varies across individuals and over time. Across sexes, the reproductive system is most often affected, resulting in menstrual disturbances in females, decreased libido in males and females and erectile dysfunction in males. Other body systems are also often affected.[293] Mental health impacts include anxiety, mood symptoms and fluctuations, sleep disturbances, exercise addiction and EDs/DE.[194]

EDs and REDs may not initially affect athletic performance but, over time, power, endurance, strength, motivation, cognitive skills, recovery, training responses and athlete availability may be reduced.[293] Elite athletes with an ED or REDs may require modifications to their training and competition until they achieve predetermined health status improvements.[293 311 314]

Management

A multidisciplinary team, which may include a sports medicine physician, dietitian and licensed mental health professional(s), should manage most elite athletes who present with DE, EDs or REDs.[314] Attention to common comorbidities such as depression and anxiety is important.[315] Suicide risk assessment is an integral component of treatment given the high suicide risk among individuals with EDs.[316] Ongoing monitoring is essential to ensure compliance and monitor clinical progress.[314] Addressing both the psychological and physiological aspects typically yields the best outcomes.[314] Treatment should be tailored to each athlete's specific presentation and comorbidities. CBT has been shown to be as effective in athletes as in non-athletes in treating EDs.[317]

Fluoxetine has been shown to reduce bingeing and purging, as well as improve mood and anxiety,[318] without affecting sports performance.[319] Lisdexamfetamine also may help binge eating disorder.[318] However, a TUE is required for the latter as lisdexamfetamine is a prohibited substance for those athletes subject to testing under the World Anti-Doping Code.[64]

Treatment of REDs should first attempt to reverse the problematic LEA via changes to energy intake and/or expenditure. If no clinical improvement occurs in 4–6 weeks, then the possibility of an underlying ED should be considered if it has not already been identified. Simultaneously, treatment should address specific presenting body system dysfunction(s) (eg, low bone mineral density and gastrointestinal dysfunction).[315 320]

To prevent EDs and REDs, it is important to proactively address body dissatisfaction, unhealthy dieting behaviours, exercise addiction and exposure to problematic LEA.[315] Research on ED prevention programmes for elite athletes is limited but promising.[321 322] Successful prevention efforts involve multiple sport partners (eg, athletes, coaches and sports administrators), interactive methods to discuss evidence-based information regarding growth and development, body composition, nutrition and training, and the creation of safe spaces for athletes to share experiences regarding body image and performance.[315 321 322] For example, an educational intervention programme developed for coaches of elite high school athletes showed that the intervention coaches had higher index scores on total knowledge, had a better understanding of how to address challenges related to weight management issues, and had improved ED recognition and management relative to control coaches at 9 months follow-up.[323]

Additionally, International Sports Federations can play an important role in prevention of EDs and REDs by developing health-focused sport regulations,[324] such as adjusting/increasing BMI criteria for certain sports where appropriate,[325] altering judging criteria in aesthetic sports and adhering to the IOC risk assessments for REDs and RTS guidelines.[311]

Attention-deficit/hyperactivity disorder

Prevalence

Attention-deficit/hyperactivity disorder (ADHD) is a brain developmental disorder characterised by persistent inattention and/or hyperactivity-impulsivity that interferes with functioning and/or development.[176] The prevalence of ADHD in elite athletes may be as high or higher than in the general population and may be even more prevalent in elite athletes in certain sports. ADHD prevalence in general populations ranges from 2.5% to 7.2% globally, but there are few studies on elite athletes.[326] The positive reinforcement of physical activity may attract children with ADHD to sports, which could lead to a higher prevalence.[327 328]

A systematic review exploring ADHD prevalence in youth and young adult athletes reported a prevalence range of 4.2–8.1% in athletes between the ages of 15–19 years,[329] and the percentage of athletes taking stimulant medications for ADHD in one university in the USA was reported as 7.1% during the 2009–2010 academic year.[330] Between 2017 and 2021, the percentage of Major League Baseball (MLB) players (from 40-man rosters) with TUEs for stimulant medications ranged from 7.5% to 8.6%. From 2022 to 2024, the percentage remained consistent at 5.1%. While the reason for the decrease in TUEs in 2022–2024 compared with 2017–2021 is unclear, these numbers likely underestimate the true prevalence in MLB players as these data reflect only those players who have received TUE approval for treatment with a stimulant medication.[331–334] The decrease in TUEs for stimulant medication among MLB players stands in contrast to global general population data that show increases in ADHD diagnosis and pharmacological treatments from 1979 to 2020.[335]

Diagnosis

Diagnosis of ADHD is based on a multifaceted approach that includes clinical interview, collateral information from multiple settings (including home, school and potentially work or sports) and a rigorous assessment completed by a licensed professional.

Br J Sports Med: first published as 10.1136/bjsports-2025-111514 on 11 August 2026. Downloaded from http://bjsm.bmj.com/ on August 12, 2026 at Universitetsbiblioteket i Tromsoe . Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies.

The assessment should include self-administered screening questionnaires completed by the individual (if an adult) as well as from sources associated with the athlete, in addition to clinician- or psychometrist-administered objective diagnostic interview instruments. Individuals meet diagnostic criteria for the inattentive or hyperactive/impulsive subtypes with demonstration of at least 6 months of at least 6 symptoms (at least 5 symptoms for age 17 and older) from the 'inattention' or 'hyperactivity and impulsivity' category, respectively, while combined type ADHD includes at least 6 symptoms (at least 5 symptoms for age 17 and older) from each of these categories.[176]

ADHD is a childhood-onset disorder but may be first diagnosed in adulthood for several reasons, including athletes leaving more structured environments (eg, homes, secondary schools) or leaving cultures or environments in which there is lack of support for ADHD being diagnosed. Additional testing may be needed to support the diagnosis and rule out other conditions.[336] Differential diagnosis and comorbidities require careful consideration, particularly in elite athletes. Common comorbidities that feature symptoms (eg, poor concentration, subjective or objective cognitive concerns) that overlap with ADHD symptoms include depression and other mood disorders; anxiety and related disorders; other neurodevelopmental disorders such as autism spectrum disorder, intellectual developmental disorder and specific learning disorders; and substance use disorders.[336] Conditions such as insufficient sleep or sleep apnoea can also create symptoms that mimic ADHD, which may result in misdiagnosis.

While ADHD is recognised globally, it may not be diagnosed as often in some regions due to different diagnostic thresholds, cultural stigma and/or less public discourse about the disorder. In addition, cultural interpretation of behaviours associated with ADHD can differ, leading to a different framework for understanding and addressing them.[337]

Impact

ADHD can have positive and negative impacts on sport performance. Behavioural deficits that can affect athletes with ADHD include inattention, motor incoordination, impulsivity, hyperactivity, psychological factors and sensory processing differences.[338] While impulsivity can confer quick reaction times and decision-making, it is also thought to increase the risk of substance misuse or substance use disorders.[330 339] Hyperfocus can filter distractions, and affected individuals can experience exhilaration and excitement in response to novel stimuli.[340] However, the impaired focus and concentration, oppositional behaviour, argumentativeness, frustration, low self-esteem and mood dysregulation found in ADHD may impair athletic performance.[330]

Management

ADHD management considerations typically include psychosocial interventions and medication management.[327 336 341] Psychosocial interventions include behavioural therapy, cognitive behavioural therapy, individual education plans, parent/caregiver teaching, training and support, and mental skills training and education for athletes, families and coaches.[328] Potential benefits of pharmacological treatment include core symptom reduction, as well as reduction in the risks of suicidal behaviours, substance misuse, transportation accidents and criminality.[342] Pharmacological management of ADHD is typically divided into stimulant and non-stimulant medications. Stimulants (methylphenidate and amphetamine compounds) are standard pharmacological treatments for ADHD in the general population and in elite athletes.[327 343 344] In 2024, sports psychiatrists reported that their top three medication choices for ADHD in athletes were all long-acting stimulant formulations (methylphenidate extended release osmotic-controlled release, lisdexamfetamine and mixed amphetamine salts extended release).[63] Non-stimulant medications that are sometimes used for ADHD include atomoxetine, viloxazine, guanfacine, clonidine and, especially if depression is comorbid, bupropion.[344]

Potential side effects of stimulants that can have a negative impact on safety and tolerability include increased heart rate and blood pressure, abdominal pain, headache, anorexia and weight loss, sleep impairment, jitteriness/anxiety and increased risk of heat illness.[345 346] Stimulants may be performance-enhancing and are prohibited by WADA *in competition*, and a TUE should be applied for and approved if stimulants are prescribed for medical treatment in most cohorts of elite athletes.[64] Stimulants are controlled substances that are regulated differently by nations across the globe, including being prohibited or illegal in some Asian countries. It is important for elite athletes to check regulations before travel and ensure the proper requirements are met, which may include travelling with a medical letter, prescription bottle and other documentation. Finally, treatment of comorbidities is important in the management plan for ADHD.

Bipolar and psychotic disorders

Bipolar and psychotic disorders are potentially serious conditions with a peak age of onset coincident with the peak age of sporting performance.[347] Bipolar disorder is characterised by mood episodes. These must include an episode of hypomania or mania where the central features are elevated mood, increased activity and energy and associated symptoms such as grandiosity, reduced need for sleep, distractibility and over-talkativeness. Depressive episodes are also typically present though not necessary to establish the diagnosis.[176] Psychotic disorders such as schizophrenia have symptoms of hallucinations and/or delusions as their central features. Additional symptoms of psychotic disorders may include disorganised speech or behaviour; negative symptoms (eg, lack of motivation); inappropriate (incongruous) affect; cognitive deficits; and problems with social cognition. There is often significant functional impairment in both groups of disorders.[176]

Prevalence

The unknown prevalence of both bipolar disorder and psychotic disorders in elite sport constitutes one of the most important gaps in the current literature,[178] and recent athlete surveys have not included measures to detect either.[348] The IOC SMHAT2 includes optional screening for both,[29] while the IOC mental health surveillance tool could aid identification.[40] The significant and enduring functional impairments of schizophrenia may mean that it is one of the few mental health disorders in elite sport with a lower prevalence than in the general population.[349]

Diagnosis

There are potential diagnostic difficulties in high-performance sport where bipolar- and psychotic disorder-type symptoms may result from another medical condition, or medication or substance use.[176] For example, medication or substance use may manifest with symptoms of mania, hypomania or psychosis. Examples include anabolic androgenic steroids,[350–353] cannabinoids and glucocorticoids[354] and mephenteremine (a sympathomimetic amine and vasopressor), which has recently attracted attention in the world of sport because of the potential adverse

Br J Sports Med: first published as 10.1136/bjsports-2025-111514 on 11 August 2026. Downloaded from http://bjsm.bmj.com/ on August 12, 2026 at Universitetsbiblioteket i Tromsoe . Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies.

effect of psychosis.[355–357] Additionally, sport-related behaviour may mask emerging mania—that is, both mania and sport can be expressed by high levels of activity and energy.[358]

Impact

Bipolar disorder mood episodes towards either pole (mania/hypomania or depression) may negatively impact sport,[349] although the condition is episodic and can be compatible with a sustained career in high-performance sport.[349] However, residual symptoms between episodes may persist,[359 360] emphasising the importance of comprehensive ongoing treatment.[361] In many cases, schizophrenia is associated with functional impairments that may eventually prove incompatible with continued sport participation at an elite level.[349] Importantly, many athletes are frequent travellers, and a majority of people with bipolar disorder are vulnerable to mood destabilisation when travelling across time zones,[361] especially via eastbound travel.[362]

Management

For bipolar disorder, evidence-based treatment guidance exists for both manic and depressive episodes, as well as for ongoing maintenance. Long-term pharmacotherapy is often needed to reduce the risk of relapse.[363–365] Sport-specific psychological and pharmacological treatments have not been evaluated, and guidance is based largely on expert opinion.[358 366 367] For example, sports psychiatrists from an international sample have reported that their top medication choices for bipolar disorder are as follows: aripiprazole for the manic/hypomanic and depressive phases and lamotrigine for the maintenance/prophylaxis phase.[63] That said, in practice, clinicians do not necessarily change medications across the course of the disorder once stability is achieved. These top choices are relatively less sedating and cause less weight gain than many other mood stabilising medications. Appropriate psychotherapy should be considered alongside medication.[364] The strongest evidence supports social rhythm therapy.[368]

For psychotic disorders, early identification and treatment are promoted, reflecting the importance of early interventions for improving the prognosis and long-term trajectory of the disorders.[369] Long-term medication is usually advised, given the chronic nature of the disorders and associated impairments.[370 371] There are no sport-specific guidelines,[349] but sports psychiatrists reported that their top medication choice is aripirazole.[63] For both bipolar and psychotic disorders, the benefits for physical health, mental health symptoms, functioning and social connections[372–379] mean that athletes should be encouraged to maintain regular exercise, which may include sports participation even if no longer competing at an elite level.[349 380] Finally, safety, stability and function may be significantly impaired in the acute phase of mania or when psychotic symptoms are present. These conditions can include loss of insight, which can sometimes necessitate involuntary exclusion of the athlete from training and/or competition.

Personality disorders

Personality disorders are enduring patterns of inner experiences and behaviours that deviate markedly from cultural norms, are pervasive and inflexible, have an onset in adolescence or early adulthood, and lead to distress or impairment. Conversely, personality traits are enduring patterns of relating to and thinking about the environment and oneself, can be harmful or adaptive, and do not cause the significant distress or impairment necessary for a personality disorder diagnosis.[176]

Prevalence

Personality disorders have received little study among elite athletes such that the prevalence is unknown. However, 'Cluster B' personality disorders and related traits are disproportionately represented within the existing publications on personality disorders in elite athletes.[381] This cluster is characterised by appearing dramatic, emotional or erratic, and three of its personality disorders have been studied in athletes: antisocial (a pattern of violation of the rights of others and impulsivity); borderline (a pattern of instability in relationships, self-image and affect and marked impulsivity, which often includes non-suicidal self-injury); and narcissistic (a pattern of grandiose need for admiration and lack of empathy).[176]

Diagnosis

No specific personality profile has been found to distinguish athletes from non-athletes.[381 382] Clinical interview—sometimes augmented with validated instruments—is the typical mode of diagnosis.[381 383] Importantly, there are cultural differences in personality styles that must be taken into account when considering personality disorder diagnoses in elite athletes.[176]

Many risk factors (eg, genetics, childhood trauma) for personality disorders and dysfunctional traits are not sport-specific. However, one sport-specific risk factor that has been described is that high degrees of success in elite sport, with resultant praise and attention, may lead to 'situational narcissism'.[384] The push for elite athletes to cultivate a social media presence may exacerbate this risk.[385 386] If 'situational' narcissistic traits become inflexible, persistent and dysfunctional, they constitute narcissistic personality disorder.[176] Pervasiveness across a broad range of settings makes narcissistic personality disorder distinctive from the public image, brand or persona that is sometimes created by an athlete in the context of sport and marketing (although these may represent narcissistic traits). An additional sport-specific risk factor for dysfunctional personality traits is anabolic androgenic steroid use, which has been associated with narcissistic personality traits.[387 388]

Impact

Personality disorders and traits may negatively impact sports performance through both individual and team effects. At the individual level, athletes with Cluster B personality disorders/traits may incur sanctions or legal penalties that affect participation.[389] There is an increased likelihood of doping/cheating in sport among athletes with antisocial and narcissistic personality traits.[390 391] Additionally, there may be an impact of personality traits on injury risk, as elite athletes with narcissistic traits[392] and higher sensation-seeking and boredom susceptibility traits[393] incur more injuries (possibly because of more risk-taking behaviours). Maladaptive perfectionism—the setting of high expectations for oneself and extreme self-criticism when expectations are not met—may be a trait of obsessive-compulsive personality disorder (a Cluster C personality disorder, which is characterised by a pattern of preoccupation with orderliness and control) and is a risk factor for eating disorders,[394–396] performance anxiety and depression in athletes.[381]

At the team level, athletes with narcissistic traits may engage in distracting, manipulative and selfish behaviours and contribute to dysfunctional team conflict, which may decrease group cohesion and performance and lead to drop out of teammates from sport.[397 398] Coaches' personality traits can also impact the team. For example, coaches with narcissistic traits are more likely to

Br J Sports Med: first published as 10.1136/bjsports-2025-111514 on 11 August 2026. Downloaded from http://bjsm.bmj.com/ on August 12, 2026 at Universitetsbiblioteket i Tromsø . Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies.

exhibit toxic controlling coaching styles, which can have a detrimental mental health impact on athletes.[399 400]

Management

Environmental supports may help athletes with personality disorders and dysfunctional traits. For example, coaches may play an important role, within appropriate boundaries,[381] in supporting athletes in regulation of negative emotions and related behavioural responses in the sport environment.[401] Management considerations typically include a focus on psychotherapy. Specifically, dialectical behavior therapy (DBT) is a first-line treatment for borderline personality disorder in general populations,[402] and the components of DBT have demonstrated efficacy in athletes.[403] Psychotherapy with athletes with personality disorders can be challenging and requires empathy, patience and persistence.[389 404] Comorbid mental health disorders are common and should be managed.[402] While primarily described in the setting of athlete eating disorders,[72 73] treatment contracts may be helpful for athletes with personality disorders in order to outline the criteria necessary for participation clearance and to ensure a shared understanding of how the clinical status of the athlete will be monitored by each member of the multidisciplinary team.

Substance misuse and use disorders

Substance misuse is the use of one or more substances in high dosages/potencies and/or in inappropriate situations that causes health or social problems.[405] A substance use disorder is a pattern of use of one or more substances that causes significant impairment and/or distress and is manifested by excessive or physically hazardous use, cravings or urges, use despite adverse consequences, failure to meet important obligations, reduced time spent on important activities and tolerance and/or withdrawal.[176] Substance misuse and use disorders pose significant challenges at elite sporting levels where the pressures to enhance performance, recover more quickly from overtraining and injuries or manage stress can entice both recreational substance misuse and use of performance-enhancing drugs (PEDs). Athletes may turn to both legal and illicit substances to gain a competitive edge, as a coping mechanism, or both, thereby potentially placing their physical and mental health at risk.

Prevalence

The most commonly used recreational substances among athletes include alcohol, tobacco/nicotine, cannabis, stimulants and prescription opioids.[227 406–417] Although the overall consumption rates of these and most other recreational substances are lower among athletes compared with the general population, certain substances such as alcohol (as binge drinking), oral tobacco/nicotine products and prescription opioids show higher prevalence in specific sports and athlete groups.[227 406–408 412–421] Specifically, rates of binge drinking (ie, ≥5 drinks per occasion for men and >4 for women) among collegiate and professional athletes from different sports and countries vary from 8% to 68%, with higher rates among men as well as across sexes in certain sports (eg, ice hockey, lacrosse, field hockey, rugby).[406–408 412 418–420 422 423] In a large sample of US collegiate athletes, 36% of men reported consuming >5 drinks while 33% of women reported >4 in typical drinking situations.[407]

Tobacco/nicotine product use (eg, oral tobacco/nicotine products, nicotine vaping) is also common among collegiate athletes and in certain professional sports (eg, skiing, ice hockey, soccer, baseball).[406 407 412 415 421] Annual use rates of different tobacco/nicotine products range from 4% to 22% among US collegiate athletes,[407] while 18% of male and 22% of female English professional footballers were current users of oral tobacco pouches or nicotine pouches.[424] Of elite international competitors from various sports, 23–29% tested positive for nicotine prior to competition.[406 407 415 421] Although research on substance use among elite Para athletes remains limited, recent evidence indicates comparatively low prevalence rates: alcohol (41.5%), nicotine (7.3%) and cannabis (8.5%).[425] Furthermore, another study reported that elite Para athletes exhibited lower rates of alcohol use than their non-Para athlete counterparts.[426]

Cannabis is typically still used less by elite athletes than non-athletes, but may be growing in popularity in some sports due to the changing legal landscapes and perceived norms that cannabis use is safe and may aid in managing anxiety and sleep difficulties.[406–412] For example, a large study of US collegiate athletes showed that annual use rates rose from 21.8% in 2013 to 25.9% in 2023.[407] A 2020 systematic review of youth, collegiate and professional athletes (n=46 202) from three continents revealed an overall annual prevalence rate of 23%, and a cross-sectional study of mixed competitive levels of endurance athletes (n=1193) reported current (past 2 weeks) use rates of 26%.[409 413] Cannabis use disorder in athletes has become increasingly common, especially among athletes who begin cannabis use at a younger age, who use daily and/or who consume high-potency cannabis.[410 411 414 418–420]

Beyond recreational substance use, PEDs such as anabolic agents, stimulants and selective androgen receptor modulators are misused by athletes aiming to improve muscle mass, endurance, recovery, appearance and overall performance.[418 420 427–431] These substances are among the most widely detected PEDs, with 2023 WADA testing data showing 0.57% positive test rates for Olympic sports and 1.88% for non-Olympic sports; 42% of positives were for anabolic agents, 16% for diuretics/masking agents and 15% for stimulants in 2022.[432 433] Carefully designed anonymous self-report surveys with validity checks have identified relatively high rates of self-reported PED use among elite international athletes (6.5–9.2%).[434] In contrast, one study of ultra-endurance athletes found positive results on blinded urine testing (eg, opioids 6.6%, diuretics 4.9%, beta-2 agonists 2.2%) that were not disclosed by any athletes on concurrent questionnaires.[435] Hormonal doping, including the use of growth hormone and erythropoiesis-stimulating agents, remains prevalent in sports that demand high physical output and endurance.[427] PED use is more common in recreational athletes than elite athletes.[430]

Risk factors for substance misuse and use disorders in athletes vary widely and include the type of sport (higher in team and collision sports), cultural norms (eg, partying lifestyle) within the sport, sex (males more than females), early age of first use and/or heavy use at an early age, mental health comorbidities (eg, anxiety, ADHD, body and muscle dysmorphia), behavioural addictions, interpersonal violence/adverse life events and the availability and legality of substances within a given country.[418–420 422 436] Substance misuse is generally more commonly encountered in elite athletes than are substance use disorders.[418–420 422] There is limited research on the prevalence rates of most substance use disorders in elite international athletes.[406 410 418 419 422 423 437]

Diagnosis

The routine use of validated recreational substance screening instruments (eg, Alcohol/Cannabis Use Disorders Identification Tests; AUDIT and CUDIT), negative consequences questionnaires

Br J Sports Med: first published as 10.1136/bjsports-2025-111514 on 11 August 2026. Downloaded from http://bjsm.bmj.com/ on August 12, 2026 at Universitetsbiblioteket i Tromsoe . Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies.

(eg, Brief Marijuana Consequences Questionnaire, Brief Adult Alcohol Consequences Questionnaire) and urine testing can lead to early detection and intervention and prevent progression to substance use disorders.[419] The evolution of methods of PED use has led to the development of sophisticated detection methods, such as the Athlete Biological Passport, aimed at monitoring changes in athletes' biomarkers to detect substances used for performance enhancement.[427 429 430]

Impact

Substance misuse and use disorders among athletes can lead to detrimental physical, mental and performance effects.[406 410 414 416–419 423–425 427–431 438 439] For example, athletes who binge drink, use cannabis or tobacco/nicotine products daily and/or who use alcohol and cannabis simultaneously have more adverse physical (eg, injuries, accidents, poor recovery) and mental (eg, sleep concerns, anxiety, depression, irritability, aggression) consequences than athletes who do not.[406 418–421]

Regarding performance, alcohol misuse can impair cognitive and motor function, reduce reaction time and negatively affect coordination and endurance.[420] Regular oral tobacco/nicotine use causes toxic physical (eg, gum irritation, nausea, dizziness, loss of appetite) and mental (eg, insomnia, mood instability, anxiety) effects that can impair quality of life and performance.[415 418–421 424] Continued use can lead to dependence/withdrawal that can produce irritability, cravings, low energy, restlessness and impaired concentration, all of which can impair performance.[415 418–421 424] The evidence on the impact of cannabis on performance is mixed.[406 408–411] Some studies have found a negative impact on physical performance, while others have found no effect.[406 408–411] Despite the perception that cannabis may aid in relaxation, sleep or recovery, there is little evidence to support its efficacy in either enhancing recovery after strenuous activity or improving sleep patterns.[406 409–411] Cannabis misuse can lower motivation, impair memory and learning, diminish socialisation and reduce academic potential.[418–420 439]

PEDs, especially anabolic agents and stimulants, increase strength, endurance, power and muscle recovery, but their use tarnishes the integrity of sport.[420 428–431] Anabolic agents and their analogues are associated with low sperm count, irregular menses and increased blood pressure/cholesterol and in higher dosages can cause cardiovascular disease, hepatotoxicity, psychosis and hormonal dysfunction.[420 429 430] Erythropoiesis-enhancing agents may lead to cardiovascular disease, polycythaemia and stroke.[420 427]

Management

Substance misuse and use disorders among athletes require specific prevention and treatment approaches. Given the unique pressures faced by this population, interventions must address not only the physical aspects of substance misuse but also the psychosocial factors contributing to substance use. The development of targeted prevention strategies such as routine screening, alongside comprehensive education on the risks associated with substance use, is crucial for protecting the health and careers of athletes.[28 406 418 419 440] Athlete education is particularly warranted regarding the use of dietary supplements, as studies indicate that up to 28% may contain undeclared prohibited ingredients, thereby posing a significant risk for unintentional doping.[441] Stigma and threats of sanctions deter self-referrals, and comorbid mental health symptoms and disorders complicate diagnosis and treatment.[418 419 440]

For substance misuse, brief motivational interventions that are individualised, use contingency management strategies and involve collaterals such as coaches or family members can shift elite athletes' attitudes and beliefs about substance misuse, reduce frequency of use and prevent progression to substance use disorders.[419] For substance use disorders, withdrawal symptoms and cravings should be addressed with both medication and behavioural strategies.[419 440 442] Higher levels of care (eg, intensive outpatient, residential) may be needed for athletes who cannot stop using with brief office-based interventions.[440 442] Addressing PED use in elite athletes requires a multifaceted approach that combines strict regulatory enforcement, biological passports, urine/blood testing, effective education and targeted intervention strategies tailored to the unique environment of elite sports.[420 430 440]

Gambling disorder and other behavioural addictions

Gambling disorder

Prevalence

The international landscape of gambling has substantially changed in the last decade. There are more land-based casinos, and online gambling via smartphone or laptop has emerged as a primary way to gamble. Recent meta-analyses have examined problem gambling among elite athletes.[443–445] These reviews suggest that, while most studies show a positive relationship between gambling/problem gambling and athletic involvement, there are some (although limited) studies that suggest no increased risk of gambling and problem gambling among athletes. Temporal differences may have resulted in prevalence variations among studies. Specifically, legalised sports gambling has only recently become readily available in the USA and, since then, prop bets (wagers on specific events within a sporting event that do not necessarily depend on the final outcome) have become commonplace.[446 447] The most recent study among NCAA collegiate athletes shows that almost 11% of male collegiate athletes reported gambling on sports at least monthly, with wagering via online sports betting showing the greatest increases. Overall, NCAA data suggest that 52% of male collegiate athletes and 36% of female collegiate athletes report gambling (of any type) in the previous 12 months,[446] with approximately 2% of male collegiate athletes and a smaller percentage of female collegiate athletes reporting some level of problem gambling.[446] This compares with global estimates for the general population of 0.12–5.8%.[448] Recent Swedish[449] and Spanish[450] studies have similarly reported a higher prevalence of gambling disorders among young male athletes relative to men in the general population.

Elite athletes may be at particular risk for a gambling disorder given their over-representation of young men, desire for competition and challenges, high levels of sensation seeking and impulsivity and risk-taking behaviours.[1 21 451–453] Their pressures to perform, travel commitments and being away from family are all factors impacting their gambling. Further, their overall sports exposure, perceived knowledge of sports, availability of prop bets and ability to make wagers in the privacy of their home may also make them prone to sports wagering.[446]

Diagnosis

Gambling disorder has been regarded as a 'hidden disorder' and thus can be difficult to detect unless athletes are specifically asked about its presence.[454] The diagnosis shares features with substance use disorders and requires at least 12 months

Br J Sports Med: first published as 10.1136/bjsports-2025-111514 on 11 August 2026. Downloaded from http://bjsm.bmj.com/ on August 12, 2026 at Universitetsbiblioteket i Tromsoe . Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies.

of persistent and recurrent problematic gambling with resultant negative consequences, including gambling with increasing amounts of money, preoccupation with gambling, 'chasing' one's losses and jeopardising of relationships, career or education.[176 455]

Impact

The consequences for elite athletes who develop problematic gambling include mental health disorders, financial difficulties, poor school and athletic performance and sometimes criminal behaviours.[451 456] Additionally, athletes and their coaches and family members have increasingly been harassed and threatened with bodily harm by gamblers via social media for not performing at a certain level.[446 457]

Other behavioural addictions

Other behavioural addictions may be of relevance to elite athletes, although they are not official DSM-5-TR diagnoses. Like gambling disorder, these may be hidden disorders that share features with substance use disorders and include exercise addiction and internet gaming disorder.[1]

Prevalence rates of exercise addiction in athletes have been reported as 3–14.2%,[458] with reports of higher risk at higher competitive levels.[459 460] While the prevalence may be higher among male athletes, the difference in risk between male and female athletes diminishes at higher competition levels.[460] It can be difficult to distinguish exercise addiction from normative elite athlete behaviour.[1 461] The diagnosis should be suspected in the presence of symptoms such as increased exercise volume over time for the purpose of avoiding negative feelings or because more is needed to satisfy 'cravings' for it; inability to reduce exercise volume even in the face of the desire to do so or negative consequences such as injury; and jeopardising of previously valued life activities and relationships because of excessive exercise.[462 463] The clinician should differentiate primary exercise addiction, where exercise is the main objective, from secondary exercise addiction, where the excessive exercise is a symptom of an eating disorder.[464] Impacts of exercise addiction may include injury,[199] worse sports performance,[465] depression[199 466] and REDs.[467]

Internet gaming disorder in elite athletes has been minimally studied, but prevalence rates in Swedish elite athletes have been reported to be 4% in males and 1% in females, approximating the general population prevalence rates.[468] Problem gaming and problem gambling may co-occur in elite athletes,[468] and time-consuming travel in this population may stimulate excessive engagement in both types of online behaviour.[468] Proposed research criteria for internet gaming disorder include at least 12 months of preoccupation with gaming and several associated symptoms.[176]

Management of gambling disorder and other behavioural addictions

Treatment for gambling disorder and other behavioural addictions is rooted in psychosocial approaches,[469] especially CBT and motivational interviewing.[199 470–472] Unlike most other addictive disorders, abstinence may not be the goal, especially for exercise addiction.[473] Currently, no medications are approved to treat behavioural addictions. Antidepressants (especially SSRIs) and opioid antagonists (nalmefene and naltrexone) have been studied, but evidence for efficacy is limited.[474]

For problem gambling specifically, prevention efforts are a major strategy.[446 454 472] At a systemic level, organisations such as universities can offer programmes to educate athletes and their entourage about the potential mental health harms associated with a gambling disorder.[475] Such programmes can enhance communication between athletes, coaches and athletic directors, increase knowledge about gambling statutes and regulations for athletes, and raise awareness about signs of problem/disordered gambling. With the growing number of athletes being threatened by gamblers, athletes should be encouraged to report threats, attempts at bribery or physical harm to their respective authorities.[446]

Neuropsychiatric aspects of sport-related concussion

Prevalence

The sport-related concussion literature commonly notes that 42 million individuals worldwide sustain concussions every year and that up to one-third of concussions are related to sports or recreational activities, especially in adolescents and young adults.[476] However, the incidence and prevalence of sport-related concussion are difficult to estimate because of differing definitions of concussion, differing methodologies in injury surveillance systems and under-reporting.[477] Concussion is especially common in contact-collision sports and likely accounts for 9–13% of high school sport injuries.[477] The concussion rate in collegiate athletes is 4.13 per 10 000 athlete-exposures (AEs), with highest rates observed in men's ice hockey (7.35 per 10 000 AEs) and women's soccer (7.15 per 10 000 AEs).[478]

Diagnosis

The sixth International Conference on Concussion in Sport[479] defines sport-related concussion as a traumatic brain injury caused by a direct blow to the head, neck or body resulting in an impulsive force being transmitted to the brain that occurs in sports and exercise-related activities. This initiates a neurotransmitter and metabolic cascade, with possible axonal injury, blood flow change and inflammation affecting the brain. Symptoms and signs may present immediately or evolve over minutes or hours, and commonly resolve within days but may be prolonged. Standard imaging studies are normal. Sport-related concussion results in a range of clinical symptoms and signs that may or may not involve loss of consciousness, and that cannot be explained solely by (but may occur concomitantly with) drug, alcohol or medication use, other injuries (eg, cervical injuries, peripheral vestibular dysfunction) or other comorbidities (such as psychological factors or coexisting medical conditions).

The Sport Concussion Assessment Tool (SCAT6)[480] is useful for assessing and diagnosing suspected concussion up to 3 days following injury. Of 22 symptoms in the Symptom Evaluation of SCAT6, 13 are related to neurobehavioural dysfunction. However, such immediate post-concussion symptoms do not predict mental health symptoms and disorders following sport-related concussion. Concussion recovery within 4 weeks of injury is common,[479 481] although 10% of athletes have ongoing symptoms that preclude RTS 6 months post injury.[482] The term 'persisting symptoms' is used to characterise symptoms that remain >4 weeks following the initial injury; such symptoms may be pre-existing, concussion-related, or both. Importantly, persisting symptoms do not necessarily reflect ongoing brain injury but rather are related to maladaptive recovery.[479 481] Common persisting symptoms include headache and pain, dizziness and imbalance, poor concentration and cognition, oculomotor dysfunction, sleep dysfunction and fatigue, anxiety and depression and dysautonomia symptoms.[479 483 484]

Pre-concussion psychological factors such as meanness, aggression and psychoticism as well as depression and anxiety

Br J Sports Med: first published as 10.1136/bjsports-2025-111514 on 11 August 2026. Downloaded from http://bjsm.bmj.com/ on August 12, 2026 at Universitetsbiblioteket i Tromsoe . Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies.

are associated with an increased risk of developing sport-related concussion.[485 486] Repetitive head impact exposure may increase the risk for developing sport-related concussion.[487] Pre-existing mental health symptoms or disorders, family history of mental health disorders and significant life stressors are associated with an increased risk of developing persisting symptoms following concussion, although studies have varying methodological approaches, and mental health disorders are not studied as a primary outcome.[488] Pre-existing depressive symptoms are associated with a higher number of depressive symptoms/persisting symptoms following concussion.[489] A 12-month observational prospective cohort study of professional rugby players found an increased association of both feeling distressed and adverse alcohol use (independent of feeling distressed) following concussion, although all findings were from self-reports.[490] Sleep dysfunction following concussion correlates with worsened neurocognitive performance, persisting symptoms and impaired attention/processing speed.[491–493] Pre-injury sleep dysfunction may kindle persisting symptoms.[494]

Impact

There are emerging high-quality studies that address long-term mental health outcomes in athletes who have been exposed to frequent contact-collision activities. These studies presume that athletes who train and compete in contact-collision sports have suffered from concussion and have been exposed to repetitive head impacts, but none provide specific measurements for concussion or repetitive head impact. American youth football players have no increased risk for depression or premature death from suicide during early and late adulthood,[495–499] and former American professional football players who played five or more seasons have a decreased risk of suicide compared with the general population.[500] Former professional soccer players are not at increased risk of death from suicide[501 502] or other mental health disorders.[501]

Management

Sub-symptom threshold aerobic exercise treatment within 2–10 days after sport-related concussion is effective for reducing the incidence of persisting symptoms after concussion and is also effective for the management of persisting symptoms, including persisting mental health symptoms.[479 503–507] Cognitive rehabilitation and neurocognitive training are effective interventions for mental health and well-being and persisting symptom impairments post-concussion.[508] Targeted sleep intervention strategies should be used for sleep dysfunction,[492] although there are no systematic studies for post-concussion sleep disorders. There are no approved pharmacological interventions for mental health symptoms and disorders following concussion, and treatment should be guided by clinical indications.

Considerations for the elite Para athlete

The concussion experience is likely to vary for the elite Para athlete due to the interaction of their primary impairment combined with the pathophysiology of concussion,[509] although there is a paucity of literature on sport-related concussion in this population.[510] Commonly used tools (eg, SCAT6) are not validated for Para athletes, although a Para SCAT-WC (for wheelchair athletes) is forthcoming. Other important considerations for the elite Para athlete include[509]:

- Elite Para athletes have a history of under-reporting concussion.
- Baseline testing may have greater value for the elite Para athlete. A medical professional who knows the athlete well should be involved in their pre- and post-concussion assessment.
- The mechanism for testing sub-symptom threshold aerobic exercise may require modification.
- When considering return-to-learn, previously existing disability accommodations should remain in place.

Stressors and environmental factors that impact elite athlete mental health

Creating an environment that supports mental well-being and resilience in elite athletes

Core elements of a supportive environment

Elite athletes function within an ecosystem that includes the athlete; the microsystem of interpersonal relationships; the exosystem of the athlete's sport; and the macrosystem of larger sport and societal influences.[18] A supportive ecosystem allows for mental health symptoms and disorders to be proactively managed and for athletes to flourish (ie, to experience positive self-regard, personal and professional growth and meaningful social connections).[14 511] On the other hand, an unsupportive ecosystem or inappropriate practices or procedures within the ecosystem (eg, an environment with low psychological safety, low social support, inappropriate body composition assessment practices, etc) may lead athletes to languish, even if they have no history of mental health symptoms or disorders.[14 426 511 512] Toxic leadership—arising from coaches, sports directors, managers, executives or anyone else acting in a leadership capacity within sport—is abusive and potentially destructive. It can generate serious and enduring effects on individuals, families, organisations, communities and societies[513] and can have a negative impact on athletes' perception of team culture, autonomy, coaches' leadership skills and coaches' concern for athletes' well-being.[514] Conversely, a supportive ecosystem can support athlete thriving, which includes a trusting sense of belonging coupled with high athlete motivation and self-belief.[515] Organisational practices including the allocation of adequate resources, leadership prioritisation and a commitment to ongoing quality improvement are important components of creating a climate that is supportive of mental health.[516 517] Finally, the media also has a responsibility to report in ways that promote mental health awareness and reduce stigma in society.[518]

Primary role of coaches and parents/caregivers

As key members of the athlete ecosystem, coaches and parents/caregivers merit particular mention. Coaches are often the primary point of contact with athletes and thus can considerably influence an athlete's attitude to mental health care seeking while also providing a supportive environment that promotes and supports well-being.[514 519] Mental health care seeking should be normalised in a manner similar to health care seeking for injuries and other medical illnesses.[519 520] Coaches should attend to stressors that may predispose athletes to mental health symptoms and disorders, including training overload and insufficient recovery,[521] injury,[521 522] burnout,[523] developmentally inappropriate activities[516 524] and REDs.[293] Coaches can identify warning signals for mental health symptoms or disorders,[28] facilitate appropriate referrals[519] and help ensure and support athlete engagement with mental health management.[519]

Whenever possible, coaches and parents/caregivers should align on mental health care seeking and fostering wellness. Parents and caregivers can positively influence athlete development by

Br J Sports Med: first published as 10.1136/bjsports-2025-111514 on 11 August 2026. Downloaded from http://bjsm.bmj.com/ on August 12, 2026 at Universitetsbiblioteket i Tromsø . Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies.

investing time, financial resources and emotional support that fosters a holistic pathway. Such support can promote intrinsic motivation, self-efficacy and adaptive skills.[525 526] Negative influences from parents and caregivers can include an overemphasis on winning and setting inappropriately high expectations, which can contribute to anxiety, decreased perceptions of competence and a decrease in sport enjoyment.[527] As with all other members of the sport ecosystem, coaches and parents/caregivers should have contextually relevant knowledge and beliefs about mental health that aid in the recognition, management and prevention of mental health symptoms and disorders.[528–530] Encouraging the involvement of coaches, parents/caregivers and other community resources as appropriate in mental health initiatives can help create a supportive sport ecosystem.

Interpersonal violence

Interpersonal violence (IV) represents an aspect of some unsupportive sport ecosystems. Also known as harassment and abuse, it is defined by the WHO as 'the intentional use of physical force or power against other persons by an individual or small group of individuals. It can occur online, be perpetrated by different actors and take different forms'.[531] IV in the sporting context can be expressed through four forms: psychological violence, physical violence, sexual violence and neglect.[532] Other forms of abuse towards athletes include hazing, bullying and grooming. IV can be manifested on or off the field of play.[532] Perpetrators of IV can be peer athletes or other members of the athlete entourage, often based on interpersonal power differentials.[533] The potential outcomes of IV in sport can be psychological, behavioural, physical and/or material damage. The specific mental health outcomes that have been described include depression,[534 535] anxiety,[535 536] PTSD,[537 538] suicidality,[283] body dissatisfaction,[250] EDs/DE[539] and risk-taking behaviours such as substance misuse and risky sexual behaviour.[540]

Members of the elite athlete health and performance team should have competence in trauma- and violence-informed care to facilitate IV prevention, identification, assessment and recovery.[541] Clinicians should screen for a history of IV within and outside the sport context. It is the responsibility of all within the sport ecosystem to work synergistically to strive for a safe sport environment to protect athletes from IV and its negative mental health impacts. To strive for a safe sport environment, sport organisations should implement safeguarding practices (eg, safeguarding policies and procedures, reporting mechanisms and athlete support) and improve awareness of IV and its impacts through educational initiatives. Additionally, members of the athlete entourage should be equipped with trauma- and violence-informed skills to build trust and prevent re-traumatisation. A healthy sport culture is grounded in mutual respect and values-based principles, with athletes meaningfully empowered and involved at all levels.[541]

Cultural considerations

All factors within the ecological systems model can impact the mental health needs of elite athletes and should be recognised and considered when working to address their mental health needs.[542 543] As one of these factors, culture can be understood as a set of shared beliefs, feelings, knowledge, norms and objects that influence people to give meaning to things, experiences and behaviours.[544] Cultures are intersectional, and all athletes are impacted by multiple cultures, including the culture of their sport, cultures related to their identity characteristics (eg, gender, race and ethnicity, sexual orientation) and the cultures of communities where they reside or with which they engage. These cultures shape the conditions that contribute to mental health symptoms and disorders, how mental health symptoms may be understood and experienced, and health care utilisation by elite athletes.[545 546] For example, gendered beliefs and norms may shape help-seeking behaviours.[547 548]

It is essential that mental health care be patient-centred, with cultural responsiveness constituting a critical component of such care. This means recognising that an athlete's culture and cultural exposures influence how distress is experienced, expressed and managed. Patient-centred culturally responsive care should be grounded in cultural humility.[549] Practising with cultural humility means that clinicians try to understand the lived experiences and cultural perspective of their patients, and dedicate themselves to lifelong learning in order to provide culturally responsive care. Such a framework demonstrates clinicians are willing to reflect on their values, beliefs and various biases, and adapt their practices accordingly. Understanding, respecting and integrating an athlete's cultural background into care helps build trust and is essential for both accurate assessment and effective management.

Central to cultural humility is an awareness of structural inequalities that shape social determinants of mental health.[550] Structural inequalities are systemic disadvantages that are experienced by various social groups, be it on grounds of race, gender, gender expression, sexual orientation, class, religion, immigration/migration status or geography.[551] Social determinants of mental health are the conditions within which people live and work and the circumstances that affect their mental health in daily living.[550–553] Social determinants can include education, income security, employment conditions, housing, food security, transportation services, social environment, conflict/war and access to health care services.[552–554]

Structural inequalities and social determinants of mental health are experienced unevenly across populations; consequently, some groups of elite athletes may experience a higher prevalence of mental health symptoms and disorders than their peers. For example, research has shown that athletes who are women, non-white or do not identify as heterosexual experience higher levels of psychological distress, often due to greater challenges with core social determinants such as housing, income and food insecurity.[553] Further, transgender athletes face higher rates of structural inequalities and discrimination compared with their cisgender peers, contributing to chronic stress and attendant higher rates of mental health symptoms and disorders.[555–557] Athletes who experience immigration challenges or live as refugees experience disparities in structural and social conditions and have high levels of mental health symptoms and disorders, which can negatively impact functioning across domains.[558 559]

The potential impact of structural inequalities and social determinants of mental health should be considered when addressing mental health needs in elite athletes. Although considerable attention has been paid to the cultural training of individual clinicians and staff within sport,[550] there is also a growing recognition of the importance of organisational practices in addressing these issues.[553] This could include organisational practices that aim to provide all mental health services in an equitable and inclusive manner.[551 553 560 561] For example, the NCAA sets out provisions for universal screening in which all athletes must be screened for mental health symptoms and disorders and offered referral pathways as needed to qualified mental health providers.[561] Organisational practices could also include taking a collaborative inter-organisation approach to addressing structural inequalities.[562] An example would be encouraging collaboration between

Br J Sports Med: first published as 10.1136/bjsports-2025-111514 on 11 August 2026. Downloaded from http://bjsm.bmj.com/ on August 12, 2026 at Universitetsbiblioteket i Tromsø . Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies.

health care providers and individuals who can support an athlete in connection to community-based resources to address stressors such as housing insecurity or relationship violence.

Barriers to mental health care seeking

Elite athletes may face significant barriers while seeking and sustaining mental health care, frequently rooted in the perceived norms of athletes not revealing their vulnerability, especially regarding mental health. Cultural perceptions that equate mental health symptoms and disorders with weakness further this stigma.[563] For example, members of certain racial and ethnic groups may prefer not to seek mental health support because of stigma that has been augmented through historical discrimination and mistreatment within medical settings.[563] In some sports, hypermasculinity and the cultural expectancy to 'tough it out' discourage athletes, particularly men, from pursuing help.[564] Moreover, elite athletes may fear that seeking help will result in undesirable judgments from peers, coaches or the public, as well as potential exclusion from future competitions.[563] Finally, elite athletes may have concerns about privacy that may lead to fear that disclosing one's problems could affect professional opportunities (eg, via deselection or loss of sponsorship contracts).[565]

In addition to stigma and cultural expectations, logistical challenges can hinder athletes' access to mental health care.[566 567] Elite athletes have rigorous training and competition schedules that often do not prioritise self-care and management of mental health symptoms.[566] This pressure to adhere to a particular periodisation schedule does not take into account mental well-being and recovery.[567] Furthermore, financial constrictions, geographic isolation of sports facilities and other logistical problems can prevent athletes from accessing mental health services.[568 569] Additional regulatory barriers, such as restrictions that prevent clinicians from practising across state, provincial or national lines, also hinder access to care; this is an important issue given the frequency of athlete travel. Moreover, much of the evidence base on which mental health recommendations are made comes from high-income countries and may not translate well to lower-income settings. For example, in some locations, psychiatric hospitals may lack basic infrastructure and resources, thereby limiting opportunities for help-seeking even when services are theoretically available.[570] Some countries have implemented systems to overcome logistical barriers. For instance, Australia has developed a free mental health network for elite athletes, and Sweden operates a no-cost specialised clinic for elite athletes.[571 572]

Improving mental health literacy and prioritising wellness and mental health care within the sport ecosystem and periodisation schedule could help to mitigate these barriers.[7 573] Mental health literacy efforts should address how individuals: (1) understand and recognise mental health symptoms and disorders, risk factors and causes, and treatment options; (2) work through personal and social stigma associated with mental health symptoms and disorders; and (3) find and use mental health services to address mental health symptoms and disorders.[542 574] Mental health literacy strategies must be constructed with knowledge of a given athlete population and their specific sociocultural and sporting contexts that may impact availability, accessibility, quality and fit of mental health services and help-seeking behaviours of athletes.[542 543] Different groups of elite athletes may have differing needs and assets when it comes to mental health literacy. For example, within elite sport, research has shown that women and LGBTQIA+ individuals have higher levels of knowledge of mental health symptoms, hold fewer stigmatising views and are more willing to seek support.[575] Within youth sport, it is recommended to design mental health literacy strategies where learning is consistent with social, psychological, mental and physical developmental readiness.[542 576] Organizational health literacy strategies, where organisations help ensure that information is accessible and understandable and that services are available, help ensure that individually-oriented mental health literacy strategies are effective.[542] Careful attention must be paid to ensure that mental health literacy strategies are designed to improve mental health knowledge, attitudes and help-seeking behaviours in an equitable, culturally responsive and effective manner.[542 575 577–580]

Social media and mental health

Social media use

Social media use has rapidly escalated in recent years, with up to 95% of 13–27-year-olds in the USA using these digital platforms.[581] The preferred social media platforms among young adults can evolve quickly. Concerns have been raised regarding the use of social media by elite athletes and their interactions with traditional media,[582 583] although the scientific evidence in this area remains limited.

Mental health impacts of social media use

Several cross-sectional studies have examined social media use (including overall screen time, social media engagement during daily activities and perceived interference from social media use) and mental health outcomes in collegiate athlete cohorts. Findings demonstrate associations between social media use and poorer general mental health and self-esteem,[584 585] heightened anxiety and diminished subjective well-being,[585 586] reduced sleep quality[587 588] and increased body image concerns.[589] Higher levels of social media use, particularly during daily activities, have been associated with poorer mental health outcomes.[584 585 587] One study found that females have both greater social media use (specifically TikTok) and associated higher levels of anxiety.[584] In adolescent athletes, a statistically significant relationship was found between time spent on social media and symptoms of addiction to digital media, as well as increased negative emotions, disordered eating symptoms and decreased quality of sleep.[590] In a study involving elite male and female athletes, cyberbullying via social media and other online mechanisms was correlated with increased eating disorder pathology (low self-esteem, negative emotions, body dissatisfaction and unhealthy eating behaviours).[591]

There are reports of athletes being targeted by cyberbullying[582] and 'cyberstalking'.[592 593] Two studies from World Athletics reported a significant number of abusive posts regarding elite track and field athletes on social media, with 20% being racially motivated and 60% directed towards female athletes.[594] Sexist and sexualised abuse comprised over 40% of all posts, with a majority targeting females.[594]

Other impacts of social media use

Qualitative studies involving elite female athletes' experiences of social media use—particularly Instagram—have highlighted risks such as sexualisation of appearance, abuse and misogyny,[595 596] all of which may result in self-censorship (ie, refraining from expressing thoughts, points of view, or beliefs that others might find objectionable).[595] Furthermore, one study demonstrated an association between increased social media use (Facebook time) and worse competitive performance.[587]

Br J Sports Med: first published as 10.1136/bjsports-2025-111514 on 11 August 2026. Downloaded from http://bjsm.bmj.com/ on August 12, 2026 at Universitetsbiblioteket i Tromsø . Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies.

Conversely, positive impacts of social media use include building connections worldwide, communicating with family and friends, gaining support, emotional connection and sponsorship opportunities.[590 596–600] Among elite Para athletes, social media can counter negative disability stereotypes[601] and, for individuals who feel isolated or stressed, including LGBTQIA+ adolescents or those experiencing mental health symptoms,[581 602 603] social media can be supportive. One study found that the overwhelming majority of Twitter responses to the disclosure of mental health experiences by prominent athletes was positive (94%), with most responses applauding the inspiring message or providing encouragement or respondent self-disclosures of mental health symptoms.[603] Comparing oneself favourably to others may mediate anxiety and well-being in athletes who otherwise report low psychological capital or positivity.[586]

Other contextual factors

Geographical differences may influence whether Instagram posts related to a specific athlete's sporting success or failure impact athletic motivation and self-efficacy.[604] Specifically, the salience or impact of Instagram posts may vary according to user identification with social comparison groups. Additionally, the type, modality and timing of social media platforms used by athletes may have a differential impact on mental health.[605] For example, Facebook was used by 68–72% of athletes during the time frame extending from 2 hours prior to competition to 2 hours after competition, and by 32% during competition; having push notifications enabled on this platform during this time frame was significantly associated with sport anxiety.[606]

Social media safety

Efforts to protect athletes from the negative effects of social media are important, and evidence-based approaches are needed. A recent study evaluating a brief behavioural intervention in collegiate athletes found no meaningful reduction in the negative mental health effects associated with social media use.[585] The Mindful Social Media e-learning course by the IOC provides elite athletes with guidance for the strategic use of social media and how to mitigate risks.[599] As of 2024, the IOC also implemented a service to detect and respond to cyber abuse during the Olympic Games.[607] Harmful posts were detected immediately and removed, often before being viewed by the athlete.[599 607]

At the individual athlete level, clinicians should discuss social media use with elite athletes within ongoing clinical care. This information can be used to help the clinician and athlete understand how it is functioning to enhance (eg, via social connection) or impair (eg, via abuse, cyberbullying, negative body image or compromised sleep) mental health and performance, and to determine whether changes in social media use would be clinically beneficial. Clinicians should provide elite athletes with guidance on the potential mental health impacts of social media use, while recognising that this is a rapidly evolving area in terms of both technology and emerging evidence.

How injury, performance and mental health intersect

The relationship between injury, performance and mental health in elite athletes is complex.[77 561 608–611] Psychosocial factors may increase the likelihood of injury.[612–617] Injury can unmask or precipitate mental health symptoms or disorders,[616 618] and if injury leads to chronic pain, the athlete may develop mental health symptoms and disorders independent of the original injury.[95 619 620] Finally, mental health symptoms or disorders can increase the likelihood of prolonged injury recovery.[612 614 615 621–623]

Psychological and sociocultural risk factors for injury

Psychological and *sociocultural* factors are potential injury risk factors in sport.[77 610 624 625] Psychological risk factors include anxiety/worry, hypervigilance, anger/hostility, lack of attention, indecision, poor body image, low self-esteem, perfectionism, limited coping resources, risk-taking behaviours and low mood state.[610 611 626 627] Sociocultural risk factors include life event stress (eg, death of a family member, new school), limited social resources, prior experiences of sexual or physical abuse,[422 628] social pressures, organisational stress,[629] stress related to negative self-appraisal of athletic and academic performance, coaching quality (eg, poor athlete–coach relationship and communication) and the culture of sport and teams (eg, 'win at all costs' vs growth/improvement mindset).[77 616 630 631]

Limited prospective studies on the association between specific mental health symptoms or disorders and injury risk exist, with conflicting results. In professional soccer players, no relationship was found between symptoms of common mental health disorders and the onset of severe musculoskeletal time-loss injuries. However, those athletes who sustained severe musculoskeletal time-loss injuries were more likely to develop subsequent symptoms of common mental health disorders.[618] In contrast, in male and female collegiate athletes from different sports, those with pre-season anxiety symptoms and males with co-occurring depressive and anxiety symptoms had a significantly higher injury incidence rate compared with athletes without pre-season anxiety or co-occurring depression and anxiety.[626]

A meta-analysis found that a high stress response (ie, negative emotional response after sport injury or other stressful events) and history of life event stressors had the strongest associations with injury rates.[614] A high stress response and life event stress can cause inattention, distraction and increased self-consciousness while increasing muscle tension and impairing coordination, all of which can impede performance and increase the risk for injury.[614 625 630–636] When teammates and coaches are sources of stress (defined using the Life Event Survey for Collegiate Athletes), there is an associated increased risk of both acute and overuse injuries.[631] Emotional reactivity (ie, involuntary and overly intense emotional reactions) is also associated with injury.[612 637 638]

Psychological response to injury

The psychological response to injury should be assessed and managed to improve the health and well-being of elite athletes. This requires ongoing contact between the injured athlete and their health and performance team. The psychological response to injury includes any combination of cognitive, emotional and behavioural responses and is different for each individual.[1] *Cognitive* responses to injury may include concerns about re-injury, doubts about competency, low self-efficacy, loss of identity and concerns about medical staff competency.[620 639 640] For example, a fear avoidance model (FAM) that includes preoperative fear of movement/reinjury, low self-efficacy and pain catastrophising found that those with a 'moderate to high FAM risk' were less likely to RTS at 12 months and had poorer patient-reported outcomes including knee function and quality of life up to 12 months after anterior cruciate ligament (ACL) reconstruction surgery.[640] The mode of ACL injury may also have an impact on psychological readiness to RTS, with lower readiness

Br J Sports Med: first published as 10.1136/bjsports-2025-111514 on 11 August 2026. Downloaded from http://bjsm.bmj.com/ on August 12, 2026 at Universitetsbiblioteket i Tromsoe . Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies.

if the injury occurs during versus outside the athlete's primary sport.[641]

Emotional responses to injury may include sadness/crying, depression, suicidal ideation, anxiety, lack of motivation, irritability/anger, changes in appetite and sleep, low vigour, disengagement and burnout.[77 221 618 642 643] Injured athletes report more symptoms of depression and generalised anxiety disorder than non-injured athletes.[221] Whereas emotional responses to injury vary, problematic emotional responses are those that do not resolve, worsen over time or in which the symptoms seem disproportionate to injury severity.[77]

Finally, injury may trigger or unmask *behavioural* responses including increased aggression, social withdrawal, problem gambling, DE or EDs and substance use disorders.[626 644–656]

Psychological factors affecting injury rehabilitation

Persisting mental health symptoms following injury can prolong injury recovery.[610 611] In contrast, athletes with more positive cognitive, emotional and behavioural responses to injury may have improved injury recovery.[182 615 620 657–666] Higher levels of optimism, motivation, resilience and self-efficacy and lower levels of depression and stress following injury are associated with improved recovery.[182 610 615 657–665] Scoring higher on psychological readiness and pain self-efficacy and lower on kinesiophobia and pain catastrophising has been associated with enhanced recovery.[667 668]

Psychological interventions to enhance injury recovery and return to sport

Screening for psychological readiness after injury is an important component of the RTS decision, and several tools are available (table 6).[28 29 77 659 668–681] Additionally, psychological interventions aiming to provide support and enhance RTS might be considered, including: (1) reducing re-injury anxieties using modelling techniques (eg, mindfulness, imagery, graded exposure therapy, videos of athletes who have been successful in RTS)[668]; (2) fostering athlete autonomy (eg, providing athletes with choices, encouraging athlete-driven goal setting, collaborating rather than directing and respecting athletes' perceived sense of control); (3) building confidence and self-efficacy through functional tests and goal setting; (4) providing social support; (5) keeping athletes involved in their sport but avoiding premature RTS; (6) stress inoculation training (a cognitive behavioural approach to manage stress-related anxiety) when surgery is required[621 665 682]; and (7) screening for psychosocial barriers and mental health symptoms.[668]

Mental health emergencies

A mental health emergency exists when an acute disturbance of mental state is associated with a high level of risk and requires an immediate response.[683–685] The disturbance may involve agitation, aggression and/or impaired insight and judgment.[685] Delayed presentation of mental health symptoms—whether due to stigma, limited resources or reluctance to seek help among elite athletes—may contribute to emergency presentations.

A broad range of mental health symptoms or disorders may be associated with an emergency presentation. Any disorder could in certain circumstances present an emergency scenario.[685–687] EDs are a particular concern in sport.[72] Specifically, suicide attempts are seen in roughly one in five bulimia nervosa sufferers and one in eight anorexia nervosa sufferers,[688] and over-exercise in individuals with DE shows a significant association with suicidality.[689] Athletes may use or misuse substances, which increase the risk of an emergency presentation linked to mental health. For example, anabolic androgenic steroids can be associated with hypomanic and psychotic symptoms, aggression and suicide.[350 352 690] Stimulants can be associated with acute anxiety, mood disturbance, aggression, impaired judgment, disorientation and psychosis.[354 691 692] The sympathomimetic agent mephentermine has also been associated with psychosis, especially among chronic users.[356 357] Cannabis now contains much higher tetrahydrocannabinol levels than in the past, conferring a greater risk for psychosis.[693] Stressors such as injury[694] may evoke emergency presentations in part due to use of substances such as alcohol for coping.[695]

Athletes may have acute emotional reactions to difficult sport circumstances such as deselection,[696] which can be experienced as a form of rejection. Those with coexisting personality traits or disorders such as borderline personality disorder may rapidly develop emotional and/or behavioural instability in such situations because of the combination of impulsivity, highly reactive mood, sensitivity to perceived rejection, anger and recurrent suicidal behaviours.[176]

Data on the prevalence of mental health emergencies among elite athletes is sparse.[686] It may be that acute symptoms resolve quickly and/or are not recorded, with injury or performance concerns taking priority for recording and reporting at major events.[685] Additionally, there has historically been a lack of reporting systems for the purposes of systematically recording mental health presentations among populations of elite athletes.[40] Efforts have been made to improve recording during large sporting events and thus to achieve parity with the quality of data that is collected in relation to general medical concerns.[40]

Table 6 Examples of screening tools for psychological readiness to return to sport (RTS) after injury

Screening tool	Description of what the tool measures
Sport Mental Health Assessment Tool 2 (SMHAT2)[29]	Mental health symptoms and disorders in elite athletes
Tampa Scale-11 for Kinesiophobia[670 671]	Pain-related fear of movement
Re-Injury Anxiety Inventory (RIAI)[672]	Re-injury anxiety
Injury-Psychological Readiness to Return to Sport Questionnaire (I-PRRS)[659]	Psychological readiness of injured athletes to return to sport participation
Fear-Avoidance Beliefs Questionnaire (FABQ)[673]	Fear avoidance beliefs about physical activity and how they contribute to low back pain and disability
ACL-Return to Sport After Injury Inventory (ACL-RSI)[674]	Psychological readiness to return to sport participation after anterior cruciate ligament reconstruction
Pain Self-Efficacy Questionnaire[675]	Confidence to complete tasks despite pain
Pain Catastrophising Scale[676]	Negative or exaggerated response to actual or perceived pain

Modified from Herring *et al*.[77]

Br J Sports Med: first published as 10.1136/bjsports-2025-111514 on 11 August 2026. Downloaded from http://bjsm.bmj.com/ on August 12, 2026 at Universitetsbiblioteket i Tromsø . Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies.

The assessment of a mental health emergency in an athlete should include an exploration of the influence of any sport-specific factors that are relevant to the athlete presentation or likely to influence their treatment and recovery plan (eg, perceived or actual performance 'failure', career-altering events or interpersonal conflict between athlete and sport personnel).[697]

The management principles of emergency situations in sport have been described as 'Predict; Prevent; Prepare'.[698] All staff who work in the sport environment should receive appropriate education and practice related to procedures, roles and skills involved with the response to emergency situations.[687] Guidelines for mental health emergencies in elite athletes elaborate on this by defining what scenarios could arise and advising written procedures for how these will be addressed.[561 685 686 699] This includes consideration of the impact of travel on existing procedures, which may need to be adapted accordingly,[700] and where familiarity with local mental health services is advised.[686] Guidelines also describe the importance of rehearsing procedures in advance[701] and reviewing procedures once emergencies have resolved.[686 699] However, many sporting institutions still lack clear guidelines on procedures for mental health emergencies.[702] To encourage and support sporting institutions to prepare for mental health emergencies, an IOC guideline document outlining recommendations for the planning of mental health services at major sporting events was published in 2025. This document outlines details for the pre-event planning of mental health services, implementation during sporting events and post-event evaluation. The publication provides sport organisations with detailed recommendations for the development of a mental health emergency action plan.[701] Regarding specific strategies for emergencies, verbal de-escalation techniques can be extremely useful in the immediate management of acute behavioural disturbance and may obviate the need for pharmacological interventions or physical restraint.[686 703] If de-escalation is not successful, pharmacological intervention may be required. Depending on the underlying cause, this may include sedative, anxiolytic or antipsychotic medication.[686]

Transitions in elite sport

Transitions in elite sport can be temporary or permanent and can include promotions or demotions from one competitive level to another, transfer to another team (which could involve a move to another country and/or a different culture), return from a major competition (eg, Olympics/Paralympics), absences for short- or long-term illness, injury or pregnancy, temporary transitions related to other life circumstances (eg, family crisis or death), the off season, deselection or retirement.[511 704–708] Retirement has been the best studied of these transitions.[709–712]

During the transition from elite sport into retirement, a variety of general and sport-specific factors interact to potentially induce mental health symptoms or impact an athlete's quality of life. Several reviews identified common barriers (eg, involuntary retirement, low social support, career dissatisfaction) associated with difficult transitions and facilitators (eg, sport-life balance, retirement planning, educational attainment) of successful ones (table 7).[709–713] Especially when barriers dominate, mental health symptoms or disorders including anxiety, depression, body dissatisfaction, EDs/DE, grief, identity confusion, problem gambling, sleep disorders and sleep concerns, social isolation and/or substance misuse may develop or worsen.[511 707–712] According to two systematic reviews, these negative consequences may occur in 15–20% of transitioning athletes, especially those who leave because of injury, deselection or with extreme career dissatisfaction.[710 711]

Table 7 Barriers to and facilitators of elite athlete transition into retirement[511 707–712]

Barriers	Facilitators
High athletic identity	Good sport/life balance
High perfectionism	Positive self-perception
Sport career dissatisfaction	Sport career satisfaction
Involuntary retirement (injury, deselection)	Voluntary retirement (eg, new life direction)
Adverse life events/stressors	Social support network
Despair about future career	Longer length of time since retirement
Lack of coping strategies	Psychological resilience
Low social support/social isolation	Transition support programme involvement
Lack of retirement career planning	Planned post-retirement occupation
Education and resource dissatisfaction	Educational attainment
Financial distress/bankruptcy	Financial planning and support
Coach/athlete conflicts	Ongoing connections with supportive coaches
Loss of structure/daily schedule	Strong organisational and coping strategies
General health problems	Good general physical health
Osteoarthritis and chronic pain	Healthy bones and joints
Multiple prior concussions	Healthy neurological/cognitive functioning
Increasing body mass index	Maintaining healthy nutrition, physical activity and fitness
Chronic pain	Absence of serious injuries or chronic pain
Harassment and abuse while competing	Safe sporting environments
Sports-centred lifestyle	Developing interests/skills outside of sports
Absence of transition mental health resources	Accessed transition mental health support (eg, psychotherapy)
Perceived lack of control over retirement	Perceived control over life and transition
No organised retirement planning	Accessed pre-transition/retirement planning
Intense training/competition until retirement	Gradual reduction of training/competition
Poorly developed transferable skills	Transferable skills (eg, communication or planning skills from leadership positions held within sport)

The leadership of sports organisations and teams as well as members of an athlete's family and entourage can ease the transition into retirement, especially for those who leave involuntarily or who are already struggling with mental health symptoms or disorders. During an athlete's career, individuals in the athlete ecosystem can advocate for a balanced emphasis on mental well-being, physical health, personal growth, identity diversification and dual career development by supporting on-site or regional networks of cost-free mental health and wellness services and player development/career assistance programmes.[15 714–728] Existing mental health and career assistance positions and programmes can facilitate sports excellence, continuing education, financial literacy, coping/life skills development, mental well-being and transition planning both before and after transitions out of sport.[729–733]

Other strategies to facilitate positive transitions from sport include: (1) creating individualised 'retirement support frameworks' (ie, establishing interests and identities outside of sport, longitudinal pre-retirement planning, building adaptive coping strategies and post-sport mentorship and support)

Br J Sports Med: first published as 10.1136/bjsports-2025-111514 on 11 August 2026. Downloaded from http://bjsm.bmj.com/ on August 12, 2026 at Universitetsbiblioteket i Tromsoe . Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies.

and designating specific individuals to monitor their effectiveness[725]; (2) strengthening the 'exit health examination' by conducting mental health screening and providing linkages to mental health services[186]; (3) providing a comprehensive longitudinal 'After Career Consultation' that promotes a healthy lifestyle, develops future employment opportunities and seeks to prevent/manage osteoarthritis and mental health/cognitive problems[733]; (4) connecting athletes with support and resources from their International Federations and National Governing Bodies, sports organisations, corporate sponsors or foundations[45 734 735]; and (5) expanding access to the IOC's or each National Olympic Committee's (NOC's) Career/Professional Development Programmes that include life skills/dual career development and psychological counselling services.[731 736 737] Since elite athletes who continue to train and compete may find it difficult or distracting to focus on life after sports, their organisations should commit to creating environments that facilitate support for all types of transitions, but especially transition into the early retirement years.[511 724 738 739] Elite female athletes, who tend to retire earlier than elite male athletes—in part because of inadequate financial support and work-family conflict—may merit special attention and support.[739]

Elite Para athletes

Although athletes with disabilities may experience the same stressors of elite sport as discussed in earlier sections of this paper, participation in elite Para sport is uniquely associated with additional risk factors for mental health symptoms and disorders that are distinct from those observed in athletes with no disability. The classification process may be a significant stressor that many elite Para athletes feel unprepared to navigate.[740 741] Athletes may also struggle with the 'superhuman' media narrative that fails to acknowledge authentic identities both in and out of sport.[740] Additional athlete stressors may include disability stigma and bias, inaccessible venues and accommodations, lack of Para sport-specific coaching, poor audience attendance at Para sport events and cost of Para sport equipment.[741] Transition out of Para sport is often a particularly challenging time for athletes due to uncertainty regarding post-sport employment and loss of athletic identity, particularly if the transition is sudden such as with an unexpected injury or unanticipated issues related to classification.[742] Para athletes are also at higher risk for psychological, physical and sexual interpersonal violence compared with non-disabled peers.[278] Importantly, elite Para athletes report a higher rate of unmet health care needs compared with athletes without disability.[743] While mental health research in Para sport has primarily focused on athletes with physical disabilities or those who are blind or have low vision, non-apparent disabilities such as learning disabilities can also have a significant impact on mental health.[744]

The prevalence of mental health symptoms and disorders in elite Para athletes varies by disability type and sport classification and is further influenced by the use of different assessment tools, many of which are not validated for this population.[745] Sleep disturbance,[92 94 95] anxiety[95 217 218] and depression[95 217 218] are the most frequently reported psychological outcomes reported in the elite Para athlete literature. Among elite Australian Olympic and Paralympic athletes, no significant differences were observed between athletes from Para and non-Para sports for most mental health symptoms, although elite Para athletes reported lower alcohol consumption and self-esteem.[438]

Routine mental health assessment is important for elite Para athletes, and targeted screening and support may be particularly important during the time of transition out of sport. There is a paucity of mental health intervention research in this population. In one pilot randomised controlled trial, athletes undergoing a 'mindfulness-acceptance-commitment' programme with a motivational interviewing framework demonstrated improvements in life satisfaction, personal growth, positive relationships with others and reduced perceptions of pain in comparison to controls.[746] There may be many factors that affect access to mental health services for elite Para athletes, including transport barriers, inaccessibility of facilities and language barriers. For example, there are relatively few mental health professionals competent to offer services in sign languages. These issues need to be considered in treatment planning.[747]

Elite young (adolescent) athletes

In the general population, an estimated 50% of common mental health disorders—such as anxiety, mood, eating, substance use and neurodevelopmental disorders—emerge before the age of 18,[748] underscoring the need for appropriate screening and assessment among elite young athletes.[749] A confluence of biopsychosocial factors contributes to mental health symptoms and disorders during this developmental phase, including puberty, social disadvantage, trauma, identity formation challenges, peer conflict and harmful social media use.[750 751] Sex is a strong correlate, with adolescent girls more likely than boys to report anxiety, mood and eating disorders, and boys more likely to report problematic substance use.[751]

Elite young athletes across different cultures also encounter sport-specific challenges that impact their mental health, including heightened performance expectations (from self and/or others), adverse effects of early sport specialisation such as overuse/musculoskeletal injury[752] and high athletic identity,[753] possible exposure to interpersonal violence in sports settings, heightened susceptibility to injury and overtraining or burnout.[754 755] Such stressors are usually experienced alongside existing academic or vocational demands during this stage of life,[15] which should be considered and supported.[756]

To complement the recent IOC consensus statement on elite young athletes,[526] Gwyther and colleagues[757] conducted a scoping review examining the mental health of high-level athletes aged 12–17 years who travel and compete internationally. Data from the 33 studies that met the inclusion criteria indicated that elite young athletes report mental health symptoms at similar rates to their adult counterparts. DE is over-represented in the literature (n=16), particularly among female gymnasts. Conversely, disorders such as ADHD and autism spectrum disorder are under-represented in the literature on elite young athletes, despite growing recognition of neurodevelopmental disorders and the potential challenges they may bring in sporting contexts.[744 758] Factors commonly associated with poorer mental health outcomes in elite youth athletes were sex (female for DE and mental well-being; male for substance use), mental health comorbidities and various forms of perceived and/or objective stress.[757] Common factors associated with positive mental health outcomes included positive self-esteem, strong social support and healthy relationships with peers, family and coaches.[757]

Research has also considered the mental health of elite-striving young athletes in development or pathways programmes, including elite sporting schools. Most of this research has focused on burnout, EDs and sleep disturbance,[312 759–762] with few studies examining the prevalence of a wider range of mental health symptoms and disorders. Moderate to severe depressive symptoms have been reported to range from 9.5% to 19%.[763 764]

Br J Sports Med: first published as 10.1136/bjsports-2025-111514 on 11 August 2026. Downloaded from http://bjsm.bmj.com/ on August 12, 2026 at Universitetsbiblioteket i Tromsoe . Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies.

There is currently no clear indication as to whether elite-striving young athletes are more or less likely than their non-athletic peers to report experiencing mental health symptoms and disorders, with research reporting mixed results in this regard.[765 766] In addition to sex[764] and younger age,[242] key risk factors for mental health symptoms and disorders in young elite-striving athletes include deselection,[696 765] experiencing performance-related shame, guilt or embarrassment,[767 768] perfectionism,[769 770] participating in individual rather than team sports[771] and negative parental style such as stigmatising attitudes towards mental health symptoms and disorders, authoritarian approach and neglect.[772]

Research conducted in low- and middle-income countries, which are less studied, indicates unique challenges for young athletes in these settings, including a shortage of mental health professionals and lack of resources.[773–775] Life satisfaction was positively correlated with mental toughness and religiosity among youth athletes from Indonesia.[776] Non-conventional approaches to addressing mental health symptoms and disorders in these locations may represent culturally acceptable pathways to healing but can also have uncertain risks versus benefits.[773–775]

Sport as a subset of physical activity to promote mental health

While this consensus statement focuses on mental health symptoms and disorders experienced by elite athletes through their sport career, it remains essential to acknowledge that sport, as a subset of physical activity, is generally beneficial for short- and long-term physical and mental health.[777] Specifically, remaining regularly physically active through sport provides benefits for mild to moderate mental health symptoms for most athletes and can lead to improved sleep, stress relief and alertness, increased energy and favourable weight management and body image.[777–779] Participation in sport is also linked to higher self-esteem and higher life satisfaction.[780] These effects might be explained by physiological reactions (eg, increased blood circulation in the brain, improvement of hypothalamus-pituitary-adrenal axis functions) as well as by distraction, self-efficacy and social interaction.[779] Mental health outcomes are likely to depend on the type and the social environment of sport, as well as the way in which it is delivered.[781]

FINAL CONSIDERATIONS

Actionable recommendations

The content of this report reflects the evidence and themes presented and discussed at the in-person meeting of the Mental Health in Elite Athletes Consensus Group. Following that meeting, a Delphi voting process was conducted. Sixty-three recommendations were rated in the first round of ratings, 12 in the second round and seven in the third and final round, with modifications occurring between rounds to statements not meeting consensus criteria for inclusion. At the end of this process, all 63 statements met the inclusion criteria. These actionable recommendations are listed in table 8.

Dissenting viewpoints

Twenty-four of the finalised recommendations had at least one score of 4 or 5; while meeting consensus criteria for inclusion, these scores nonetheless represented lower enthusiasm or questions by panel members about a recommendation for inclusion. Open-ended comments corresponding to these ratings are presented in online supplemental file 5 and clustered into six thematic categories. Ten of these recommendations had feedback related to wording clarity or specificity. These comments were closely reviewed by the working group to determine whether minor revisions could improve readability without changing the substance of the recommendation. Five recommendations had comments related to feasibility, specifically regarding the resources required for implementation across settings. Four recommendations prompted questions about alignment with the current evidence base or with experts' clinical experience. Three comments addressed the intended audience for recommendation—for example, whether the responsibility for preventing interpersonal violence should extend to athletes. Two recommendations received feedback regarding the appropriateness of the recommendation as worded, including suggestions that athlete consent may be necessary before involving coaches or parents in mental health initiatives. For two additional recommendations, respondents questioned whether the proposed screening and assessment tools are valid and reliable across contexts.

DISCUSSION

Although research on elite athlete mental health has expanded rapidly in recent years, there remains no universally accepted gold standard guidance that is applicable across populations and settings. As such, a consensus approach using Delphi methodology was undertaken to aggregate expert opinion. This paper describes the state of the literature on elite athlete mental health and summarises the clinical key points in the form of actionable recommendations for health care providers of elite athletes.

Potential benefits of this consensus statement include increased awareness of the importance of mental health in elite sport. Furthermore, it aspires to improve the quality of health care afforded to elite athletes across settings. That is, some degree of standardisation of care - aligned with the evidence base described in this statement - across countries and sports could improve care for many elite athletes. However, variations in the cultures that exist within different geographic locations, demographic groups and sports remain challenging. These variations must be acknowledged so that these recommendations are considered in contextually appropriate ways. The clear actionable nature of the recommendations and the attention to making them relevant across diverse contexts will help support effective implementation. Barriers to application include widely varying resource levels of target audiences and continued mental health stigma. More research about elite athlete mental health in low- and middle-income countries was identified as a critical area in the research agenda generation that occurred in tandem with this consensus process.[782]

This research has some important limitations. There was potential bias in findings based on the panel membership and the evidence base, both of which disproportionately came from high-income countries. Furthermore, while athlete and International Sports Federation representation was included, other members of the elite athlete entourage, including elite athlete coaches and the full spectrum of high-performance support staff, were not represented. Additionally, the wide variation in types of quantitative and qualitative research included in this statement precluded formal systematic assessment of quality of the included studies. Finally, given the rapid evolution of this field, it will be important to update this statement regularly, ensuring that recommendations remain current and reflect emerging evidence.

To address these limitations, future research should continue to involve athletes as partners, employ robust and culturally responsive methodologies and emphasise real-world implementation

Br J Sports Med: first published as 10.1136/bjsports-2025-111514 on 11 August 2026. Downloaded from http://bjsm.bmj.com/ on August 12, 2026 at Universitetsbiblioteket i Tromsoe . Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies.

Table 8 Actionable recommendations for health care providers of elite athletes

Topic	Recommendation
General	Supporting elite athlete mental health means not only preventing or treating mental health symptoms and disorders but also promoting mental well-being.
	Elite athletes' mental health is dynamic and can change over time in response to sport-specific and everyday life demands. Elite athletes may also develop mental health symptoms and disorders. Timely support matched to individual needs can help improve mental well-being and address clinical problems.
	Treatment options to consider for mental health symptoms and disorders in elite athletes include psychotherapy, medication and other culturally or contextually appropriate modalities. The optimal approach should be individualised and person-centred, guided by the specific diagnosis, the severity of distress or dysfunction and the evidence base for particular management strategies.
Prevalence of mental health symptoms and disorders in elite athletes and their entourage members	Everyone involved in elite sport should be aware that mental health symptoms and disorders are experienced by not only athletes but also coaches, members of the elite athlete health and performance team and others in clinical and non-clinical roles.
Screening and surveillance of mental health symptoms and disorders in elite athletes	Screening and surveillance undertaken at regularly scheduled time points and at the time of significant occurrences/events can help identify elite athletes in need of further mental health evaluation and support.
	Screening and surveillance of mental health symptoms and disorders in elite athletes should be conducted using validated approaches.
	The IOC Sport Mental Health Assessment Tool 2 includes validated disorder-specific screening tools and clinical assessment and management guidance for physicians and other licensed/registered health professionals working with elite athletes.
Psychotherapy	Psychotherapy, either as monotherapy or combined with other non-pharmacological and pharmacological strategies, is a foundational component of the management of mental health symptoms and disorders in elite athletes.
	Psychotherapy for elite athletes should be culturally responsive, recognising that culture influences how distress is experienced, expressed and managed.
Pharmacological treatment	When prescribing medications for mental health symptoms and disorders in elite athletes, health care providers should understand the potential negative impacts on sports performance, safety risks, and whether the medication is prohibited by any sport governing body, such as the World Anti-Doping Agency.
Mental health considerations for removal from and return to sport	With appropriate support and treatment, many athletes experiencing mental health symptoms or disorders can continue to train and compete safely.
	If elite athletes experience mental health symptoms and disorders that compromise safety, stability or function, it may be necessary to modify sport participation. For some athletes, this may include ceasing sport training and competition until they are cleared by a licensed clinician to return to sport.
	When planning for return to sport participation, it is important to consider supports and stressors in the elite athlete's sport ecosystem, including teammates, coaches, staff, family, organisations and other aspects of their sport environment.
Sleep	Assessment for sleep disorders and sleep concerns in elite athletes typically involves self-reports of symptoms, which may be supplemented by at-home or in-laboratory tests, depending on the disorder of concern and the available resources.
	Non-pharmacological treatments are recommended for many sleep disorders in elite athletes because they are effective and because many medications for sleep disorders are associated with increased injury risk and/or performance impairment.
	Sub-clinical sleep issues (eg, insufficient sleep, daytime sleepiness) can often be ameliorated through behavioural and environmental interventions.
Major depressive disorder and depressive symptoms; suicide	Differentiating between major depressive disorder, burnout and overtraining syndrome in elite athletes is important for providing the most effective treatment.
	Prevention of suicide in elite athletes may include strategies that foster mental health literacy, suicide risk factor identification, targeted support and structured safety plans.
Anxiety disorders and obsessive-compulsive and related disorders	It is important to distinguish anxiety and obsessive-compulsive disorder (OCD) symptoms from other common experiences within elite sport. For example, distinguish elevated arousal associated with competition from symptoms of an anxiety disorder, and distinguish athlete superstitions from OCD symptoms.
	Early detection and intervention are critical to ensuring that anxiety, OCD and related disorders do not become persistent and debilitative in elite athletes.
Trauma- and stressor-related disorders	Clinicians should recognise the crucial nature of the therapeutic alliance when working with elite athletes with trauma- and stressor-related disorders, as a positive therapeutic alliance increases the likelihood of interventions being effective.
	Trauma- and stressor-related disorders in elite athletes should be treated early, as that may reduce the risk of progression to more chronic mental health disorders such as post-traumatic stress disorder.
	When managing elite athletes with trauma- and stressor-related disorders, clinicians should consider team-based and/or individual athlete-based approaches, depending on the nature of the trauma and the impact on those involved.

Continued

Br J Sports Med: first published as 10.1136/bjsports-2025-111514 on 11 August 2026. Downloaded from http://bjsm.bmj.com/ on August 12, 2026 at Universitetsbiblioteket i Tromsoe . Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies.

Table 8 Continued

Topic	Recommendation
Eating disorders and Relative Energy Deficiency in Sport (REDs)	REDs is caused by severe and prolonged low energy availability (inadequate intake relative to energy needs) and manifests as impaired physiological and psychological functioning. Clinicians should assess athletes for REDs even in the absence of disordered eating or an eating disorder.
	When possible, a multidisciplinary team should manage most elite athletes who present with disordered eating, eating disorders or REDs. This may include a sports medicine physician, dietitian, licensed mental health professional(s) and other culturally or contextually appropriate carers.
	Screening for eating disorders in elite athletes poses challenges (under-reporting and difficulties in distinguishing between normative and pathological behaviours in an elite sport context). Validated screening instruments such as the Eating Disorder Examination Questionnaire (EDE-Q) can identify elite athletes at risk for eating disorders, but clinical interviews are needed to diagnose eating disorders.
	The 2023 International Olympic Committee's REDs Clinical Assessment Tool (IOC REDs CAT2) can help facilitate diagnosis of REDs, stratify severity, and guide treatment and sport participation decisions.
	Treatment of REDs should address presenting physiological dysfunction(s) (eg, low bone mineral density, GI dysfunction) as well as psychological outcomes from exposure to problematic low energy availability.
Attention-deficit/hyperactivity disorder (ADHD)	If stimulants are prescribed for medical treatment of elite athletes, clinicians should check if a Therapeutic Use Exemption is required.
	When evaluating individuals for ADHD, the assessment should include consideration of common comorbidities such as mood disorders, anxiety and related disorders, learning disorders, substance use disorders and sleep disorders.
Bipolar and psychotic disorders	It is important to distinguish bipolar and psychotic disorder symptoms from other common experiences within elite sport. For example, distinguish the effects of commonly misused substances in sport from primary bipolar/psychotic disorders, and distinguish high levels of activity associated with sport from mania.
	Athletes with severe and chronic mental health disorders who have transitioned out of elite competition often benefit from regular exercise, which may include continued participation in sport, for the purposes of improved physical health, mental health symptoms and functioning.
Personality disorders	Personality disorders should be considered in elite athletes who present with a pattern of disruptive behaviour, sport-related sanctions, legal difficulties and/or maladaptive perfectionism.
Substance use	Misuse of alcohol, tobacco/nicotine and cannabis is common among elite athletes. Early identification using validated screening tools, combined with brief interventions, can help prevent progression to substance use disorders.
	For some elite athletes, brief individualised motivational interventions—incorporating contingency management and involving coaching staff and/or family members—can shift attitudes and beliefs about substance misuse and reduce frequency of use.
	Clinicians need to be aware of the most commonly used performance-enhancing drugs and their mental health effects.
Gambling disorder and other behavioural addictions	Behavioural addictions are often 'hidden disorders' and thus are often difficult to detect unless elite athletes are asked targeted questions.
	Management of behavioural addictions in elite athletes primarily consists of cognitive behavioural therapy and/or motivational interviewing.
Concussion	Mental health symptoms should be monitored after concussion in elite athletes.
	Sub-symptom threshold aerobic exercise is recommended as an early intervention to manage post-concussive mental health symptoms in elite athletes.
	Cognitive rehabilitation and neurocognitive training are effective interventions for persisting mental health symptoms in elite athletes post-concussion.
Creating an environment that supports mental well-being and resilience in elite athletes	Everyone involved in elite sport should have a basic understanding of mental health, including how to recognise signs and symptoms of mental health disorders, access appropriate support and promote mental well-being. Clinicians with mental health expertise are well positioned to provide and interpret evidence-based information. Others in the sport ecosystem—such as organisational leaders, coaches and staff in non-clinical roles—can play a key role in supporting the dissemination and reinforcement of this information, helping to create environments that prioritise mental health.
	Encouraging the involvement of coaches, parents/caregivers and community resources as appropriate in mental health initiatives can help create a supportive sport ecosystem.
	Interpersonal violence in elite athletes is associated with an elevated risk of mental health symptoms and disorders as well as risk-taking behaviours. Clinicians should screen for a history of interpersonal violence both within and outside the sport context.
	Everyone in the sport ecosystem is responsible for protecting athletes from interpersonal violence.
Cultural considerations and inclusivity	Understanding, respecting and integrating an athlete's cultural background into care helps build trust and is essential for both accurate assessment and effective management.
	The potential impact of structural inequalities and social determinants of mental health should be considered when addressing mental health needs in elite athletes.
Barriers to mental health care seeking	Elite athletes may encounter numerous barriers to seeking and sustaining mental health care—including stigma, low mental health literacy, cultural perceptions, fear of negative judgment, concern about losing sport opportunities and logistical challenges. Clinicians should actively address these barriers with athletes and their support team.
	Interventions to decrease barriers to care seeking should be tailored to the unique needs and assets of a given athlete population and their context, and should consider the availability, accessibility, quality and fit of care.
Social media and mental health	Clinicians should provide elite athletes with guidance on the potential mental health impacts of social media use, while recognising that this is a rapidly evolving area in terms of both technology and emerging evidence.
	Clinicians should discuss social media use with elite athletes within ongoing clinical care. This information can be used to help the clinician and athlete understand how it is functioning to enhance (eg, via social connection) or impair (eg, via abuse, cyberbullying, negative body image or compromised sleep) mental health and performance, and to determine whether changes in social media use would be clinically beneficial.
How injury, performance and mental health intersect	The psychological response to injury should be assessed and managed to improve the health and well-being of elite athletes. This requires ongoing contact between the injured athlete and their health and performance team and provision of opportunities for support following injury.
	Screening for psychological readiness to return to sport can help with return to sport decisions following injury.

Continued

Br J Sports Med: first published as 10.1136/bjsports-2025-111514 on 11 August 2026. Downloaded from http://bjsm.bmj.com/ on August 12, 2026 at Universitetsbiblioteket i Tromsoe . Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies.

Table 8 Continued

Topic	Recommendation
Mental health emergencies	A broad range of mental health symptoms or disorders can lead to mental health emergencies in elite athletes, requiring an immediate response from trained personnel.
	Planning ahead (eg, creating and rehearsing an emergency action plan) is important for ensuring an appropriate response if a mental health emergency occurs.
Transitions in elite sport	Athletes are at risk of mental health challenges in the transition from elite sport, with risk heightened for athletes facing involuntary retirement, career dissatisfaction, persistent illness or injury, chronic pain, inadequate post-retirement planning or limited external support. Athletes should receive targeted mental health support both before and after their transition out of sport.
	Structured support, including career assistance programmes, is helpful for many athletes in their transition from elite sport. The International Olympic Committee and some International Federations and National Governing Bodies have well-developed career assistance programmes that can be shared with elite athletes.
Elite Para athletes	Health care providers for elite Para athletes should consider that participation may be associated with unique mental health stressors including disability stigma and bias, the classification system, inaccessible venues and accommodations and cost of Para sport equipment.
	Routine mental health assessment is important for elite Para athletes, who report a higher likelihood of having unmet health care needs compared with athletes with no disability.
	The transition out of Para sport can be a particularly high-risk period for elite Para athletes due to the loss of athletic identity and uncertainty around post-sport employment. Targeted screening and support should be available during this time.
Elite young (adolescent) athletes	Mental health symptoms and disorders frequently develop before 18 years of age, underscoring the need for appropriate screening and assessment among elite young athletes.
	The demands of high-performance sport often co-occur with academic and/or vocational commitments, contributing to potential cumulative stressors during the elite young athlete phase that should be considered and supported.
How sport can be used to promote mental health	Support elite athletes to maintain a healthy, sustainable relationship with their sport in a positive environment, so as to promote the positive mental health outcomes of continued sport participation.

and evaluation across varied sport settings. Priorities include generating stronger evidence for preventive and system-level interventions, advancing measurement tools appropriate for diverse athlete groups and exploring scalable approaches in low-resource contexts. Broadly, this work will strengthen the evidence base and support ongoing refinement of consensus-driven guidance. A systematic approach to aggregating expert feedback about areas for future research, addressing these limitations to the current evidence base, was undertaken in parallel to this consensus process. The results are reported separately.[782]

CONCLUSION

Mental health is an integral and essential component of elite athlete health and overall well-being. The responsibility for prevention, recognition and management of mental health symptoms and disorders and for promoting well-being extends across the entire elite sport ecosystem. Multiple interacting biopsychosocial and ecological factors influence athlete well-being, necessitating system-level attention and coordinated strategies. Elite athletes experience mental health symptoms and disorders at rates similar to, or higher than, the general population, and their presentations and care needs may differ from those of non-athletes. Ensuring that clinicians, coaches and sport organisations are prepared to recognise, respond to and proactively support mental health is essential to safe, ethical and sustainable elite sport participation.

Author affiliations
[1]Department of Psychiatry, University of Wisconsin-Madison School of Medicine and Public Health, Madison, Wisconsin, USA
[2]University Health Services, University of Wisconsin-Madison, Madison, Wisconsin, USA
[3]Section Sports Medicine, University of Pretoria, Pretoria, South Africa
[4]Amsterdam UMC location University of Amsterdam, Department of Orthopedic Surgery and Sports Medicine, Amsterdam, Netherlands
[5]Department of Pediatrics, University of Washington School of Medicine, Seattle, Washington, USA
[6]Center for Child Health, Behavior and Development, Seattle Children's Research Institute, Seattle, Washington, USA
[7]Health, Medicine and Science Department, International Olympic Committee, Lausanne, Switzerland
[8]Oslo Sports Trauma Research Center, Department of Sports Medicine, Norwegian School of Sports Sciences, Oslo, Norway
[9]Department of Physical Medicine and Rehabilitation, Shirley Ryan AbilityLab, Northwestern University Feinberg School of Medicine, Chicago, Illinois, USA
[10]Department of Neuroscience, Faculdade de Medicina do ABC, Santo Andre, Brazil
[11]Universidade de Sao Paulo Faculdade de Medicina Hospital das Clinicas Instituto de Psiquiatria, Sao Paulo, Brazil
[12]Athlete Representative, Hong Kong, China
[13]UK Sports Institute, The Manchester Institute of Health and Performance, Manchester, UK
[14]International Centre for Youth Gambling Problems and High-Risk Behaviors, McGill University, Montreal, Quebec, Canada
[15]Department of Psychiatry and Behavioral Neurosciences, McMaster University Michael G DeGroote School of Medicine, Hamilton, Ontario, Canada
[16]International Ski and Snowboard Federation, Oberhofen, Switzerland
[17]School of Human Sciences, University of Greenwich, London, UK
[18]Department of Psychiatry, University of Arizona Medical Center - University Campus, Tucson, Arizona, USA
[19]Department of Psychiatry, Chung Ang University Hospital, Seoul, Korea (the Republic of)
[20]Ebling Library for the Health Sciences, University of Wisconsin-Madison, Madison, Wisconsin, USA
[21]Graduate Institute of Sport Coaching Science, Chinese Culture University, Taipei City, Taiwan
[22]Medical Department, Federation Internationale de Football Association, Zurich, Switzerland
[23]Department of Psychiatry, University of Maryland School of Medicine, Baltimore, Maryland, USA
[24]Department of Family Medicine-Sport, McMaster University, Hamilton, Ontario, Canada
[25]Elite Sport and Mental Health, Centre for Youth Mental Health, The University of Melbourne, Melbourne, Victoria, Australia
[26]Orygen, Melbourne, Victoria, Australia
[27]Team Physician, US Women's National Soccer Team (independent contractor), Princeton, New Jersey, USA
[28]Movember Institute of Men's Health, Richmond, Victoria, Australia
[29]Department of Sports Medicine, Norwegian School of Sports Sciences, Oslo, Norway
[30]Department of Psychology, Stellenbosch University, Stellenbosch, Western Cape, South Africa
[31]Center for Public Health and Family Medicine, Western University, London, Ontario, Canada

Br J Sports Med: first published as 10.1136/bjsports-2025-111514 on 11 August 2026. Downloaded from http://bjsm.bmj.com/ on August 12, 2026 at Universitetsbiblioteket i Tromsoe . Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies.

[32]Department of Kinesiology and Health Promotion, University of Kentucky, Lexington, Kentucky, USA
[33]NYU Grossman School of Medicine, New York, New York, USA

Acknowledgements The authors acknowledge Dr Robin Mitchell (IOC Health, Medicine and Science Commission Chair) for his leadership and participation in the consensus meeting and Fiona Trabelsi (IOC Health, Medicine and Science Department) for her expert organisation of the consensus meeting logistics.

Contributors CLR, BH and VG were responsible for the design, development and collation of contributions. All authors were responsible for drafting the manuscript and were involved in the conception, revising and approval of the manuscript prior to submission. EK-H, MEH, CLR, BH and VG were responsible for drafting and implementing the methods. All authors agreed to be accountable for the work in ensuring that questions related to the accuracy or integrity of the work are appropriately investigated and resolved. The guarantor, CLR, accepts responsibility for the finished work and the conduct of the study and controlled the decision to publish.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Disclaimer This consensus paper provides an overview of mental health issues in elite athletes that are important to physicians and other clinicians who treat elite athletes. It is not intended as a clinical practice guideline or legal standard of care and should not be interpreted as such. This consensus paper serves as a guide and, as such, is of a general nature, consistent with the reasonable practice of the health care professional. Individual treatment will depend on the facts and circumstances specific to each individual case.

Competing interests CLR is Co-chair of the IOC Mental Health in Elite Athletes Consensus Group. VG is Co-chair of the IOC Mental Health in Elite Athletes Consensus Group and Chair of the IOC Mental Health Working Group. EK-H is a paid consultant of the National Collegiate Athletic Association. CB is an Associate Editor of the British Journal of Sports Medicine and a member of the IOC Health, Medicine and Science Commission. MG has received grants from Harmony Biosciences and has served as a consultant to Google, Vytalogy, Smartypants Vitamins, Kenvue, Haleon, Canyon Ranch and Wndr Hlth. MM is a member of the IOC Mental Health Working Group and is a Deputy Editor and member of the Injury Prevention & Health Promotion Editorial Board of the British Journal of Sports Medicine. RP was an external consultant on mental health to the IOC and is a member of the IOC Mental Health Working Group. JST is Director of Health, Medicine and Science at the International Olympic Committee and is a member of the Editorial Board of the British Journal of Sports Medicine. BH is Co-chair of the IOC Mental Health in Elite Athletes Consensus Group.

Patient consent for publication Not applicable.

Ethics approval Not applicable.

Provenance and peer review Not commissioned; externally peer reviewed.

Supplemental material This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: https://creativecommons.org/licenses/by-nc/4.0/.

ORCID iDs
Claudia L Reardon https://orcid.org/0000-0002-1767-1500
Vincent Gouttebarge https://orcid.org/0000-0002-0126-4177
Emily Kroshus-Havril https://orcid.org/0000-0002-7484-2601
Cindy Miller Aron https://orcid.org/0000-0002-6618-4653
Cheri Blauwet https://orcid.org/0000-0001-8568-1009
João Mauricio Castaldelli-Maia https://orcid.org/0000-0001-9621-2291
Jeffrey Lee Derevensky https://orcid.org/0000-0001-6947-2987
Carla Edwards https://orcid.org/0000-0002-2256-1963
Doug Hyun Han https://orcid.org/0000-0002-8314-0767
Mary E Hitchcock https://orcid.org/0000-0002-8254-388X
Andrew Massey https://orcid.org/0000-0002-8253-932X
David McDuff https://orcid.org/0000-0003-4088-3920
Rosemary Purcell https://orcid.org/0000-0002-7912-6073
Margot Putukian https://orcid.org/0000-0002-1478-8068
Simon M Rice https://orcid.org/0000-0003-4045-8553
Scott Sloan https://orcid.org/0009-0001-0255-0271
Torbjørn Soligard https://orcid.org/0000-0001-8863-4574
Jorunn Kaiander Sundgot-Borgen https://orcid.org/0000-0002-5291-5486
Leslie Swartz https://orcid.org/0000-0003-1741-5897
Jane S Thornton https://orcid.org/0000-0002-3519-7101
Brian Hainline https://orcid.org/0000-0002-0233-2434

REFERENCES

1 Reardon CL, Hainline B, Aron CM, *et al*. Mental health in elite athletes: International Olympic Committee consensus statement (2019). *Br J Sports Med* 2019;53:667–99.
2 Gouttebarge V, Bilgoe SC, Gorczynski P, *et al*. Supporting the IOC Consensus Statement on Mental Health in Elite Athletes: A systematic review and meta-analysis on the prevalence of mental health symptoms in elite sports. *Sports* 2026;14:296.
3 Kegelaers J, Wylleman P, *et al*. Mental health in elite-level coaches: Prevalence rates and associated impact of coach stressors and psychological resilience. *Int Sport Coaching J* 2021;8:338–47.
4 Pilkington V, Rice SM, Walton CC, *et al*. Prevalence and correlates of mental health symptoms and well-being among elite sport coaches and high-performance support staff. *Sports Med Open* 2022;8:89.
5 Ruddock S, Rahimi-Golkhanden S, Ruddock-Hudson M, *et al*. Tracking the mental health outcomes of occupational burnout with Australian Rules Football coaches: A 2-year longitudinal study. *J Sci Med Sport* 2019;22:S52.
6 Ruddock S, Ruddock-Hudson M, Rahimi-Golkhandan S. The impact of job-burnout on Australian Football League coaches: Mental health and well-being. *J Sci Med Sport* 2017;20:73–4.
7 Ruddock S, Ruddock-Hudson M, Rahimi-Golkhandan S. Examining within-season change of job-burnout and psychological distress for Australian Rules Football coaches. *J Sci Med Sport* 2018;21:S41–2.
8 Kim SSY, Hamiliton B, Beable S, *et al*. Elite coaches have a similar prevalence of depressive symptoms to the general population and lower rates than elite athletes. *BMJ Open Sport Exerc Med* 2020;6:e000719.
9 Åkesdotter C, Kenttä G, Eloranta S, *et al*. Prevalence and comorbidity of psychiatric disorders among treatment-seeking elite athletes and high-performance coaches. *BMJ Open Sport Exerc Med* 2022;8:e001264.
10 Eken M, Runciman P, Badenhorst M, *et al*. Evaluating workforce well-being: An investigation of healthcare professionals' mental health and burnout symptoms at the Tokyo 2020 and Beijing 2022 Paralympic Games. *Br J Sports Med* 2023;57:1382–7.
11 International Olympic Committee (IOC). IOC Mental Health in Elite Athletes Toolkit. 2021. Available: https://olympics.com/athlete365/articles/mentally-fit/ioc-mental-health-in-elite-athletes-toolkil-x3908
12 International Olympic Committee (IOC). Mental Health Action Plan. 2023. Available: https://stillmed.olympics.com/media/Documents/News/2023/07/Mental-Health-Action-Plan-2023.pdf?_ga=2.35011553.173243837.1693809757-2074473781.169156822
13 Henriksen K, Diment G, Kuettel A. The Team Denmark applied model of athlete mental health. *Int J Sport Exerc Psychol* 2024;22:2325–41.
14 Kuettel A, Pedersen AK, Larsen CH. To flourish or languish, that is the question: Exploring the mental health profiles of Danish elite athletes. *Psychol Sport Exerc* 2021;52:101837.
15 Stambulova NB, Ryba TV, Henriksen K. Career development and transitions of athletes: The International Society of Sport Psychology Position Stand revisited. *Int J Sport Exerc Psychol* 2021;19:524–50.
16 Lundqvist C, Andersson G. Let's talk about mental health and mental disorders in elite sports: A narrative review of theoretical perspectives. *Front Psychol* 2021;12:700829.
17 Morgan WP, Brown DR, Raglin JS, *et al*. Psychological monitoring of overtraining and staleness. *Br J Sports Med* 1987;21:107–14.
18 Purcell R, Gwyther K, Rice SM. Mental health In elite athletes: Increased awareness requires an early intervention framework to respond to athlete needs. *Sports Med Open* 2019;5:46.
19 Henriksen K, Schinke R, Moesch K, *et al*. Consensus statement on improving the mental health of high performance athletes. *Int J Sport Exerc Psychol* 2020;18:553–60.
20 Agnew D, Pill S. Creating caring environments: An exploration of football managers and coaching. *Sports Coaching Review* 2023;12:27–46.
21 Kuettel A, Larsen CH. Risk and protective factors for mental health in elite athletes: A scoping review. *Int Rev Sport Exerc Psychol* 2020;13:231–65.
22 Campbell R, Dworkin E, Cabral G. An ecological model of the impact of sexual assault on women's mental health. *Trauma, Violence, & Abuse* 2009;10:225–46.
23 Taylor E. We agree, don't we? The Delphi method for health environments research. *HERD* 2020;13:11–23.
24 Trevelyan EG, Robinson PN. Delphi methodology in health research: How to do it? *Eur J Integr Med* 2015;7:423–8.

Br J Sports Med: first published as 10.1136/bjsports-2025-111514 on 11 August 2026. Downloaded from http://bjsm.bmj.com/ on August 12, 2026 at Universitetsbiblioteket i Tromsø . Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies.

25 BMJ. BJSM author guidelines for consensus statements. 2022. Available: https://bjsm.bmj.com/pages/authors#consensus_statement
26 Diamond IR, Grant RC, Feldman BM, *et al*. Defining consensus: A systematic review recommends methodologic criteria for reporting of Delphi studies. *J Clin Epidemiol* 2014;67:401–9.
27 Silvester C, Hume PA, Clark T, *et al*. The perspectives of high-performance athletes on mental health assessment: A mixed-methods exploratory study. *Int J Sport Exerc Psychol* 2025;1–22.
28 Gouttebarge V, Bindra A, Blauwet C, *et al*. International Olympic Committee (IOC) Sport Mental Health Assessment Tool 1 (SMHAT-1) and Sport Mental Health Recognition Tool 1 (SMHRT-1): Towards better support of athletes' mental health. *Br J Sports Med* 2021;55:30–7.
29 Gouttebarge V, Bindra A, Blauwet C, *et al*. International Olympic Committee Sport Mental Health Assessment Tool 2 (IOC SMHAT2): development, content validity and feasibility. *Br J Sports Med* 2026;60:1131–42.
30 Anderson T, Adams WM, Bartley JD, *et al*. Analysis of the Sport Mental Health Assessment Tool 1 (SMHAT-1) in Team USA athletes. *Br J Sports Med* 2023;57:1187–94.
31 Bahr R, Clarsen B, Derman W, *et al*. International Olympic Committee Consensus Statement: Methods for Recording and Reporting of Epidemiological Data on Injury and Illness in Sports 2020 (Including the STROBE Extension for Sports Injury and Illness Surveillance (STROBE-SIIS)). *Orthop J Sports Med* 2020;8:2325967120902908.
32 Verhagen E, Clarsen B, Capel-Davies J, *et al*. Tennis-specific extension of the International Olympic Committee consensus statement: Methods for recording and reporting of epidemiological data on injury and illness in sport 2020. *Br J Sports Med* 2021;55:9–13.
33 Clarsen B, Pluim BM, Moreno-Pérez V, *et al*. Methods for epidemiological studies in competitive cycling: An extension of the IOC consensus statement on methods for recording and reporting of epidemiological data on injury and illness in sport 2020. *Br J Sports Med* 2021;55:1262–9.
34 Murray A, Junge A, Robinson PG, *et al*. International consensus statement: Methods for recording and reporting of epidemiological data on injuries and illnesses in golf. *Br J Sports Med* 2020;54:1136–41.
35 Greenspan S, Munro D, Nicholas J, *et al*. Circus-specific extension of the International Olympic Committee 2020 consensus statement: Methods for recording and reporting of epidemiological data on injury and illness in sport. *BMJ Open Sport Exerc Med* 2022;8:e001394.
36 Waldén M, Mountjoy M, McCall A, *et al*. Football-specific extension of the IOC consensus statement: Methods for recording and reporting of epidemiological data on injury and illness in sport 2020. *Br J Sports Med* 2023;57:1341–50.
37 Spörri J, McGawley K, Alhammoud M, *et al*. Snow sports-specific extension of the IOC consensus statement: Methods for recording and reporting epidemiological data on injury and illness in sports. *Br J Sports Med* 2024;59:8–23.
38 Derman W, Badenhorst M, Blauwet C, *et al*. Para sport translation of the IOC consensus on recording and reporting of data for injury and illness in sport. *Br J Sports Med* 2021;55:1068–76.
39 Moore IS, Crossley KM, Bo K, *et al*. Female athlete health domains: A supplement to the International Olympic Committee consensus statement on methods for recording and reporting epidemiological data on injury and illness in sport. *Br J Sports Med* 2023;57:1164–74.
40 Mountjoy M, Junge A, Bindra A, *et al*. Surveillance of athlete mental health symptoms and disorders: A supplement to the International Olympic Committee's consensus statement on injury and illness surveillance. *Br J Sports Med* 2023;57:1351–60.
41 Ströhle A. Sports psychiatry: Mental health and mental disorders in athletes and exercise treatment of mental disorders. *Eur Arch Psychiatry Clin Neurosci* 2019;269:485–98.
42 Stillman MA, Glick ID, McDuff D, *et al*. Psychotherapy for mental health symptoms and disorders in elite athletes: A narrative review. *Br J Sports Med* 2019;53:767–71.
43 StillmanA, Ritvo EC, Glick ID. Psychotherapeutic treatment of athletes and their significant others. In: BaronDA, Reardon CL, Baron SH, eds. *Clinical Sports Psychiatry: An International Perspective*. New York: Wiley, 2013: 115–23.
44 Ackerman SR. Attitudes toward help-seeking and mental health among college athletes: Impact of a psycho-educational workshop. 2012. Available: https://scholarship.shu.edu/dissertations/37/
45 Hong HJ, Coffee P. A psycho-educational curriculum for sport career transition practitioners: Development and evaluation. *Eur Sport Management Quarterly* 2018;18:287–306.
46 Hays KF. The psychology of performance in sport and other domains. In: Murphy SM, ed. *The Oxford Handbook of Sport and Performance Psychology*. London: Oxford University Press, 2012: 24–45.
47 Pelz B. The science of accepting yourself and implementing sports and positive psychology in the clinical setting. *J Psychol Neurosci* 2024;6:1–9.
48 Edwards CD. Management of mental health challenges in athletes: Screening, pharmacology, and behavioral approaches. *Clin Sports Med* 2024;43:13–31.
49 Stull T, Glick I, Kamis D. The role of a sport psychiatrist on the sports medicine team, Circa 2021. *Psychiatr Clin North Am* 2021;44:333–45.
50 Anderson SA, Haraldsdottir K, Watson D. Mindfulness in athletes. *Curr Sports Med Rep* 2021;20:655–60.
51 Hassanmirzaei B, Khalifeh Soltani S, Haratian Z, *et al*. Web-based supportive psychotherapy to prevent posttraumatic stress disorder: A cross sectional study on the Iranian national under-14 girls' football team after Nepal earthquake in 2015. *Asian J Sports Med* [Preprint] 2017.
52 Madigan S, Racine N, Cooke JE, *et al*. COVID-19 and telemental health: Benefits, challenges, and future directions. *Canadian Psychology / Psychologie Canadienne* 2021;62:5–11.
53 Swartz L, Hunt X, Bantjes J, *et al*. Mental health symptoms and disorders in Paralympic athletes: A narrative review. *Br J Sports Med* 2019;53:737–40.
54 Quartiroli A, Schinke RJ, Giffin C, *et al*. Cultural competence in a multinational group of sport psychology professionals. *J Appl Sport Psychol* 2023;35:919–40.
55 Ryba TV, Schinke RJ, Quartiroli A, *et al*. ISSP position stand on cultural praxis in sport psychology: Reaffirming our commitments to the ethics of difference, cultural inclusion, and social justice. *Int J Sport Exerc Psychol* 2024;22:533–52.
56 Elvis Hagan J, Schack T, Schinke R. Sport psychology practice in Africa: Do culture-specific religion and spirituality matter? *ASSRJ* 2019;6:183–97.
57 Ikulayo PB, Semidara JA. Culturally informed sport psychology practice: Nigeria in perspective. *J Clin Sport Psychol* 2011;5:339–49.
58 Stillman MA, Manocchio AM, Glick ID. Combining pharmacotherapy with psychotherapeutic management for the treatment of psychiatric disorders among athletes. *Sports Psychiatry* 2024;3:117–23.
59 Bauer M, Pfennig A, Linden M, *et al*. Efficacy of an algorithm-guided treatment compared with treatment as usual: A randomized, controlled study of inpatients with depression. *J Clin Psychopharmacol* 2009;29:327–33.
60 Trivedi MH, Rush AJ, Crismon ML, *et al*. Clinical results for patients with major depressive disorder in the Texas Medication Algorithm Project. *Arch Gen Psychiatry* 2004;61:669–80.
61 Adli M, Wiethoff K, Baghai TC, *et al*. How effective Is algorithm-guided treatment for depressed inpatients? Results from the randomized controlled multicenter German Algorithm Project 3 trial. *Int J Neuropsychopharmacol* 2017;20:721–30.
62 Adli M, Bauer M, Rush AJ. Algorithms and collaborative-care systems for depression: Are they effective and why? A systematic review. *Biol Psychiatry* 2006;59:1029–38.
63 Reardon CL, Claussen MC, Creado S, *et al*. Psychiatric treatment preferences of sports psychiatrists for athlete-patients across competitive levels: A 2024 update. *Phys Sportsmed* 2025;53:406–16.
64 WADA. The Prohibited List: World Anti-Doping Agency. 2025. Available: https://www.wada-ama.org/en/prohibited-list/ [Accessed 4 Jun 2025].
65 WADA. Therapeutic use exemptions. 2025. Available: https://www.wada-ama.org/en/athletes-support-personnel/therapeutic-use-exemptions-tues
66 Hirschbeck A, Leao DS, Wagner E, *et al*. Psychiatric medication and physical performance parameters - Are there implications for treatment? *Front Psychiatry* 2022;13:985983.
67 Halsey ES, Evans DP. CDC yellow book: Health information for international travel, Available: https://www.incb.org/incb/en/travellers/index.html
68 Portes MAM, Werle I, Bertoglio LJ. Is there a place for psychedelics in sports practice? *Acta Neuropsychiatr* 2025;37:e55.
69 Pinheiro L, Verhagen E, Ocarino J, *et al*. Periodic health evaluation in Para athletes: A position statement based on expert consensus. *BMJ Open Sport Exerc Med* 2024;10:e001946.
70 Fedoruk MN, Teah CS. Unique anti-doping and therapeutic use exemption considerations for the Paralympic athlete. *Aspetar Sports Med J* 2019;7:180–5.
71 Baron DA, Reardon CL, Baron SH. Clinical Sports Psychiatry: An International Perspective. John Wiley & Sons, 2013:132–46.Available: https://onlinelibrary.wiley.com/doi/book/10.1002/9781118404904
72 Edwards CD. Mental health considerations for athlete removal from play and return to play planning. *Sports Psychiatry* 2024;3:125–39.
73 De Souza MJ, Nattiv A, Joy E, *et al*. 2014 Female athlete triad coalition consensus statement on treatment and return to play of the female athlete triad. *Clin J Sport Med* 2014;24:96–119.
74 Joy E, Kussman A, Nattiv A. 2016 update on eating disorders in athletes: A comprehensive narrative review with a focus on clinical assessment and management. *Br J Sports Med* 2016;50:154–62.
75 Wells KR, Jeacocke NA, Appaneal R, *et al*. The Australian Institute of Sport (AIS) and National Eating Disorders Collaboration (NEDC) position statement on disordered eating in high performance sport. *Br J Sports Med* 2020;54:1247–58.
76 Sundgot-Borgen J, Meyer NL, Lohman TG, *et al*. How to minimise the health risks to athletes who compete in weight-sensitive sports review and position statement on behalf of the Ad Hoc Research Working Group on Body Composition, Health and Performance, under the auspices of the IOC Medical Commission. *Br J Sports Med* 2013;47:1012–22.
77 Herring SA, Putukian M, Kibler WB, *et al*. Team physician consensus statement: Return to sport/return to play and the team physician: A team physician consensus statement—2023 update. *Med Sci Sports Exerc* 2024;56:767–75.
78 Dijkstra HP, Pollock N, Chakraverty R, *et al*. Return to play in elite sport: A shared decision-making process. *Br J Sports Med* 2017;51:419–20.

Br J Sports Med: first published as 10.1136/bjsports-2025-111514 on 11 August 2026. Downloaded from http://bjsm.bmj.com/ on August 12, 2026 at Universitetsbiblioteket i Tromsø . Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies.

79 Shrier I. Strategic Assessment of Risk and Risk Tolerance (StARRT) framework for return-to-play decision-making. *Br J Sports Med* 2015;49:1311–5.
80 Kroshus E, Wagner J, Wyrick D, *et al*. Wake up call for collegiate athlete sleep: Narrative review and consensus recommendations from the NCAA Interassociation Task Force on Sleep and Wellness. *Br J Sports Med* 2019;53:731–6.
81 Walsh NP, Halson SL, Sargent C, *et al*. Sleep and the athlete: narrative review and 2021 expert consensus recommendations. *Br J Sports Med* 2021;55:356–68.
82 Leeder J, Glaister M, Pizzoferro K, *et al*. Sleep duration and quality in elite athletes measured using wristwatch actigraphy. *J Sports Sci* 2012;30:541–5.
83 Lastella M, Roach GD, Halson SL, *et al*. Sleep/wake behaviours of elite athletes from individual and team sports. *Eur J Sport Sci* 2015;15:94–100.
84 Sargent C, Lastella M, Halson SL, *et al*. The impact of training schedules on the sleep and fatigue of elite athletes. *Chronobiol Int* 2014;31:1160–8.
85 Turner RW II, Vissa K, Hall C, *et al*. Sleep problems are associated with academic performance in a national sample of collegiate athletes. *J Am Coll Health* 2021;69:74–81.
86 Swinbourne R, Gill N, Vaile J, *et al*. Prevalence of poor sleep quality, sleepiness and obstructive sleep apnoea risk factors in athletes. *Eur J Sport Sci* 2016;16:850–8.
87 Tsunoda K, Mutsuzaki H, Hotta K, *et al*. Correlation between sleep and psychological mood states in female wheelchair basketball players on a Japanese national team. *J Phys Ther Sci* 2017;29:1497–501.
88 Hoshikawa M, Uchida S, Hirano Y. A subjective assessment of the prevalence and factors associated with poor sleep quality amongst elite Japanese athletes. *Sports Med - Open* 2018;4:10.
89 Mah CD, Kezirian EJ, Marcello BM, *et al*. Poor sleep quality and insufficient sleep of a collegiate student-athlete population. *Sleep Health* 2018;4:251–7.
90 Drew M, Vlahovich N, Hughes D, *et al*. Prevalence of illness, poor mental health and sleep quality and low energy availability prior to the 2016 Summer Olympic Games. *Br J Sports Med* 2018;52:47–53.
91 Drew MK, Vlahovich N, Hughes D, *et al*. A multifactorial evaluation of illness risk factors in athletes preparing for the Summer Olympic Games. *J Sci Med Sport* 2017;20:745–50.
92 Badenhorst M, Brown J, Runciman P, *et al*. Promotion of Para Athlete Well-being (PROPEL II): A cross-sectional study of sleep in Para athletes across two nations. *Am J Phys Med Rehabil* 2024;103:261–70.
93 Silva A, Queiroz SS, Winckler C, *et al*. Sleep quality evaluation, chronotype, sleepiness and anxiety of Paralympic Brazilian athletes: Beijing 2008 Paralympic Games. *Br J Sports Med* 2012;46:150–4.
94 Silva A, Pinto Pinheiro LS, Silva S, *et al*. Sleep in Paralympic athletes and its relationship with injuries and illnesses. *Phys Ther Sport* 2022;56:24–31.
95 Fagher K, Dahlström Ö, Lexell J. Mental health, sleep, and pain in elite Para athletes and the association with injury and illness-A prospective study. *PM R* 2023;15:1130–9.
96 Gupta L, Morgan K, Gilchrist S. Does elite sport degrade sleep quality? A systematic review. *Sports Med* 2017;47:1317–33.
97 Khalladi K, Farooq A, Souissi S, *et al*. Inter-relationship between sleep quality, insomnia and sleep disorders in professional soccer players. *BMJ Open Sport Exerc Med* 2019;5:e000498.
98 Dobrosielski DA, Nichols D, Ford J, *et al*. Estimating the prevalence of sleep-disordered breathing among collegiate football players. *Respir Care* 2016;61:1144–50.
99 Tuomilehto H, Vuorinen V-P, Penttilä E, *et al*. Sleep of professional athletes: Underexploited potential to improve health and performance. *J Sports Sci* 2017;35:704–10.
100 Dunican IC, Walsh J, Higgins CC, *et al*. Prevalence of sleep disorders and sleep problems in an elite super rugby union team. *J Sports Sci* 2019;37:950–7.
101 Fagundes SBR, Fagundes DJL, Luna AA, *et al*. Prevalence of restless legs syndrome in runners. *Sleep Med* 2012;13:771.
102 Heller HC, Herzog E, Brager A, *et al*. The negative effects of travel on student athletes through sleep and circadian disruption. *J Biol Rhythms* 2024;39:5–19.
103 Crowley SJ, Acebo C, Carskadon MA. Sleep, circadian rhythms, and delayed phase in adolescence. *Sleep Med* 2007;8:602–12.
104 Schmidt C, Peigneux P, Cajochen C, *et al*. Adapting test timing to the sleep-wake schedule: Effects on diurnal neurobehavioral performance changes in young evening and older morning chronotypes. *Chronobiol Int* 2012;29:482–90.
105 Roenneberg T, Kuehnle T, Pramstaller PP, *et al*. A marker for the end of adolescence. *Curr Biol* 2004;14:R1038–9.
106 Roenneberg T, Kuehnle T, Juda M, *et al*. Epidemiology of the human circadian clock. *Sleep Med Rev* 2007;11:429–38.
107 Charest J, Grandner MA. Sleep and athletic performance: Impacts on physical performance, mental performance, injury risk and recovery, and mental health: An update. *Sleep Med Clin* 2022;17:263–82.
108 Meltzer LJ, Mindell JA. Sleep and sleep disorders in children and adolescents. *Psychiatric Clinics of North America* 2006;29:1059–76.
109 Meltzer LJ, Johnson C, Crosette J, *et al*. Prevalence of diagnosed sleep disorders in pediatric primary care practices. *Pediatrics* 2010;125:e1410–8.
110 Singh S, Kaur H, Singh S, *et al*. Parasomnias: A comprehensive review. *Cureus* 2018;10:e3807.
111 Ohayon MM, Duhoux S, Grieco J, *et al*. Prevalence and incidence of narcolepsy symptoms in the US general population. *Sleep Med X* 2023;6:100095.
112 American Academy of Sleep Medicine. The AASM International Classification of Sleep Disorders – Third Edition, Text Revision (ICSD-3-TR). Darien, IL: ASSM Press, 2023. Available: https://aasm.org/clinical-resources/international-classification-sleep-disorders/
113 Edinger JD, Bonnet MH, Bootzin RR, *et al*. Derivation of research diagnostic criteria for insomnia: Report of an American Academy of Sleep Medicine Work Group. *Sleep* 2004;27:1567–96.
114 Gossard TR, Trotti LM, Videnovic A, *et al*. Restless legs syndrome: Contemporary diagnosis and treatment. *Neurotherapeutics* 2021;18:140–55.
115 Picchietti MA, Picchietti DL. Advances in pediatric restless legs syndrome: Iron, genetics, diagnosis and treatment. *Sleep Med* 2010;11:643–51.
116 Grandner MA, Winkelman JW. Nocturnal leg cramps: Prevalence and associations with demographics, sleep disturbance symptoms, medical conditions, and cardiometabolic risk factors. *PLoS ONE* 2017;12:e0178465.
117 Salas RE, Rasquinha R, Gamaldo CE. All the wrong moves: A clinical review of restless legs syndrome, periodic limb movements of sleep and wake, and periodic limb movement disorder. *Clin Chest Med* 2010;31:383–95.
118 Reid KJ, Zee PC. Circadian rhythm disorders. *Semin Neurol* 2009;29:393–405.
119 Zhu L, Zee PC. Circadian rhythm sleep disorders. *Neurol Clin* 2012;30:1167–91.
120 Ruoff C, Rye D. The ICSD-3 and DSM-5 guidelines for diagnosing narcolepsy: Clinical relevance and practicality. *Curr Med Res Opin* 2016;32:1611–22.
121 Krahn LE, Arand DL, Avidan AY, *et al*. Recommended protocols for the Multiple Sleep Latency Test and Maintenance of Wakefulness Test in adults: Guidance from the American Academy of Sleep Medicine. *J Clin Sleep Med* 2021;17:2489–98.
122 Brauer AA, Athey AB, Ross MJ, *et al*. Sleep and health among collegiate student athletes. *Chest* 2019;156:1234–45.
123 Charest J, Bastien CH, Ellis JG, *et al*. The impact of perceived sleep, mood and alcohol use on verbal, physical and sexual assault experiences among student athletes and student non-athletes. *Int J Environ Res Public Health* 2021;18:2883.
124 Grandner MA, Hall C, Jaszewski A, *et al*. Mental health in student athletes: Associations with sleep duration, sleep quality, insomnia, fatigue, and sleep apnea symptoms. *Athl Train Sports Health Care* 2021;13:e159–67.
125 Khader WS, Tubbs AS, Haghighi A, *et al*. Onset insomnia and insufficient sleep duration are associated with suicide ideation in university students and athletes. *J Affect Disord* 2020;274:1161–4.
126 Kong Y, Yu B, Guan G, *et al*. Effects of sleep deprivation on sports performance and perceived exertion in athletes and non-athletes: A systematic review and meta-analysis. *Front Physiol* 2025;16:1544286.
127 Messman BA, Wiley JF, Feldman E, *et al*. Irregular sleep is linked to poorer mental health: A pooled analysis of eight studies. *Sleep Health* 2024;10:493–9.
128 Huang T, Mariani S, Redline S. Sleep irregularity and risk of cardiovascular events: The multi-ethnic study of atherosclerosis. *J Am Coll Cardiol* 2020;75:991–9.
129 Zuraikat FM, Makarem N, Redline S, *et al*. Sleep regularity and cardiometabolic heath: Is variability in sleep patterns a risk factor for excess adiposity and glycemic dysregulation? *Curr Diab Rep* 2020;20:38.
130 Healy KL, Morris AR, Liu AC. Circadian synchrony: Sleep, nutrition, and physical activity. *Front Netw Physiol* 2021;1:732243.
131 Galinde AA, Al-Mughales F, Oster H, *et al*. Different levels of circadian (de)synchrony -- Where does it hurt? *F1000Res* 2022;11:1323.
132 Raikes AC, Athey A, Alfonso-Miller P, *et al*. Insomnia and daytime sleepiness: Risk factors for sports-related concussion. *Sleep Med* 2019;58:66–74.
133 Hertenstein E, Feige B, Gmeiner T, *et al*. Insomnia as a predictor of mental disorders: A systematic review and meta-analysis. *Sleep Med Rev* 2019;43:96–105.
134 Baglioni C, Riemann D. Is chronic insomnia a precursor to major depression? Epidemiological and biological findings. *Curr Psychiatry Rep* 2012;14:511–8.
135 Baglioni C, Battagliese G, Feige B, *et al*. Insomnia as a predictor of depression: A meta-analytic evaluation of longitudinal epidemiological studies. *J Affect Disord* 2011;135:10–9.
136 Dolsen EA, Prather AA, Lamers F, *et al*. Suicidal ideation and suicide attempts: Associations with sleep duration, insomnia, and inflammation. *Psychol Med* 2021;51:2094–103.
137 Perlis ML, Pigeon WR, Grandner MA, *et al*. Why treat insomnia? *J Prim Care Community Health* 2021;12:21501327211014084.
138 Perlis ML, Grandner MA, Chakravorty S, *et al*. Suicide and sleep: Is it a bad thing to be awake when reason sleeps? *Sleep Med Rev* 2016;29:101–7.
139 Tubbs AS, Taneja K, Ghani SB, *et al*. Sleep continuity, timing, quality, and disorder are associated with suicidal ideation and suicide attempts among college students. *J Am Coll Health* 2024;72:3357–65.
140 Lorenzi-Filho G, Drager LF. Is the cardiovascular system the primary target of obstructive sleep apnea? *Am J Respir Crit Care Med* 2008;178:892–3.
141 Bakker JP, Sharma B, Malhotra A. Obstructive sleep apnea: The elephant in the cardiovascular room. *Chest* 2012;141:580–1.
142 Selim B, Won C, Yaggi HK. Cardiovascular consequences of sleep apnea. *Clin Chest Med* 2010;31:203–20.
143 Aloia MS, Arnedt JT, Smith L, *et al*. Examining the construct of depression in obstructive sleep apnea syndrome. *Sleep Med* 2005;6:115–21.

Br J Sports Med: first published as 10.1136/bjsports-2025-111514 on 11 August 2026. Downloaded from http://bjsm.bmj.com/ on August 12, 2026 at Universitetsbiblioteket i Tromsoe . Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies.

144 Pandi-Perumal SR, Monti JM, Burman D, *et al*. Clarifying the role of sleep in depression: A narrative review. *Psychiatry Res* 2020;291:113239.
145 Malhotra RK. Sleep, recovery, and performance in sports. *Neurol Clin* 2017;35:547–57.
146 Martínez Revuelta L, Flores-Fraile J, Zubizarreta-Macho Á, *et al*. Relationship between obstructive sleep apnea and sports: Systematic review and meta-analysis. *J Clin Med* 2024;13:6814.
147 Howarth NE, White AJ, Pearce AJ, *et al*. Obstructive sport apnea (OSA) and contact sports: A systematic review and meta-analysis. *Sleep Epidemiology* 2022;2:100036.
148 Adams JW, Alosco ML, Mez J, *et al*. Association of probable REM sleep behavior disorder with pathology and years of contact sports play in chronic traumatic encephalopathy. *Acta Neuropathol* 2020;140:851–62.
149 Grandner MA, Mills J, Clarke M, *et al*. Sleep and circadian health promotion programs for athletes. In: *Sleep and Sport*. Elsevier, 2024: 313–26.
150 Grandner MA, Lujan MR, Ghani SB. Sleep-tracking technology in scientific research: Looking to the future. *Sleep* 2021;44:zsab071.
151 Lujan MR, Perez-Pozuelo I, Grandner MA. Past, present, and future of multisensory wearable technology to monitor sleep and circadian rhythms. *Front Digit Health* 2021;3:721919.
152 Sargent C, Lastella M, Halson SL, *et al*. The validity of activity monitors for measuring sleep in elite athletes. *J Sci Med Sport* 2016;19:848–53.
153 Wanberg LJ, Thomas SJ, Reid M, *et al*. Adapting brief behavioral treatment for insomnia for former National Football League players: A pilot study. *Behav Sleep Med* 2023;21:273–90.
154 Ludwig R, Rippee M, D'Silva L, *et al*. Cognitive behavioral therapy for insomnia improves sleep outcomes in individuals with concussion: A preliminary randomized wait-list control study. *J Head Trauma Rehabil* 2024;39:318–27.
155 Simpson NS, Gibbs EL, Matheson GO. Optimizing sleep to maximize performance: Implications and recommendations for elite athletes. *Scandinavian Med Sci Sports* 2017;27:266–74.
156 Perlis ML, Smith MT, Benson-Jungquist C, *et al*. *Cognitive Behavioral Treatment of Insomnia: A Session-by-Session Guide*. Springer, 2005.
157 Qaseem A, Kansagara D, Forciea MA, *et al*. Management of chronic insomnia disorder in adults: A clinical practice guideline from the American College of Physicians. *Ann Intern Med* 2016;165:125–33.
158 Edinger JD, Arnedt JT, Bertisch SM, *et al*. Behavioral and psychological treatments for chronic insomnia disorder in adults: An American Academy of Sleep Medicine clinical practice guideline. *J Clin Sleep Med* 2021;17:255–62.
159 Gorovoy SB, Campbell RL, Fox RS, *et al*. App-supported sleep coaching: Implications for sleep duration and sleep quality. *Front Sleep* 2023;2:1156844.
160 Reardon CL, Creado S. Psychiatric medication preferences of sports psychiatrists. *Phys Sportsmed* 2016;44:397–402.
161 Celorrio San Miguel AM, Roche E, Herranz-López M, *et al*. Impact of melatonin supplementation on sports performance and circulating biomarkers in highly trained athletes: A systematic review of randomized controlled trials. *Nutrients* 2024;16:1011.
162 Halson SL. Sleep in elite athletes and nutritional interventions to enhance sleep. *Sports Med* 2014;44:13–23.
163 Sletten TL, Magee M, Murray JM, *et al*. Efficacy of melatonin with behavioural sleep-wake scheduling for delayed sleep-wake phase disorder: A double-blind, randomised clinical trial. *PLoS Med* 2018;15:e1002587.
164 Baird MB, Asif IM. Medications for sleep schedule adjustments in athletes. *Sports Health* 2018;10:35–9.
165 Leonardo-Mendonça RC, Martinez-Nicolas A, de Teresa Galván C, *et al*. The benefits of four weeks of melatonin treatment on circadian patterns in resistance-trained athletes. *Chronobiol Int* 2015;32:1125–34.
166 Choi K, Lee YJ, Park S, *et al*. Efficacy of melatonin for chronic insomnia: Systematic reviews and meta-analyses. *Sleep Med Rev* 2022;66:101692.
167 Sateia MJ, Buysse DJ, Krystal AD, *et al*. Clinical practice guideline for the pharmacologic treatment of chronic insomnia in adults: An American Academy of Sleep Medicine clinical practice guideline. *J Clin Sleep Med* 2017;13:307–49.
168 Patil SP, Ayappa IA, Caples SM, *et al*. Treatment of adult obstructive sleep apnea with positive airway pressure: An American Academy of Sleep Medicine clinical practice guideline. *J Clin Sleep Med* 2019;15:335–43.
169 Ilea A, Timuș D, Höpken J, *et al*. Oral appliance therapy in obstructive sleep apnea and snoring - Systematic review and new directions of development. *CRANIO®* 2021;39:472–83.
170 Ramar K, Dort LC, Katz SG, *et al*. Clinical practice guideline for the treatment of obstructive sleep apnea and snoring with oral appliance therapy: An update for 2015. *J Clin Sleep Med* 2015;11:773–827.
171 Auger RR, Burgess HJ, Emens JS, *et al*. Clinical practice guideline for the treatment of intrinsic circadian rhythm sleep-wake disorders: Advanced Sleep-Wake Phase Disorder (ASWPD), Delayed Sleep-Wake Phase Disorder (DSWPD), Non-24-Hour Sleep-Wake Rhythm Disorder (N24SWD), and Irregular Sleep-Wake Rhythm Disorder (ISWRD). An Update for 2015. *J Clin Sleep Med* 2015;11:1199–236.
172 Bassetti CLA, Kallweit U, Vignatelli L, *et al*. European guideline and expert statements on the management of narcolepsy in adults and children. *J Sleep Res* 2021;30:e13387.
173 Maski K, Trotti LM, Kotagal S, *et al*. Treatment of central disorders of hypersomnolence: An American Academy of Sleep Medicine clinical practice guideline. *J Clin Sleep Med* 2021;17:1881–93.
174 Maski K, Trotti LM, Kotagal S, *et al*. Treatment of central disorders of hypersomnolence: an American Academy of Sleep Medicine systematic review, meta-analysis, and GRADE assessment. *J Clin Sleep Med* 2021;17:1895–945.
175 Nixdorf R, Madigan DJ, Kenttä G, *et al*. Depression, athlete burnout, and overtraining: A review of similarities and differences. In: *Routledge Handbook of Mental Health in Elite Sport*. 2023: 188–200.
176 American Psychiatric Association. *Diagnostic and Statistical Manual of Mental Disorders, Text Revision (DSM-5-TR)*. 5th edn. Washington, DC: American Psychiatric Association, 2022. Available: https://doi.org/10.1176/appi.books.9780890425787
177 Currie A, Rogers T. Helping the athlete with depression. In: *Sport and Mental Health: From Research to Everyday Practice*. Springer, 2023: 169–93.
178 Currie A, Blauwet C, Bindra A, *et al*. Athlete mental health: Future directions. *Br J Sports Med* 2021;55:1243–4.
179 Tahtinen RE, Shelley J, Morris R. Gaining perspectives: A scoping review of research assessing depressive symptoms in athletes. *Psychol Sport Exerc* 2021;54:101905.
180 Ely G, Woodman T, Roberts R, *et al*. A three-year longitudinal study of athlete mental health: A cricket case study. *Scandinavian Med Sci Sports* 2025;35:e70125.
181 McGuire LC, Ingram YM, Sachs ML, *et al*. Temporal changes in depression symptoms in male and female collegiate student-athletes. *J Clin Sport Psychol* 2017;11:337–51.
182 Hammond T, Gialloreto C, Kubas H, *et al*. The prevalence of failure-based depression among elite athletes. *Clin J Sport Med* 2013;23:273–7.
183 Harenberg S, Ouellet-Pizer C, Nieto M, *et al*. Anonymous vs. non-anonymous administration of depression scales in elite athletes: A meta-analysis. *Int Rev Sport Exerc Psychol* 2024;17:882–902.
184 Golding L, Gillingham RG, Perera NKP. The prevalence of depressive symptoms in high-performance athletes: A systematic review. *Phys Sportsmed* 2020;48:247–58.
185 Gorczynski PF, Coyle M, Gibson K. Depressive symptoms in high-performance athletes and non-athletes: A comparative meta-analysis. *Br J Sports Med* 2017;51:1348–54.
186 Gouttebarge V, Castaldelli-Maia JM, Gorczynski P, *et al*. Occurrence of mental health symptoms and disorders in current and former elite athletes: A systematic review and meta-analysis. *Br J Sports Med* 2019;53:700–6.
187 Schorb A, Aichhorn J, Schiepek G, *et al*. Burnout in adolescent elite athletes: A pilot study on an Austrian student sample. *Sports Psychiatry: Journal of Sports and Exercise Psychiatry* 2023;2:51–6.
188 Madigan DJ, Olsson LF, Hill AP, *et al*. Athlete burnout symptoms are increasing: A cross-temporal meta-analysis of average levels from 1997 to 2019. *J Sport Exerc Psychol* 2022;44:153–68.
189 Meeusen R, Duclos M, Foster C, *et al*. Prevention, diagnosis, and treatment of the overtraining syndrome: Joint consensus statement of the European College of Sport Science and the American College of Sports Medicine. *Med Sci Sports Exerc* 2013;45:186–205.
190 Raglin J, Wilson G. Overtraining and staleness in athletes. In: Hanin Y, ed. *Emotions in Sports*. Champaign, IL: Human Kinetics, 2000: 191–207.
191 Ecklund RE, DeFreese JD. Athlete burnout: What we know, what we could know, and how we can find out more. *IJASS (International Journal of Applied Sports Sciences)* 2015;27:63–75.
192 Goodger K, Gorely T, Lavallee D, *et al*. Burnout in sport: A systematic review. *Sport Psychol* 2007;21:127–51.
193 Soligard T, Schwellnus M, Alonso JM, *et al*. How much is too much. *Br J Sports Med* 2026;50:1030–41.
194 Pensgaard AM, Sundgot-Borgen J, Edwards C, *et al*. Intersection of mental health issues and Relative Energy Deficiency in Sport (REDs): A narrative review by a subgroup of the IOC consensus on REDs. *Br J Sports Med* 2023;57:1127–35.
195 Halioua R, Wasserfurth P, Toepffer D, *et al*. Exploring the relationship between low energy availability, depression and eating disorders in female athletes: A cross-sectional study. *BMJ Open Sport Exerc Med* 2024;10:e002035.
196 Gorczynski P. Major depressive disorder and depressive symptoms. In: Reardon C, ed. *Mental Health Care for Elite Athletes*. 2022: 51–9.
197 Moreira D, Azeredo A, Dias P. Risk factors for gambling disorder: A systematic review. *J Gambl Stud* 2023;39:483–511.
198 Brenner P, Brandt L, Li G, *et al*. Substance use disorders and risk for treatment resistant depression: A population-based, nested case-control study. *Addiction* 2020;115:768–77.
199 Lichtenstein MB, Hinze CJ, Emborg B, *et al*. Compulsive exercise: Links, risks and challenges faced. *Psychol Res Behav Manag* 2017;10:85–95.
200 Stillman MA, Farmer H. *Psychotherapeutic Approaches to Addressing Mental Health Problems Among Elite Athletes*. London: IntechOpen, 2021.
201 Gleeson M. Immune function in sport and exercise. *J Appl Physiol* 2007;103:693–9.
202 Arteaga-Henríquez G, Simon MS, Burger B, *et al*. Low-grade inflammation as a predictor of antidepressant and anti-inflammatory therapy response in MDD patients: A systematic review of the literature in combination with an analysis of experimental data collected in the EU-MOODINFLAME Consortium. *Front Psychiatry* 2019;10:458.

Br J Sports Med: first published as 10.1136/bjsports-2025-111514 on 11 August 2026. Downloaded from http://bjsm.bmj.com/ on August 12, 2026 at Universitetsbiblioteket i Tromsø . Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies.

203 Vreijling SR, Chin Fatt CR, Williams LM, *et al*. Features of immunometabolic depression as predictors of antidepressant treatment outcomes: Pooled analysis of four clinical trials. *Br J Psychiatry* 2024;224:89–97.
204 Gill VS, Sullivan G, Stearns H, *et al*. Mental health in elite athletes: A systematic review of suicidal behaviour as compared to the general population. *Sports Med* 2024;54:1–18.
205 Whelan BM, Kliethermes SA, Schloredt KA, *et al*. Suicide in National Collegiate Athletic Association athletes: A 20-year analysis. *Br J Sports Med* 2024;58:531–7.
206 Anchuri K MHS, Davoren AK PhD, Shanahan A PhD, *et al*. Nonsuicidal self-injury, suicidal ideation, and suicide attempt among collegiate athletes: Findings from the National College Health Assessment. *J Am Coll Health* 2020;68:815–23.
207 Rao AL, Asif IM, Drezner JA, *et al*. Suicide in National Collegiate Athletic Association (NCAA) Athletes: A 9-Year Analysis of the NCAA Resolutions Database. *Sports Health* 2015;7:452–7.
208 Tran A. Looking forward to student-athlete mental health: Racial/ethnic trends from 2010 to 2015. *J Am College Health* 2021;69:942–50.
209 Hemati Alamdarloo G, Sheikhani Shahin H, Gholami M, *et al*. The role of general health in predicting the social harms of elite athletes. *J Study of Sports and Athletes in Education* 2024;18:257–67.
210 Sun G, Zhao J, Tian S, *et al*. Psychological strain and suicidal ideation in athletes: The multiple mediating effects of hopelessness and depression. *IJERPH* 2020;17:8087.
211 Pichler E, Ewers S, Ajdacic-Gross V, *et al*. Athletes are not at greater risk for death by suicide: A review. *Scandinavian Med Sci Sports* 2023;33:569–85.
212 Gorczynski P, McCabe T. Suicide in elite sport. *Advances in Psychiatry and Behavioral Health* 2024;4:33–9.
213 Shaman J, Kandula S, Pei S, *et al*. Quantifying suicide contagion at population scale. *Sci Adv* 2024;10:eadq4074.
214 Calvo S, Carrasco JP, Conde-Pumpido C, *et al*. Does suicide contagion (Werther effect) take place in response to social media? A systematic review. *Span J Psychiatry Ment Health* 2026;19:55–65.
215 Ivey-Stephenson AZ, Ballesteros MF, Trinh E, *et al*. CDC guidance for community response to suicide clusters, United States, 2024. *MMWR Suppl* 2024;73:17–26.
216 Ramamurthy C, Fraser T, Krysinska K, *et al*. Effectiveness of suicide postvention service models and guidelines 2014-2024: A scoping review. *Prev Med* 2025;195:108279.
217 Badenhorst M, Runciman P, Brown JC, *et al*. Promotion of Para athlete well-being in South Africa (the PROPEL studies): Profiles and prevalence of psychological distress. *J Sci Med Sport* 2021;24:616–21.
218 Bentzen M, Kenttä G, Karls T, *et al*. Monitoring mental distress in Para athletes in preparation, during and after the Beijing Paralympic Games 2022: A 22 week prospective mixed-method study. *Front Sports Act Living* 2022;4:945073.
219 Junge A, Wellmann K, Zech A. Prevalence of symptoms of depression and generalized anxiety disorder in field hockey players of German first and lower leagues. *Scandinavian Med Sci Sports* 2023;33:1866–73.
220 Mountjoy M, Edwards C, Cheung CP, *et al*. Implementation of the International Olympic Committee Sport Mental Health Assessment Tool 1: Screening for mental health symptoms in a Canadian multisport university program. *Clin J Sport Med* 2023;33:5–12.
221 Gulliver A, Griffiths KM, Mackinnon A, *et al*. The mental health of Australian elite athletes. *J Sci Med Sport* 2015;18:255–61.
222 Kuettel A, Durand-Bush N, Larsen CH. Mental health profiles of Danish youth soccer players: The influence of gender and career development. *J Clin Sport Psychol* 2022;16:276–93.
223 Lima Y, Denerel N, Devran S, *et al*. Which athletes are more vulnerable to mental health symptoms during the COVID-19 crisis? A cross-sectional study. *Research in Sports Medicine* 2024;32:248–59.
224 Sanfilippo JL, Haralsdottir K, Watson AM. Anxiety and depression prevalence in incoming Division I collegiate athletes from 2017 to 2021. *Sports Health: A Multidisciplinary Approach* 2024;16:776–82.
225 Tahtinen RE, Kristjansdottir H. The influence of anxiety and depression symptoms on help-seeking intentions in individual sport athletes and non-athletes: The role of gender and athlete status. *J Clin Sport Psychol* 2019;13:134–51.
226 Tran A. In or out of the game? Counter-stereotype paradoxes and Asian-identified student-athlete mental health. *Cultural Diversity & Ethnic Minority Psychology* 2021;27:579–92.
227 Andersson MJ, Kenttä G, Claesdotter-Knutsson E, *et al*. Mental health symptom burden in elite ice hockey players and its association with self-reported concussive events. *BMC Sports Sci Med Rehabil* 2024;16:197.
228 Kraus AD, Tibbetts E. Depression, anxiety, and help-seeking among NCAA Division III athletes at a historically women's college. *J Clin Sport Psychol* 2022;16:1–22.
229 Ito T, Fetters MD, Kipps C, *et al*. Depressive symptoms among male professional soccer players in Japan. *Asian J Sport Exercise Psychology* 2023;3:154–60.
230 Junge A, Feddermann-Demont N. Prevalence of depression and anxiety in top-level male and female football players. *BMJ Open Sport Exerc Med* 2016;2:e000087.
231 Junge A, Prinz B. Depression and anxiety symptoms in 17 teams of female football players including 10 German first league teams. *Br J Sports Med* 2019;53:471–7.
232 Kurnellas R, Ward RM, Taylor E, *et al*. Alcohol demand in college students: The roles of athletic involvement and gender. *Exp Clin Psychopharmacol* 2024;32:54–67.
233 Poucher ZA, Tamminen KA, Sabiston CM, *et al*. A longitudinal examination of changes in mental health among elite Canadian athletes. *J Sports Sci* 2022;40:733–41.
234 Salles JI, Silva C, Wolff A, *et al*. Anxiety, insomnia, and depression during COVID-19 lockdown in elite individual and team sport athletes. *Psicol Reflex Crit* 2022;35:33.
235 Scheid JL, Basile S, West SL. Low energy availability risk Is associated with anxiety in female collegiate athletes. *Sports* 2014;12:269.
236 Zhao X, Liu Z, Zhao L, *et al*. Post-traumatic stress disorder and mental health in Chinese college athletes during and after the COVID-19 pandemic: The multiple mediating effects of basic psychological needs and generalized anxiety disorder. *Behav Sci (Basel)* 2023;13:567.
237 Li C, Fan R, Sun J, *et al*. Risk and protective factors of generalized anxiety disorder of elite collegiate athletes: A cross-sectional study. *Front Public Health* 2021;9:607800.
238 Liu M, Zhao X, Liu Z. Relationship between psychological distress, basic psychological needs, anxiety, mental pressure, and athletic burnout of Chinese college football ahletes during the COVID-19 pandemic. *Sustainability* 2022;14:7100.
239 Poucher ZA, Tamminen KA, Sabiston CM, *et al*. Prevalence of symptoms of common mental disorders among elite Canadian athletes. *Psychol Sport Exerc* 2021;57:102018.
240 Roberts SD, Champigny C, Feldman SJ, *et al*. Screening for anxiety and depression symptoms using concussion symptom scales among varsity athletes. *Clin J Sport Med* 2024;34:105–11.
241 Simon MB, Barczak-Scarboro NE, Pexa B, *et al*. Collegiate student-athlete mental health symptom trajectories across a competitive season: Do training load and resilience matter? *Int J Sport Exerc Psychol* 2025;23:501–19.
242 Schaal K, Tafflet M, Nassif H, *et al*. Psychological balance in high level athletes: Gender-based differences and sport-specific patterns. *PLoS ONE* 2011;6:e19007.
243 Bonet L, Benito A, Usó H, *et al*. Mental health in first- and second-division soccer players: A cross-sectional study. *Sports* 2024;12:106.
244 Fiorilli G, Iuliano E, Aquino G, *et al*. Mental health and social participation skills of wheelchair basketball players: A controlled study. *Res Dev Disabil* 2013;34:3679–85.
245 Yildiz K, Pepe O, Saritaş N, *et al*. An investigation over the levels of social anxiety according to the gender factors of hearing-impaired adolescent athletes between the ages of 12-16. *Nigde University Journal of Physical Education & Sport Sciences / Nigde Üniversitesi Beden Egitimi ve Spor Bilimleri Dergisi* 2012;6:56–61.
246 Storch EA, Storch JB, Killiany EM, *et al*. Self-reported psychopathology in athletes: A comparison of intercollegiate student-athletes and non-athletes. *J Sport Behav* 2005;28:86–98.
247 Reardon CL, Gorczynski P, Hainline B, *et al*. Anxiety disorders in athletes: A clinical review. *Adv Psychiatry Behavioral Health* 2021;1:149–60.
248 Gay JL, Monsma EV, Torres-McGehee TM. Developmental and contextual risks of social physique anxiety among female athletes. *Res Q Exerc Sport* 2011;82:168–77.
249 Haase AM. Weight perception in female athletes: Associations with disordered eating correlates and behavior. *Eat Behav* 2011;12:64–7.
250 Kosmidou E, Giannitsopoulou E, Kountouratzi N, *et al*. Maltreatment (psychological, physical), social physique, anxiety, body dissatisfaction and drive for thinness in Greek female athletes (rhythmic gymnasts and other sports) after dropout. *SGJ* 2022;14:271–84.
251 Kosmidou E, Giannitsopoulou E, Moysidou D. Social physique anxiety and pressure to be thin in adolescent ballet dancers, rhythmic gymnastics and swimming athletes. *Research in Dance Education* 2017;18:23–33.
252 Krane V, Stiles-Shipley JA, Waldron J, *et al*. Relationships among body satisfaction, social physique anxiety, and eating behaviors in female athletes and exercisers. *J Sport Behav* 2001;24:247–64.
253 Wolverton J, Dobersek U. Affective forecasting and social physique anxiety among female athletes: A pilot study. *Sport J* 2023;24. Available: https://thesportjournal.org/article/affective-forecasting-and-social-physique-anxiety-among-female-athletes-a-pilot-study
254 Haase AM, Prapavessis H, Glynn Owens R. Perfectionism, social physique anxiety and disordered eating: A comparison of male and female elite athletes. *Psychol Sport Exerc* 2002;3:209–22.
255 Mülazimoğlu-Balli Ö, Koca C, Aşçi F. An examination of social physique anxiety with regard to sex and level of sport involvement. *J Hum Kinet* 2010;26:115–22.
256 Demirel H. Social appearance anxiety and Rosenberg Self-esteem Scores in young physical disabled athletes. *UJER* 2019;7:664–7.
257 Üstün NA. Social physical anxiety and nutritional attitude in athlete and non-athlete university students. *Progress in Nutrition* 2020;22.
258 Ford JL, Ildefonso K, Jones ML, *et al*. Sport-related anxiety: Current insights. *Open Access J Sports Med* 2017;8:205–12.
259 Cromer L, Kaier E, Davis J, *et al*. OCD in college athletes. *Am J Psychiatry* 2017;174:595–7.
260 Oevreboe TH, Ivarsson A, Sundgot-Borgen J, *et al*. Mental health problems in elite sport: The difference in the distribution of mental distress and mental disorders among a sample of Norwegian elite athletes. *BMJ Open Sport Exerc Med* 2023;9:e001538.

Br J Sports Med: first published as 10.1136/bjsports-2025-111514 on 11 August 2026. Downloaded from http://bjsm.bmj.com/ on August 12, 2026 at Universitetsbiblioteket i Tromsoe . Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies.

261 Fawcett EJ, Power H, Fawcett JM. Women are at greater risk of OCD than men: A meta-analytic review of OCD prevalence worldwide. *J Clin Psychiatry* 2020;81:19r13085.
262 Marazziti D, Parra E, Amadori S, *et al*. Obsessive-compulsive and depressive symptoms in professional tennis players. *Clin Neuropsychiatry* 2021;18:304–11.
263 Dömötör Z, Ruíz-Barquín R, Szabo A. Superstitious behavior in sport: A literature review. *Scand J Psychol* 2016;57:368–82.
264 Knecht JC. The prevalence of body dysmorphic disorder in female collegiate athletes. West Virginia University; 2007.
265 Baghurst T, Lirgg C. Characteristics of muscle dysmorphia in male football, weight training, and competitive natural and non-natural bodybuilding samples. *Body Image* 2009;6:221–7.
266 Caner SM, Çevïker A. An investigation on the relationship between social appearance concerns and body perceptions of weightlifting athletes at national team level. *Akdeniz Spor Bilimleri Dergisi* 2023;6:147–58.
267 Hernández-Martínez A, González-Martí I, Contreras Jordán OR. Detection of muscle dysmorphia symptoms in male weightlifters. *AN PSICOL-SPAIN* 2017;33:204.
268 Lantz CD, Rhea DJ, Cornelius AE. Muscle dysmorphia in elite-level power lifters and bodybuilders: A test of differences within a conceptual model. *J Strength Cond Res* 2002;16:649–55.
269 Vardardottir B, Olafsdottir AS, Gudmundsdottir SL. Body dissatisfaction, disordered eating and exercise behaviours: Associations with symptoms of REDs in male and female athletes. *BMJ Open Sport Exerc Med* 2023;9:e001731.
270 Skemp KM, Mikat RP, Schenck KP, *et al*. Muscle dysmorphia: Risk may be influenced by goals of the weightlifter. *J Strength Cond Res* 2013;27:2427–32.
271 DeWolfe CEJ, Watt MC, Galbraith MK, *et al*. Mental health in athletes: Anxiety sensitivity as a transdiagnostic risk factor. *Int J Sport Exerc Psychol* 2025;23:1608–27.
272 Bukhari FK, Fahd S, Tahira R, *et al*. Impact of sports anxiety on sports performance of players. *Pak J Humanit Soc Sci* 2021;9:581–6.
273 Mesagno C, Hammond AA, Goodyear MA. An initial investigation into the mental health difficulties in athletes who experience choking under pressure. *Psychol Sport Exerc* 2024;74:102663.
274 Edwards CD, Aron CM. Obsessive–compulsive disorder in sports–Beyond superstitions. *Advances in Psychiatry and Behavioral Health* 2023;3:43–55.
275 Thomson P, Jaque S. Visiting the muses: Creativity, coping, and PTSD in talented dancer and athletes. *Am J Play* 2016;8:363–78.
276 Aron CM, LeFay SM. Post-traumatic stress disorder and other trauma-related disorders. In: Reardon C, ed. *Mental Health Care for Elite Athletes*. Champaign, IL: Springer, 2022: 69–78.
277 Aron CM, Harvey S, Hainline B, *et al*. Post-traumatic stress disorder (PTSD) and other trauma-related mental disorders in elite athletes: A narrative review. *Br J Sports Med* 2019;53:779–84.
278 Tuakli-Wosornu YA, Sun Q, Gentry M, *et al*. Non-accidental harms ('abuse') in athletes with impairment ('para athletes'): A state-of-the-art review. *Br J Sports Med* 2020;54:129–38.
279 Brassil HE, Salvatore AP. The frequency of post-traumatic stress disorder symptoms in athletes with and without sports related concussion. *Clin Translational Med* 2018;7:25.
280 Thomson P, Jaque SV. Cumulative psychological trauma, emotional regulation, and orthopedic injury in a sample of pre-professional and professional dancers and college athletes. *Med Probl Perform Art* 2020;35:89–95.
281 Yang SX, Cheng S, Su DL. Sports injury and stressor-related disorder in competitive athletes: A systematic review and a new framework. *Burns Trauma* 2022;10:tkac017.
282 O'Neill DF. Injury contagion in Alpine ski racing: The effect of injury on teammates' performance. *J Clin Sport Psychol* 2008;2:278–92.
283 Timpka T, Spreco A, Dahlstrom O, *et al*. Suicidal thoughts (ideation) among elite athletics (track and field) athletes: Associations with sports participation, psychological resourcefulness and having been a victim of sexual and/or physical abuse. *Br J Sports Med* 2021;55:198–205.
284 Wenzel T, Zhu LJ. Posttraumatic stress in athletes. In: *Clinical Sports Psychiatry: An International Perspective*. 2013: 102–14.
285 Roberts AL, Gilman SE, Breslau J, *et al*. Race/ethnic differences in exposure to traumatic events, development of post-traumatic stress disorder, and treatment-seeking for post-traumatic stress disorder in the United States. *Psychol Med* 2011;41:71–83.
286 Steinfeldt JA, Foltz BD, Mungro J, *et al*. Masculinity socialization in sports: Influence of college football coaches. *Psychology of Men & Masculinity* 2011;12:247–59.
287 Curdt A, Eggleston B. EMDR-based interventions for athletic traumas: A case study of two female golfers. *J EMDR Pract and Res* 2023;17:70–82.
288 Ramaeker J, Petrie TA. "Man up!": Exploring intersections of sport participation, masculinity, psychological distress, and help-seeking attitudes and intentions. *Psychology of Men & Masculinity* 2019;20:515–27.
289 McMahon J, McGannon KR. Implementing an organizational trauma-informed approach to Olympic sites: An urgent priority to protect elite athlete well-being during Olympic games participation. *Am Behavioral Scientist* 2024;00027642241261253.
290 Courtois CA, Sonis J, Brown L, *et al*. Clinical practice guideline for the treatment of posttraumatic stress disorder (PTSD) in adults. Washington DC American Psychological Association; 2017.
291 Ursano RJ, Bell C, Eth S, *et al*. Practice guideline for the treatment of patients with acute stress disorder and posttraumatic stress disorder. *Am J Psychiatry* 2004;161:3–31.
292 Reynoso-Sánchez LF, Hoyos-Flores JR. A single-session eye movement desensitization and reprocessing (EMDR) therapy reduces anxiety and improves self-confidence in athletes with post-traumatic stress associated with injury. *Int J Sport Stud Hlth* 2022;5:1–9.
293 Mountjoy M, Ackerman KE, Bailey DM, *et al*. 2023 International Olympic Committee's (IOC) consensus statement on Relative Energy Deficiency in Sport (REDs). *Br J Sports Med* 2023;57:1073–98.
294 Karrer Y, Halioua R, Mötteli S, *et al*. Disordered eating and eating disorders in male elite athletes: A scoping review. *BMJ Open Sport Exerc Med* 2020;6:e000801.
295 Sundgot-Borgen J, Torstveit MK. Prevalence of eating disorders in elite athletes is higher than in the general population. *Clin J Sport Med* 2004;14:25–32.
296 Torstveit MK, Rosenvinge JH, Sundgot-Borgen J. Prevalence of eating disorders and the predictive power of risk models in female elite athletes: A controlled study. *Scand J Med Sci Sports* 2008;18:108–18.
297 Martinsen M, Sundgot-borgen J. Higher prevalence of eating disorders among adolescent elite athletes than controls. *Med Sci Sports Exerc* 2013;45:1188–97.
298 Lichtenstein MB, Johansen KK, Runge E, *et al*. Behind the athletic body: A clinical interview study of identification of eating disorder symptoms and diagnoses in elite athletes. *BMJ Open Sport Exerc Med* 2022;8:e001265.
299 Brook EM, Tenforde AS, Broad EM, *et al*. Low energy availability, menstrual dysfunction, and impaired bone health: A survey of elite para athletes. *Scand J Med Sci Sports* 2019;29:678–85.
300 Condo D, Lohman R, Kelly M, *et al*. Nutritional intake, sports nutrition knowledge and energy availability in female Australian rules football players. *Nutrients* 2019;11:971.
301 Staal S, Sjödin A, Fahrenholtz I, *et al*. Low RMR_{ratio} as a surrogate marker for energy deficiency, the choice of predictive equation vital for correctly identifying male and female ballet dancers at risk. *Int J Sport Nutr Exerc Metab* 2018;28:ijsnem.2017-0327:412–8:.
302 Sygo J, Coates AM, Sesbreno E, *et al*. Prevalence of indicators of low energy availability in elite female sprinters. *Int J Sport Nutr Exerc Metab* 2018;28:490–6.
303 Jesus F, Castela I, Silva AM, *et al*. Risk of low energy availability among female and male elite runners competing at the 26th European Cross-Country Championships. *Nutrients* 2021;13:873.
304 McCormack WP, Shoepe TC, LaBrie J, *et al*. Bone mineral density, energy availability, and dietary restraint in collegiate cross-country runners and non-running controls. *Eur J Appl Physiol* 2019;119:1747–56.
305 Dasa MS, Friborg O, Kristoffersen M, *et al*. Risk and prevalence of Relative Energy Deficiency in Sport (REDs) among professional female football players. *Eur J Sport Sci* 2024;24:1032–41.
306 Dambacher L, Pritchett K, Pritchett R, *et al*. Risk of low energy availability, disordered eating, and menstrual dysfunction in female collegiate runners. *J Athl Train* 2025;60:177–84.
307 Keay N, Overseas A, Francis G. Indicators and correlates of low energy availability in male and female dancers. *BMJ Open Sport Exerc Med* 2020;6:e000906.
308 Moris JM, Olendorff SA, Zajac CM, *et al*. Collegiate male athletes exhibit conditions of the Male Athlete Triad. *Appl Physiol Nutr Metab* 2022;47:328–36.
309 Keay N, Francis G, Hind K. Low energy availability assessed by a sport-specific questionnaire and clinical interview indicative of bone health, endocrine profile and cycling performance in competitive male cyclists. *BMJ Open Sport Exerc Med* 2018;4:e000424.
310 Fairburn CG, Beglin SJ. Assessment of eating disorders: Interview or self-report questionnaire? *Int J Eat Disord* 1994;16:363–70.
311 Stellingwerff T, Mountjoy M, McCluskey WT, *et al*. Review of the scientific rationale, development and validation of the International Olympic Committee Relative Energy Deficiency in Sport Clinical Assessment Tool: V.2 (IOC REDs CAT2)-by a subgroup of the IOC consensus on REDs. *Br J Sports Med* 2023;57:1109–18.
312 Sundgot-Borgen C, Sundgot-Borgen J, Sølvberg N, *et al*. Factors predicting disordered eating and the prevalence of eating disorders in adolescent elite athletes, trained athletes and a reference group: a prospective controlled two-step study. *Br J Sports Med* 2025;59:1043–54.
313 Hambleton A, Pepin G, Le A, *et al*. Psychiatric and medical comorbidities of eating disorders: Findings from a rapid review of the literature. *J Eat Disord* 2022;10:132.
314 Flatt RE, Miller AJ, Foreman T, *et al*. Considerations for athletes with eating disorders based on levels of care. *Curr Sports Med Rep* 2022;21:92–9.
315 Torstveit MK, Ackerman KE, Constantini N, *et al*. Primary, secondary and tertiary prevention of Relative Energy Deficiency in Sport (REDs): A narrative review by a subgroup of the IOC consensus on REDs. *Br J Sports Med* 2023;57:1119–28.
316 Arcelus J, Mitchell AJ, Wales J, *et al*. Mortality rates in patients with anorexia nervosa and other eating disorders. A meta-analysis of 36 studies. *Arch Gen Psychiatry* 2011;68:724–31.

Br J Sports Med: first published as 10.1136/bjsports-2025-111514 on 11 August 2026. Downloaded from http://bjsm.bmj.com/ on August 12, 2026 at Universitetsbiblioteket i Tromsø . Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies.

317 Ibáñez-Caparrós A, Sánchez I, Granero R, *et al*. Athletes with eating disorders: Analysis of their clinical characteristics, psychopathology and response to treatment. *Nutrients* 2023;15:3003.
318 Costandache GI, Munteanu O, Salaru A, *et al*. An overview of the treatment of eating disorders in adults and adolescents: Pharmacology and psychotherapy. *Postep Psychiatr Neurol* 2023;32:40–8.
319 Meeusen R, Piacentini MF, Van Den Eynde S, *et al*. Exercise performance is not influenced by a 5-HT reuptake inhibitor. *Int J Sports Med* 2001;22:329–36.
320 Schulz JM, White J, Edwards C, *et al*. Responding to Relative Energy Deficiency in Sport (REDs): A multidisciplinary care pathway for safe return to sport. *Sports Psychiatry* 2025;4:95–107.
321 Martinsen M, Bahr R, Børresen R, *et al*. Preventing eating disorders among young elite athletes: A randomized controlled trial. *Med Sci Sports Exerc* 2014;46:435–47.
322 Sundgot-Borgen C, Wisting L, Sundgot-Borgen J, *et al*. The "Young Athlete Body Project": A pilot study evaluating the acceptability of and results from an eating disorder prevention program for adolescent athletes. *Int J Eat Disord* 2024;57:568–80.
323 Martinsen M, Sherman RT, Thompson RA, *et al*. Coaches' knowledge and management of eating disorders: A randomized controlled trial. *Med Sci Sports Exerc* 2015;47:1070–8.
324 Vettoretti M. Official website of the International Federation of Sport Climbing. 2024. Available: https://www.ifsc-climbing.org/news/new-competition-policy-places-ifsc-at-forefront-of-fight-against-reds [Accessed 2 Nov 2024].
325 Virmavirta M, Kivekäs J. Is it still important to be light in ski jumping? *Sports Biomech* 2021;20:407–18.
326 Thomas R, Sanders S, Doust J, *et al*. Prevalence of attention-deficit/hyperactivity disorder: A systematic review and meta-analysis. *Pediatrics* 2015;135:e994–1001.
327 Putukian M, Kreher JB, Coppel DB, *et al*. Attention deficit hyperactivity disorder and the athlete: An American Medical Society for Sports Medicine position statement. *Clin J Sport Med* 2011;21:392–401.
328 Subcommittee on Attention-Deficit/Hyperactivity Disorder, Steering Committee on Quality Improvement and Management. ADHD: Clinical practice guideline for the diagnosis, evaluation, and treatment of attention-deficit/hyperactivity disorder in children and adolescents. *Pediatrics* 2011;128:1007–22.
329 Poysophon P, Rao AL. Neurocognitive deficits associated with ADHD in athletes: A systematic review. *Sports Health* 2018;10:317–26.
330 Parr JW. Attention-deficit hyperactivity disorder and the athlete: New advances and understanding. *Clin Sports Med* 2011;30:591–610.
331 Martin T. Public report of Major League Baseball's joint drug prevention and treatment program. 2020. Available: https://img.mlbstatic.com/mlb-images/image/upload/mlb/ylj7vghznkikcpolyiu3.pdf [Accessed 29 Aug 2025].
332 Martin T. Public report of Major League Baseball's joint drug prevention and treatment program. 2021. Available: https://img.mlbstatic.com/mlb-images/image/upload/mlb/o0zfqmwykwhyhbyfjhzn.pdf [Accessed 29 Aug 2025].
333 Martin T. Public report of Major League Baseball's joint drug prevention and treatment program. 2023. Available: https://img.mlbstatic.com/mlb-images/image/upload/mlb/o0zfqmwykwhyhbyfjhzn.pdf [Accessed 29 Aug 2025].
334 Martin T. Public report of Major League Baseball's joint drug prevention and treatment program. 2024. Available: https://www.mlb.com/press-release/release-independent-program-administrator-issues-annual-report-11-29-24 [Accessed 29 Aug 2025].
335 Kazda L, Bell K, Thomas R, *et al*. Overdiagnosis of attention-deficit/hyperactivity disorder in children and adolescents: A systematic scoping review. *JAMA Netw Open* 2021;4:e215335.
336 Han DH, McDuff D, Thompson D, *et al*. Attention-deficit/hyperactivity disorder in elite athletes: A narrative review. *Br J Sports Med* 2019;53:741–5.
337 Smith M. Hyperactive around the world? The history of ADHD in global perspective. *Soc Hist Med* 2017;30:767–87.
338 White RD, Harris GD, Gibson ME. Attention deficit hyperactivity disorder and athletes. *Sports Health* 2014;6:149–56.
339 Regnart J, Truter I, Meyer A. Critical exploration of co-occurring attention-deficit/hyperactivity disorder, mood disorder and substance use disorder. *Expert Rev Pharmacoecon Outcomes Res* 2017;17:275–82.
340 Cloninger CR. A systematic method for clinical description and classification of personality variants. A proposal. *Arch Gen Psychiatry* 1987;44:573–88.
341 Katzman MA, Bilkey TS, Chokka PR, *et al*. Adult ADHD and comorbid disorders: Clinical implications of a dimensional approach. *BMC Psychiatry* 2017;17:302.
342 Zhang L, Zhu N, Sjölander A, *et al*. ADHD drug treatment and risk of suicidal behaviours, substance misuse, accidental injuries, transport accidents, and criminality: Emulation of target trials. *BMJ* 2025;390:e083658.
343 Conant-Norville DO, Tofler IR. Attention deficit/hyperactivity disorder and psychopharmacologic treatments in the athlete. *Clin Sports Med* 2005;24:viii:829–43:.
344 Stewman CG, Liebman C, Fink L, *et al*. Attention deficit hyperactivity disorder: Unique considerations in athletes. *Sports Health: A Multidisciplinary Approach* 2018;10:40–6.
345 Hamilton RM, Rosenthal E, Hulpke-Wette M, *et al*. Cardiovascular considerations of attention deficit hyperactivity disorder medications: A report of the European Network on Hyperactivity Disorders work group, European Attention Deficit Hyperactivity Disorder Guidelines Group on attention deficit hyperactivity disorder drug safety meeting. *Cardiol Young* 2012;22:63–70.
346 Procyshyn RM, Bezchlibnyk-Butler KZ, Jeffries JJ. Drugs for ADHD. In: *Clinical Handbook of Psychotropic Drugs*. 23rd edn. Boston, MA: Hogrefe Publishing, 2019.
347 Moesch K, Kenttä G, Kleinert J, *et al*. FEPSAC position statement: Mental health disorders in elite athletes and models of service provision. *Psychol Sport Exerc* 2018;38:61–71.
348 Åkesdotter C, Kenttä G, Eloranta S, *et al*. The prevalence of mental health problems in elite athletes. *J Sci Med Sport* 2020;23:329–35.
349 Currie A, Gorczynski P, Rice SM, *et al*. Bipolar and psychotic disorders in elite athletes: A narrative review. *Br J Sports Med* 2019;53:746–53.
350 Bahrke MS, Yesalis CE III, Wright JE. Psychological and behavioural effects of endogenous testosterone and anabolic-androgenic steroids. An update. *Sports Med* 1996;22:367–90.
351 Daley MM, Reardon CL. Bipolar disorder and athletes: A narrative review. *Curr Sports Med Rep* 2021;20:638–44.
352 Piacentino D, Kotzalidis G, Casale A, *et al*. Anabolic-androgenic steroid use and psychopathology in athletes. A systematic review. *CN* 2015;13:101–21.
353 Trenton AJ, Currier GW. Behavioural manifestations of anabolic steroid use. *CNS Drugs* 2005;19:571–95.
354 Baron DA, Reardon CL, Baron SH. Doping in Sport. Oxford: John Wiley & Sons, 2013:21–32.
355 Bhardwaj A, Yadav J, Arya S, *et al*. Mephentermine misuse: An impending crisis among sportspersons. *J Psychoactive Drugs* 2022;54:196–8.
356 Kumari S, Chaudhary V, Makota VF, *et al*. Mephentermine use and adverse effects among athletes: A systematic review. *J Subst Use* 2024;29:1167–72.
357 Patel V, Kumar A, *et al*. Intravenous non-medical mephentermine use among young men: A consecutive case series from a tertiary addiction treatment facility in India. *Asian J Psychiatr* 2025;108:104525.
358 Markser V, Currie A, McAllister-WIlliams RH. Mood disorders. In: Currie A, Owen B, eds. *Sports Psychiatry*. Oxford: Oxford University Press, 2016: 31–51.
359 Judd LL, Akiskal HS, Schettler PJ, *et al*. A prospective investigation of the natural history of the long-term weekly symptomatic status of bipolar II disorder. *Arch Gen Psychiatry* 2003;60:261–9.
360 Judd LL, Akiskal HS, Schettler PJ, *et al*. The long-term natural history of the weekly symptomatic status of bipolar I disorder. *Arch Gen Psychiatry* 2002;59:530–7.
361 Morriss R. Bipolar disorder: treatment, psychological, social and physical health approaches. In: Kingdon D, Rowlands P, Stein G, eds. *Seminars in General Adult Psychiatry*. Cambridge University Press, 2024: 183–97.
362 Inder ML, Crowe MT, Porter R. Effect of transmeridian travel and jetlag on mood disorders: Evidence and implications. *Aust N Z J Psychiatry* 2016;50:220–7.
363 Goodwin GM, Haddad PM, Ferrier IN, *et al*. Evidence-based guidelines for treating bipolar disorder: Revised third edition recommendations from the British Association for Psychopharmacology. *J Psychopharmacol* 2016;30:495–553.
364 Hirschfeld RMA, Bowden CL, Gitlin MJ, *et al*. Practice guideline for the treatment of patients with bipolar disorder (revision). *FOC* 2003;1:64–110.
365 National Institute for Health and Care Excellence (NICE). Clinical guideline CG185: Bipolar disorder: Assessment and management. 2025. Available: https://www.nice.org.uk/guidance/cg185
366 Reardon CL. The sports psychiatrist and psychiatric medication. *Int Rev Psychiatry* 2016;28:606–13.
367 Johnston A, McAllister-Williams RH. Psychotropic drug prescribing. In: Currie A, Owen B, eds. *Sports psychiatry*. Oxford: Oxford University Press, 2016: 133–43.
368 Frank E, Soreca I, Swartz HA, *et al*. The role of interpersonal and social rhythm therapy in improving occupational functioning in patients with bipolar I disorder. *AJP* 2008;165:1559–65.
369 Barnes TR, Drake R, Paton C, *et al*. Evidence-based guidelines for the pharmacological treatment of schizophrenia: Updated recommendations from the British Association for Psychopharmacology. *J Psychopharmacol* 2020;34:3–78.
370 Lehman AF, Lieberman JA, Dixon LB. *Practice Guideline for the Treatment of Patients with Schizophrenia*. 2nd edn.2004. Available: https://psychiatryonline.org/pb/assets/raw/sitewide/practice_guidelines/guidelines/bipolar-1410197656063.pdf
371 National Institute for Health and Care Excellence (NICE). Guideline CG178: Psychosis and schizophrenia in adults: Prevention and management. 2019. Available: https://www.nice.org.uk/Guidance/CG178
372 Bredin SSD, Kaufman KL, Chow MI, *et al*. Effects of aerobic, resistance, and combined exercise training on psychiatric symptom severity and related health measures in adults living with schizophrenia: A systematic review and meta-analysis. *Front Cardiovasc Med* 2021;8:753117.
373 Bueno-Antequera J, Munguía-Izquierdo D. Exercise and schizophrenia. *Adv Exp Med Biol* 2020;1228:317–32.
374 Correll CU, Solmi M, Croatto G, *et al*. Mortality in people with schizophrenia: A systematic review and meta-analysis of relative risk and aggravating or attenuating factors. *World Psychiatry* 2022;21:248–71.
375 Dauwan M, Begemann MJH, Heringa SM, *et al*. Exercise improves clinical symptoms, quality of life, global functioning, and depression in schizophrenia: A systematic review and meta-analysis. *Schizophr Bull* 2016;42:588–99.

Br J Sports Med: first published as 10.1136/bjsports-2025-111514 on 11 August 2026. Downloaded from http://bjsm.bmj.com/ on August 12, 2026 at Universitetsbiblioteket i Tromsø . Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies.

376 Martinez-Calderon J, Villar-Alises O, García-Muñoz C, *et al*. Exercise or mind-body exercises for psychiatry symptoms and quality of life in schizophrenia spectrum and other psychotic disorders: An overview of systematic reviews with meta-analysis. *Disabil Rehabil* 2024;46:6034–50.
377 Firth J, Cotter J, Elliott R, *et al*. A systematic review and meta-analysis of exercise interventions in schizophrenia patients. *Psychol Med* 2015;45:1343–61.
378 Gallardo-Gómez D, Noetel M, Álvarez-Barbosa F, *et al*. Exercise to treat psychopathology and other clinical outcomes in schizophrenia: A systematic review and meta-analysis. *Eur Psychiatry* 2023;66:e40.
379 McCartan CJ, Yap J, Best P, *et al*. Factors that influence participation in physical activity for people with bipolar disorder: A synthesis of qualitative evidence. *Cochrane Database Syst Rev* 2024;6:CD013557.
380 Currie A, Malik R. Exercise participation and mental helath. In: Currie A, Owen B, eds. *Sport Psychiatry*. Oxford: Oxford University Press, 2016: 107–16.
381 Owen B. The athletic personality and personality disorders. In: CurrieA, OwenB, eds. *Sports Psychiatry*. Oxford, U.K.: Oxford University Press, 2016: 97–106.
382 Chang CJ, Putukian M, Aerni G, *et al*. Mental health issues and psychological factors in athletes: Detection, management, effect on performance, and prevention: American Medical Society for Sports Medicine position statement. *Clin J Sport Med* 2020;30:e61–87.
383 Widiger TA, Boyd SE. Personality disorders assessment instruments. In: Butcher JN, ed. *Oxford Handbook of Personality Disorders*. Oxford, UK, 2012: 336–63.
384 Glick ID, Stillman MA, Reardon CL, *et al*. Managing psychiatric issues in elite athletes. *J Clin Psychiatry* 2012;73:640–4.
385 Casale S, Banchi V. Narcissism and problematic social media use: A systematic literature review. *Addict Behav Rep* 2020;11:100252.
386 McCain JL, Campbell WK. Narcissism and social media use: A meta-analytic review. *Psychol Pop Media Cult* 2018;7:308–27.
387 Ceto E, Yigitoglu PH, Yavuz HU. Relationship between anabolic–androgenic steroid use, aggression, and narcissism in male bodybuilders. *Medicina (B Aires)* 2025;61:241.
388 Narcissism and empathy in steroid users. *AJP* 1995;152:1672–4.
389 Andersen MB, Denson EL, Brewer BW, *et al*. Disorders of personality and mood in athletes: Recognition and referral. *J Appl Sport Psychol* 1994;6:168–84.
390 Nicholls AR, Madigan DJ, Backhouse SH, *et al*. Personality traits and performance enhancing drugs: The dark triad and doping attitudes among competitive athletes. *Pers Individ Dif* 2017;112:113–6.
391 Nicholls AR, Madigan DJ, Duncan L, *et al*. Cheater, cheater, pumpkin eater: The dark triad, attitudes towards doping, and cheating behaviour among athletes. *Eur J Sport Sci* 2020;20:1124–30.
392 Plante TG, Booth J. Personality correlates of athletic injuries among elite collegiate baseball players: The role of narcissism, anger and locus of control. *J Human Movement Studies* 1997;32:47–59.
393 Osborn ZH, Blanton PD, Schwebel DC. Personality and injury risk among professional hockey players. *J Inj Violence Res* 2009;1:15–9.
394 Hendawy HMF, Awad EAA. Personality and personality disorders in athletes. In: Baron DA, Reardon CL, Baron SH, eds. *Clinical Sports Psychiatry: An International Perspective*. New York: Wiley, 2013: 53–64.
395 Galli N, A Petrie T, Greenleaf C, *et al*. Personality and psychological correlates of eating disorder symptoms among male collegiate athletes. *Eat Behav* 2014;15:615–8.
396 Shanmugam V, Davies B. Clinical perfectionism and eating psychopathology in athletes: The role of gender. *Pers Individ Dif* 2015;74:99–105.
397 Leggat FJ, Smith MJ, Figgins SG. Talented but disruptive: An exploration of problematic players in sports teams. *J Appl Sport Psychol* 2020;32:357–76.
398 Boulter MW, Roberts R, Hardy J. Upsetting the apple cart: Within-team profiles of intragroup conflict and their associations with narcissism. *Psychol Sport Exerc* 2022;63:102291.
399 Matosic D, Ntoumanis N, Boardley ID, *et al*. Narcissism, beliefs about controlling interpersonal style, and moral disengagement in sport coaches. *Int J Sport Exerc Psychol* 2020;18:592–606.
400 Chen Z, Wang D, Wang K, *et al*. Effects of coaching style on prosocial and antisocial behavior among Chinese athletes. *Soc Behav Pers* 2016;44:1889–900.
401 Biber DD, Czech DR. Comparing anxiety, aggression, and hostility between NCAA Division I athletes and non-athletes. *Applied Research in Coaching & Athletics Annual* 2021;36:1–22.
402 Leichsenring F, Heim N, Leweke F, *et al*. Borderline personality disorder: A review. *JAMA* 2023;329:670–9.
403 Ferraro JP. *Dialectical Behavior Therapy Application Efficacy in the World of an Athlete: Addressing Special Considerations*. Eastern Kentucky University, 2023
404 Marchant D, Gibbs P. Ethical considerations in treating borderline personality in sport: A case example. *Sport Psychol* 2004;18:317–23.
405 McLellan AT. Substance misuse and substance use disorders: Why do they matter in healthcare? *Trans Am Clin Climatol Assoc* 2017;128:112–30.
406 Exner J, Bitar R, Berg X, *et al*. Use of psychotropic substances among elite athletes - A narrative review. *Swiss Med Wkly* 2021;151.
407 National Collegiate Athletic Association. Research on student-athlete substance use, 2024. Available: https://ncaaorg.s3.amazonaws.com/research/wellness/Jan2024RES_HW-SubstanceUseRelease.pdf
408 Moore M, Abbe A. The National Association of Intercollegiate Athletics substance use and abuse survey. *J Issues Intercoll Athl* 2021;14:95–114.
409 Docter S, Khan M, Gohal C, *et al*. Cannabis use and sport: A systematic review. *Sports Health* 2020;12:189–99.
410 Kennedy M. Cannabis, cannabidiol and tetrahydrocannabinol in sport: An overview. *Intern Med J* 2022;52:1471–7.
411 Benoy R, Ramirez C, Hitchcock M, *et al*. Cannabis use in adolescent and young adult athletes: A clinical review. *Sports Health* 2024;16:279–84.
412 Charest J, Grandner MA, Athey AB, *et al*. Substance use among collegiate athletes versus non-athletes. *Athl Train Sports Health Care* 2021;13:e443–52.
413 Zeiger JS, Silvers WS, Fleegler EM, *et al*. Cannabis use in active athletes: Behaviors related to subjective effects. *PLoS One* 2019;14:e0218998.
414 Parisi CE, Bugbee BA, Vincent KB, *et al*. Risks associated with alcohol and marijuana use among college student athletes: The case for involving athletic personnel in prevention and intervention. *J Issues Intercoll Athl* 2019;12:343–64.
415 Mündel T. Nicotine: Sporting friend or foe? A review of athlete use, performance consequences and other considerations. *Sports Med* 2017;47:2497–506.
416 Ekhtiari S, Yusuf I, AlMakadma Y, *et al*. Opioid use in athletes: A systematic review. *Sports Health* 2020;12:534–9.
417 Ford JA, Pomykacz C, Veliz P, *et al*. Sports involvement, injury history, and non-medical use of prescription opioids among college students: An analysis with a national sample. *Am J Addict* 2018;27:15–22.
418 McDuff D, Stull T, Castaldelli-Maia JM, *et al*. Recreational and ergogenic substance use and substance use disorders in elite athletes: A narrative review. *Br J Sports Med* 2019;53:754–60.
419 McDuff DR, Garvin M, Chang J, *et al*. Substance misuse in elite athletes: Early detection, brief intervention and referral to treatment. *Clin Sports Med* 2024;43:127–44.
420 Stull T, Morse E, McDuff DR. Substance use and its impact on athlete health and performance. *Psychiatr Clin North Am* 2021;44:405–17.
421 Read D, Carter S, Hopley P, *et al*. Snus use in football: The threat of a new addiction? *Biol Sport* 2024;41:201–5.
422 Mountjoy M, Adriaens K, Junge A, *et al*. Association of self-reported health problems and interpersonal violence in sport: A cross-sectional study in world-level performing athletes. *Br J Sports Med* 2024;58:1360–8.
423 Dorfling T, Fulcher ML. High prevalence of harmful drinking habits and gambling among professional rugby players: Mental health symptoms and lifestyle risks among New Zealand Super Rugby players—A cross-sectional survey. *BMJ Open Sport Exerc Med* 2024;10:e002002.
424 Read D, Cope E, Taylor L. Snus Use in English Professional Football: Report Prepared for the English Professional Footballers' Association. Loughborough University, UK, 2024.
425 Hartmann T, Kiselev N, Claussen M, *et al*. Mental health and substance use among elite para-athletes: A cross-sectional survey. *CISS* 2024;9:016.
426 Olive LS, Rice SM, Gao C, *et al*. Risk and protective factors for mental ill-health in elite Para- and non-Para athletes. *Front Psychol* 2022;13:939087.
427 Handelsman DJ, *et al*. Performance Enhancing Hormone Doping in Sport. EndoText, 2020.
428 Coquet R, Roussel P, Ohl F. Understanding the paths to appearance- and performance-enhancing drug use in bodybuilding. *Front Psychol* 2018;9:1431.
429 Wenbo Z, Yan Z. The uses of anabolic androgenic steroids among athletes; its positive and negative aspects: A literature review. *J Multidiscip Healthc* 2023;16:4293–305.
430 Bhasin S, Hatfield DL, Hoffman JR, *et al*. Anabolic-androgenic steroid use in sports, health, and society. *Med Sci Sports Exerc* 2021;53:1778–94.
431 Mingxing L, Yanfei Y. Adverse effects of anabolic androgenic steroid abuse in athletes and physically active individuals: A systematic review and meta-analysis. *Substance Use & Misuse* 2025;60:873–87.
432 WADA. Anti-doping testing figures executive summary, 2023. Available: https://www.wada-ama.org/sites/default/files/2025-06/2023_anti-doping_testing_figures_executive_summary.pdf
433 WADA. Anti-doping testing figures executive summary, 2022. Available: https://www.wada-ama.org/sites/default/files/2024-04/2022_anti-doping_testing_figures_en.pdf
434 Davoren AK, Rulison K, Milroy J, *et al*. Doping prevalence among U.S. elite athletes subject to drug testing under the World Anti-Doping Code. *Sports Med - Open* 2024;10:57.
435 Robach P, Trebes G, Buisson C, *et al*. Prevalence of drug use in ultraendurance athletes. *Med Sci Sports Exerc* 2024;56:828–38.
436 Lipkens NR, Qeadan F. Athlete wellness: Examining the risk of alcohol use disorder in collegiate athletes and non-athletes. *J Sci Med Sport* 2025;28:447–53.
437 Bilgoe SC, Moen MH, Raedts E, *et al*. Mental health symptoms among Dutch elite athletes and their coaches: A cross-sectional study. *BMJ Open Sport Exerc Med* 2024;10:e001821.

Br J Sports Med: first published as 10.1136/bjsports-2025-111514 on 11 August 2026. Downloaded from http://bjsm.bmj.com/ on August 12, 2026 at Universitetsbiblioteket i Tromsoe . Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies.

438 Olive LS, Rice S, Butterworth M, *et al*. Do rates of mental health symptoms in currently competing elite athletes in Paralympic sports differ from non-Para-athletes? *Sports Med - Open* 2021;7:62.
439 Gorelick DA. Cannabis-related disorders and toxic effects. *N Engl J Med* 2023;389:2267–75.
440 McDuff DR DR, Garvin M M, Thompson DL DL. Substance use and substance use disorders. In: Reardon CL, ed. *Mental Health Care for Elite Athletes*. Champaign, IL, 2022: 131–43.
441 Kozhuharov VR, Ivanov K, Ivanova S. Dietary supplements as source of unintentional doping. *Biomed Res Int* 2022;2022:8387271.
442 Walters P, Hillier B, Passetti F, *et al*. Diagnosis and management of substance use disorders in athletes. *Advances in Psychiatry and Behavioral Health* 2021;1:135–43.
443 Ghergel A-M, Woodhouse D, Ramchandani G. A systematic review of gambling amongst elite athletes with a focus on females. *JGI* 2026.
444 Håkansson A, Durand-Bush N, Kenttä G. Problem gambling and problem gaming in elite athletes: A literature review. *Int J Ment Health Addiction* 2023;21:1837–53.
445 Nelson SE, Slabczynski JM, Lee TG, *et al*. All in: A scoping review of the association between gambling and athletic participation. *Int Gambl Stud* 2024;24:185–209.
446 Paskus TS, Derevensky J. Trends in NCAA student-athlete gambling and sports betting behaviors, 2025. Available: https://ncaaorg.s3.amazonaws.com/research/wagering/2025RES_WageringReport.pdf
447 Winters KC, Derevensky JL. A review of sports wagering: Prevalence, characteristics of sports bettors, and association with problem gambling. *JGI* 2020;43:102–27.
448 Calado F, Griffiths MD. Problem gambling worldwide: An update and systematic review of empirical research (2000-2015). *J Behav Addict* 2016;5:592–613.
449 Vinberg M, Durbeej N, Rosendahl I. Young male athletes at risk for problem gambling: Findings from a cross-sectional study in Sweden. *Scand J Psychol* 2023;64:766–75.
450 González-Moret R, Almodóvar-Fernández I, Gimeno M, *et al*. Gambling and gaming: A comparative study of professional footballers versus the general population. *Sports (Basel)* 2025;13:34.
451 Valenciano-Mendoza E, Mora-Maltas B, Mestre-Bach G, *et al*. Clinical correlates of sports betting: A systematic review. *J Gambl Stud* 2023;39:579–624.
452 Richard J, Paskus TS, Derevensky JL. Trends in gambling behavior among college student-athletes: A comparison of 2004, 2008, 2012 and 2016 NCAA survey data. *JGI* 2019;41:73–100.
453 Poucher ZA, Tamminen KA, Kerr G, *et al*. A commentary on mental health research in elite sport. *J Appl Sport Psychol* 2021;33:60–82.
454 Derevensky JL. *Teen gambling*. 2012. Available: https://www.nomos-elibrary.de/index.php?doi=10.5771/9781442202283
455 World Health Organization. ICD-11: International statistical classification of diseases and related health problems, 2021. Available: https://www.who.int/standards/classifications/classification-of-diseases
456 Wardle H, Degenhardt L, Ceschia A, *et al*. The Lancet Public Health Commission on gambling. *Lancet Public Health* 2021;6:e2–3.
457 National Collegiate Athlete Association. Online abuse in NCAA championships: NCAA pilot study 2023-2024, 2024. Available: https://ncaaorg.s3.amazonaws.com/ncaa/wagering/NCAAThreatMatrixPilotStudyPublicReport.pdf
458 Di Lodovico L, Poulnais S, Gorwood P. Which sports are more at risk of physical exercise addiction: A systematic review. *Addict Behav* 2019;93:257–62.
459 Szabo A PhD, De La Vega R, Ruiz-Barquín R, *et al*. Exercise addiction in Spanish athletes: Investigation of the roles of gender, social context and level of involvement. *J Behav Addict* 2013;2:249–52.
460 Mayolas-Pi C, Sitko S, Pano-Rodriguez A, *et al*. Exercise addiction and psychosocial health risks among adolescent athletes: Focus on sport type and performance level. *J Behav Addict* 2025;14:1095–106.
461 Lichtenstein MB, Melin AK, Szabo A, *et al*. The prevalence of exercise addiction symptoms in a sample of national level elite athletes. *Front Sports Act Living* 2021;3:635418.
462 Colledge F, Cody R, Buchner UG, *et al*. Excessive exercise—A meta-review. *Front Psychiatry* 2020;11:521572.
463 Egorov AY, Szabo A. The exercise paradox: An interactional model for a clearer conceptualization of exercise addiction. *J Behav Addict* 2013;2:199–208.
464 de Coverley Veale DM. Exercise dependence. *Br J Addict* 1987;82:735–40.
465 Çetin E, Bulğay C, Demir GT, *et al*. The examination of the relationship between exercise addiction and performance enhancement in elite athletes. *Int J Ment Health Addiction* 2021;19:1019–30.
466 Lichtenstein MB, Nielsen RO, Gudex C, *et al*. Exercise addiction is associated with emotional distress in injured and non-injured regular exercisers. *Addict Behav Rep* 2018;8:33–9.
467 Torstveit MK, Fahrenholtz IL, Lichtenstein MB, *et al*. Exercise dependence, eating disorder symptoms and biomarkers of Relative Energy Deficiency in Sports (RED-S) among male endurance athletes. *BMJ Open Sport Exerc Med* 2019;5:e000439.
468 Håkansson A, Kenttä G, Åkesdotter C. Problem gambling and gaming in elite athletes. *Addict Behav Rep* 2018;8:79–84.
469 Mestre-Bach G, Granero R, Mora-Maltas B, *et al*. Sports-betting-related gambling disorder: Clinical features and correlates of cognitive behavioral therapy outcomes. *Addict Behav* 2022;133:107371.
470 Hodgins DC, Stea JN, Grant JE. Gambling disorders. *Lancet* 2011;378:1874–84.
471 Adams J. Understanding exercise dependence. *J Contemp Psychother* 2009;39:231–40.
472 Richard J, Derevensky JL. Counseling Interventions for Youth Problem Gambling. Ontario: CDS Press, 2023.
473 Freimuth M, Moniz S, Kim SR. Clarifying exercise addiction: Differential diagnosis, co-occurring disorders, and phases of addiction. *IJERPH* 2011;8:4069–81.
474 Ginley MK, Zajac K, Rash CJ, *et al*. Pharmacological Treatment of Behavioral Addictions. Washington DC: American Psychology Association, 2019.
475 Zhao Y, Marchica L, Derevensky JL, *et al*. The scope, focus and types of gambling policies among Canadian colleges and universities. *Canadian Psychology / Psychologie Canadienne* 2017;58:187–93.
476 Cassidy JD, Carroll L, Peloso P, *et al*. Incidence, risk factors and prevention of mild traumatic brain injury: Results of the WHO collaborating centre task force on mild traumatic brain injury. *J Rehabil Med* 2004;36:28–60.
477 Gardner AJ, Quarrie KL, Iverson GL. The epidemiology of sport-related concussion: What the rehabilitation clinician needs to know. *J Orthopaed Sports Physical Ther* 2019;49:768–78.
478 Chandran A, Boltz AJ, Morris SN, *et al*. Epidemiology of concussions in National Collegiate Athletic Association (NCAA) sports: 2014/15-2018/19. *Am J Sports Med* 2022;50:526–36.
479 Patricios JS, Schneider KJ, Dvorak J, *et al*. Consensus statement on concussion in sport. *The 6th International Conference on Concussion in Sport-Amsterdam* 2022;57:695–711.
480 Echemendia RJ, Brett BL, Broglio S. Sport Concussion Assessment Tool 6 (SCAT6). *Br J Sports Med* 2023;57:622–31.
481 Henry LC, Elbin RJ, Collins MW, *et al*. Examining recovery trajectories after sport-related concussion with a multimodal clinical assessment approach. *Neurosurgery* 2016;78:232–41.
482 McAllister TW, Broglio SP, Katz BP, *et al*. Characteristics and outcomes of athletes with slow recovery from sports-related concussion. *Neurology* 2023;100:e1510–9.
483 Harmon KG, Clugston JR, Dec K, *et al*. American Medical Society for Sports Medicine position statement on concussion in sport. *Br J Sports Med* 2019;53:213–25.
484 Rice SM, Parker AG, Rosenbaum S, *et al*. Sport-related concussion and mental health outcomes in elite athletes: A systematic review. *Sports Med* 2018;48:447–65.
485 Trinh LN, Brown SM, Mulcahey MK. The influence of psychological factors on the incidence and severity of sports-related concussions: A systematic review. *Am J Sports Med* 2020;48:1516–25.
486 Thomas GA, Bradson ML, Riegler KE, *et al*. Association between co-occurring anxiety and depressive symptoms at baseline and risk for sports-related concussion in collegiate athletes. *Orthop J Sports Med* 2024;12:23259671241255932.
487 Broglio SP, Lapointe A, O'Connor KL, *et al*. Head impact density: A model to explain the elusive concussion threshold. *J Neurotrauma* 2017;34:2675–83.
488 Iverson GL, Williams MW, Gardner AJ, *et al*. Systematic review of preinjury mental health problems as a vulnerability factor for worse outcome after sport-related concussion. *Orthop J Sports Med* 2020;8:2325967120950682.
489 Vargas G, Rabinowitz A, Meyer J, *et al*. Predictors and prevalence of postconcussion depression symptoms in collegiate athletes. *J Athl Train* 2015;50:250–5.
490 Kilic Ö, Hopley P, Kerkhoffs G, *et al*. Impact of concussion and severe musculoskeletal injuries on the onset of mental health symptoms in male professional rugby players: A 12-month study. *BMJ Open Sport Exerc Med* 2019;5:e000693.
491 Blaney NA, Trbovich A, Ernst N, *et al*. Influence of sleep dysfunction on concussion assessment outcomes among adolescent athletes after concussion and healthy controls. *Clin J Sport Med* 2021;31:481–7.
492 Brustman K, Eagle SR, Mucha A, *et al*. Association of sleep symptoms with mood and vestibular subtypes following sport-related concussion. *Appl Neuropsychol Child* 2022;11:235–9.
493 Thomas GA, Riegler KE, Bradson ML, *et al*. Depressive symptoms, sleep somplaints, and cognitive performance: Examining the mediating role of sleep in college athletes. *Arch Clin Neuropsychol* 2025;40:1348–56.
494 Sufrinko A, Pearce K, Elbin RJ, *et al*. The effect of preinjury sleep difficulties on neurocognitive impairment and symptoms after sport-related concussion. *Am J Sports Med* 2015;43:830–8.
495 Iverson GL, Terry DP. High school football and risk for depression and suicidality in adulthood: Findings from a national longitudinal study. *Front Neurol* 2021;12:812604.
496 Iverson GL, Merz ZC, Terry DP. Playing high school football is not associated with an increased risk for suicidality in early adulthood. *Clin J Sport Med* 2021;31:469–74.
497 Deshpande SK, Hasegawa RB, Weiss J, *et al*. The association between adolescent football participation and early adulthood depression. *PLoS One* 2020;15:e0229978.
498 Deshpande SK, Hasegawa RB, Rabinowitz AR, *et al*. Association of playing high school football with cognition and mental health later in life. *JAMA Neurol* 2017;74:909–18.
499 Bohr AD, Boardman JD, McQueen MB. Association of adolescent sport participation with cognition and depressive symptoms in early adulthood. *Orthop J Sports Med* 2019;7:2325967119868658.
500 Lehman EJ, Hein MJ, Gersic CM. Suicide mortality among retired National Football League players who played 5 or more seasons. *Am J Sports Med* 2016;44:2486–91.

Br J Sports Med: first published as 10.1136/bjsports-2025-111514 on 11 August 2026. Downloaded from http://bjsm.bmj.com/ on August 12, 2026 at Universitetsbiblioteket i Tromsø . Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies.

501 Russell ER, McCabe T, Mackay DF, *et al*. Mental health and suicide in former professional soccer players. *J Neurol Neurosurg Psychiatry* 2020;91:1256–60.
502 Taioli E. All causes of mortality in male professional soccer players. *Eur J Public Health* 2007;17:600–4.
503 Leddy JJ, Haider MN, Ellis MJ, *et al*. Early subthreshold aerobic exercise for sport-related concussion: A randomized clinical trial. *JAMA Pediatr* 2019;173:319–25.
504 Leddy JJ, Burma JS, Toomey CM, *et al*. Rest and exercise early after sport-related concussion: A systematic review and meta-analysis. *Br J Sports Med* 2023;57:762–70.
505 Leddy JJ, Master CL, Mannix R, *et al*. Early targeted heart rate aerobic exercise versus placebo stretching for sport-related concussion in adolescents: A randomised controlled trial. *Lancet Child Adolesc Health* 2021;5:S2352-4642(21)00267-4:792–9:.
506 Art K, Ridenour C, Durbin S, *et al*. The effectiveness of physical therapy interventions for athletes post-concussion: A systematic review. *Int J Sports Phys Ther* 2023;18:26–38.
507 Schneider KJ, Critchley ML, Anderson V, *et al*. Targeted interventions and their effect on recovery in children, adolescents and adults who have sustained a sport-related concussion: A systematic review. *Br J Sports Med* 2023;57:771–9.
508 Tracey AJ, Bateman AG, Baez SE, *et al*. Effectiveness of interventions for the improvement of mental health and well-being post-concussion: A systematic review. *Brain Inj* 2023;37:1135–58.
509 Weiler R, Blauwet C, Clarke D, *et al*. Concussion in para sport: The first position statement of the Concussion in Para Sport (CIPS) Group. *Br J Sports Med* 2021;55:1187–95.
510 Smetana RM, Kaplan DT, Magill RT, *et al*. Sport-related concussion in Para athletes: A scoping review of concussion incidence, assessment, and management. *Am J Phys Med Rehabil* 2024;103:858–65.
511 Purcell R, Pilkington V, Carberry S, *et al*. An evidence-informed framework to promote mental wellbeing in elite sport. *Front Psychol* 2022;13:780359.
512 Mathisen TF, Ackland T, Burke LM, *et al*. Best practice recommendations for body composition considerations in sport to reduce health and performance risks: A critical review, original survey and expert opinion by a subgroup of the IOC consensus on Relative Energy Deficiency in Sport (REDs). *Br J Sports Med* 2023;57:1148–60.
513 Lundqvist C, Camps J, Vertommen T, *et al*. Toxic leadership in high-performance sports and its consequences for mental health and performance: A scoping review. *Int Rev Sport Exerc Psychol* 2025;1–27.
514 Runquist EB III, Adenaiye OO, Sarzaeim M, *et al*. Associations of abusive supervision among collegiate athletes from equity-deserving groups. *Br J Sports Med* 2025;59:1124–34.
515 Brown DJ, Arnold R, Standage M, *et al*. The prediction of thriving in elite sport: A prospective examination of the role of psychological need satisfaction, challenge appraisal, and salivary biomarkers. *J Sci Med Sport* 2021;24:373–9.
516 Bean C, Forneris T. Examining the importance of intentionally structuring the youth sport context to facilitate positive youth development. *J Appl Sport Psychol* 2016;28:410–25.
517 Giles S. Evaluating the effectiveness of the implementation of elite athlete mental health support services in the United Kingdom. International Olympics Committee; 2025. Available: https://library.olympics.com/Default/doc/SYRACUSE/3788456/evaluating-the-effectiveness-of-the-implementation-of-elite-athlete-mental-health-support-services-i
518 Hapig M, Zurstiege G, Mayer J, *et al*. Exploring print media coverage of elite athletes' mental illness between 2010 and 2023 in Germany: A quantitative content analysis. *Front Sports Act Living* 2024;6:1446680.
519 Bissett JE, Kroshus E, Hebard S. Determining the role of sport coaches in promoting athlete mental health: A narrative review and Delphi approach. *BMJ Open Sport Exerc Med* 2020;6:e000676.
520 Sankey C, Wallace L, Caperchione CM. Understanding the role of coaches in supporting the mental health of elite athletes. *J Sci Med Sport* 2023;26:399–404.
521 Armstrong SN, Burcin MM, Bjerke W, *et al*. Depression in student athletes: A particularly at-risk group? A systematic review of the literature. *Athletic Insight* 2015;7:177–93. Available: https://digitalcommons.sacredheart.edu/cgi/viewcontent.cgi?article=1186&context=pthms_fac
522 Prinz B, Dvořák J, Junge A. Symptoms and risk factors of depression during and after the football career of elite female players. *BMJ Open Sport Exerc Med* 2016;2:e000124.
523 Lonsdale C, Hodge K, Rose E. Athlete burnout in elite sport: A self-determination perspective. *J Sports Sci* 2009;27:785–95.
524 Bergeron MF, Mountjoy M, Armstrong N, *et al*. International Olympic Committee consensus statement on youth athletic development. *Br J Sports Med* 2015;49:843–51.
525 Kramers S, Thrower SN, Steptoe K, *et al*. Parental strategies for supporting children's psychosocial development within and beyond elite sport. *J Appl Sport Psychol* 2023;35:498–520.
526 Bergeron MF, Côté J, Cumming SP, *et al*. IOC consensus statement on elite youth athletes competing at the Olympic Games: Essentials to a healthy, safe and sustainable paradigm. *Br J Sports Med* 2024;58:946–65.
527 Wendling E, Flaherty M, Sagas M, *et al*. Youth athletes' sustained involvement in elite sport: An exploratory examination of elements affecting their athletic participation. *Int J Sports Sci Coaching* 2018;13:658–73.
528 Anderson RJ, Pierce D. Assumptions associated with mental health literacy training – Insights from initiatives in rural Australia. *Advances in Mental Health* 2012;10:258–67.
529 Kitchener BA, Jorm AF. Mental health first aid training for the public: Evaluation of effects on knowledge, attitudes and helping behavior. *BMC Psychiatry* 2002;2:10.
530 Hadlaczky G, Hökby S, Mkrtchian A, *et al*. Mental Health First Aid is an effective public health intervention for improving knowledge, attitudes, and behaviour: A meta-analysis. *Int Rev Psychiatry* 2014;26:467–75.
531 Krug EG, Mercy JA, Dahlberg LL, *et al*. The world report on violence and health. *The Lancet* 2002;360:1083–8.
532 Mountjoy M, Brackenridge C, Arrington M, *et al*. International Olympic Committee consensus statement: Harassment and abuse (non-accidental violence) in sport. *Br J Sports Med* 2016;50:1019–29.
533 Vertommen T, Kampen J, Schipper-van Veldhoven N, *et al*. Profiling perpetrators of interpersonal violence against children in sport based on a victim survey. *Child Abuse Negl* 2017;63:172–82.
534 Timon CE, Dallam SJ, Hamilton MA, *et al*. Child sexual abuse of elite athletes: Prevalence, perceptions, and mental health. *J Child Sex Abus* 2022;31:672–91.
535 Kerr G, Willson E, Stirling A. "It was the worst time in my life": The effects of emotionally abusive coaching on female Canadian national team athletes. *Women in Sport and Physical Activity Journal* 2020;28:81–9.
536 Greblo JurakiĆ Z, LjubičIĆ V. Sexual harrassment of young athletes by their coaches and depression, anxiety, and stress in later life. *Društvena Istraživanja* 2022;31:135–54.
537 Parent S, Vaillancourt-Morel M-P, Gillard A. Interpersonal violence (IV) in sport and mental health outcomes in teenagers. *Journal of Sport and Social Issues* 2022;46:323–37.
538 Salim J, Winter S. "I still wake up with nightmares" … The long-term psychological impacts from gymnasts' maltreatment experiences. *Sport Exerc Perform Psychol* 2022;11:429–43.
539 Mountjoy M, Junge A, Magnusson C, *et al*. Beneath the surface: Mental health and harassment and abuse of athletes participating in the FINA (Aquatics) World Championships, 2019. *Clin J Sport Med* 2022;32:95–102.
540 McMahon J, McGannon KR. 'I hurt myself because it sometimes helps': Former athletes' embodied emotion responses to abuse using self-injury. *Sport Educ Soc* 2021;26:161–74.
541 Wathen CN, Schmitt B, MacGregor JCD. Measuring trauma- (and violence-) informed care: A scoping review. *Trauma Violence Abuse* 2023;24:261–77.
542 Gorczynski P, Currie A, Gibson K, *et al*. Developing mental health literacy and cultural competence in elite sport. *J Appl Sport Psychol* 2021;33:387–401.
543 Gorczynski P, Baron D. Designing and delivering mental health literacy strategies in elite sport. *World Psychiatry* 2024;23:169–71.
544 d'Andrade RG. *The Development of Cognitive Anthropology*. Cambridge University Press, 1995.
545 Office of the Surgeon General, Center for Mental Health Services. Culture counts: The influence of culture and society on mental health: Culture, race, and ethnicity: A supplement to mental health. A report of the Surgeon General. Substance Abuse and Mental Health Services Administration (US); 2001.
546 Coyle M, Gorczynski P, Gibson K. "You have to be mental to jump off a board any way": Elite divers' conceptualizations and perceptions of mental health. *Psychol Sport Exerc* 2017;29:10–8.
547 Moreland JJ, Coxe KA, Yang J. Collegiate athletes' mental health services utilization: A systematic review of conceptualizations, operationalizations, facilitators, and barriers. *J Sport Health Sci* 2018;7:58–69.
548 Ojio Y, Amemiya R, Oliffe JL, *et al*. Mental health help-seeking knowledge, attitudes and behaviour among male elite rugby players: The role of masculine health-related values. *BMJ Open Sport Exerc Med* 2025;11:e002275.
549 Lekas HM, Pahl K, Fuller Lewis C. Rethinking cultural competence: Shifting to cultural humility. *Health Serv Insights* 2020;13:1178632920970580.
550 Picha KJ, Welch Bacon CE, Normore C, *et al*. Social determinants of health: Considerations for athletic health care. *J Athl Train* 2022;57:521–31.
551 National Academies of Sciences, Engineering, Medicine, Health. *Communities in Action: Pathways to Health Equity*. Washington, DC: National Academies Press (US), 2017.
552 Kirkbride JB, Anglin DM, Colman I, *et al*. The social determinants of mental health and disorder: Evidence, prevention and recommendations. *World Psychiatry* 2024;23:58–90.
553 Kroshus E, Bell L, Gurganus-Wright K, *et al*. Structural and social determinants of mental health inequities among collegiate athletes during the COVID-19 pandemic. *Br J Sports Med* 2023;57:1435–41.
554 World Health Organization. Social determinants of health. 2026 Available: https://www.who.int/health-topics/social-determinants-of-health#tab=tab_1
555 Chan ASW, Choong A, Phang KC, *et al*. Societal discrimination and mental health among transgender athletes: A systematic review and meta-analysis. *BMC Psychol* 2024;12:24.

Br J Sports Med: first published as 10.1136/bjsports-2025-111514 on 11 August 2026. Downloaded from http://bjsm.bmj.com/ on August 12, 2026 at Universitetsbiblioteket i Tromsø . Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies.

556 Gorczynski P, Reardon CL, Aron CM. Lesbian, gay, bisexual, trans, and queer mental health in elite sport: A review. *Advances in Psychiatry and Behavioral Health* 2022;2:9–16.
557 Rovira-Font M, Vilanova-Soler A. LGTBIQA+, mental health and the sporting context: A systematic review. *Apunts Educación Física y Deportes* 2022;147:1–16.
558 Giffin CE, Schinke RJ, Larivière M, *et al*. Migration and meaning: An exploration of elite refugee athletes' transitions into the Canadian sports system. *Int J Sport Exerc Psychol* 2024;22:1460–79.
559 Michelini E. War, migration, resettlement and sport socialization of young athletes: The case of Syrian elite water polo. *Eur J Sport Society* 2018;15:5–21.
560 Ray V. A theory of racialized organizations. *Am Sociol Rev* 2019;84:26–53.
561 NCAA. Mental health practices: Understanding and supporting student-athlete mental health. An inter-association consensus document. NCAA Sport Science Institute; 2024. Available: https://ncaaorg.s3.amazonaws.com/ssi/mental/SSI_MentalHealthBestPractices.pdf
562 Jeste DV, Smith J, Lewis-Fernández R, *et al*. Addressing social determinants of health in individuals with mental disorders in clinical practice: Review and recommendations. *Transl Psychiatry* 2025;15:120.
563 Castaldelli-Maia JM, Gallinaro JG de ME, Falcão RS, *et al*. Mental health symptoms and disorders in elite athletes: A systematic review on cultural influencers and barriers to athletes seeking treatment. *Br J Sports Med* 2019;53:707–21.
564 Malchrowicz-Mośko E, Wieliński D, Adamczewska K. Perceived benefits for mental and physical health and barriers to horseback riding participation. The analysis among professional and amateur athletes. *Int J Environ Res Public Health* 2020;17:3736.
565 Breslin G, Shannon S, Haughey T, *et al*. A systematic review of interventions to increase awareness of mental health and well-being in athletes, coaches and officials. *Syst Rev* 2017;6:177.
566 Yoon JJ, Petrie T. Barriers to and facilitators of collegiate athletes seeking mental health services. *J Athl Train* 2023;58:715–21.
567 Roche E, Richardson N, Sweeney J, *et al*. Workplace interventions targeting mental health literacy, stigma, help-seeking, and help-offering in male-dominated industries: A systematic review. *Am J Mens Health* 2024;18:15579883241236223.
568 Kim SY, Choi C. Differences in stress, stress-coping behavior, and quality of life based on the performance of Korean ladies Professional Golf Association tour players. *IJERPH* 2022;19:6623.
569 Gorczynski P, Reardon CL, Aron CM. Mental health in golf. In: *The Psychology of Golf Performance Under Pressure*. Routledge, 2023: 198–209.
570 Rathod S, Pinninti N, Irfan M, *et al*. Mental health service provision in low- and middle-income countries. *Health Serv Insights* 2017;10:1178632917694350.
571 Pilkington V, Walton CC, Gwyther K, *et al*. Mental health service provision in elite sport: An evaluation of the Australian Institute of Sport Mental Health Referral Network. *J Appl Sport Psychol* 2025;37:397–421.
572 Andersson MJ, Abdul Rahim Y, Kenttä G, *et al*. Mental health challenges in elite sports, barriers to treatment, and quality of psychiatric care at an elite sports-centered mental health clinic: A mixed-methods study. *Psychol Sport Exerc* 2025;79:102859.
573 Chrétien A, D'arripe-Longueville F, Haffner K, *et al*. Using a socioecological approach to explore healthy lifestyle in elite sport: A qualitative study of French athletes', coaches', and managers' perspectives. *Int J Qual Stud Health Well-Being* 2023;18:2251235.
574 Cosh SM, McNeil DG, Jeffreys A, *et al*. Athlete mental health help-seeking: A systematic review and meta-analysis of rates, barriers and facilitators. *Psychol Sport Exerc* 2024;71:102586.
575 Oftadeh-Moghadam S, Weston N, Gorczynski P. Mental health literacy intervention to reduce stigma toward mental health symptoms and disorders in women rugby players: A feasibility study. *Sport Psychol* 2023;37:119–27.
576 Diamond S, Wallace L, English M, *et al*. The impact of mental health literacy initiatives on youth elite athletes: A systematic review. *Performance Enhancement & Health* 2022;10:100226.
577 Bu D, Chung P-K, Zhang C-Q, *et al*. Mental health literacy intervention on help-seeking in athletes: A systematic review. *Int J Environ Res Public Health* 2020;17:7263.
578 Chow GM, Bird MD, Gabana NT, *et al*. A program to reduce stigma toward mental illness and promote mental health literacy and help-seeking in National Collegiate Athletic Association Division I student-athletes. *J Clin Sport Psychol* 2021;15:185–205.
579 Gorczynski P, Aron CM, Moore M, *et al*. The epidemiology of mental health symptoms and disorders among elite athletes and the evolution of mental health literacy. *Clin Sports Med* 2024;43:1–11.
580 Moore MA, Gorczynski P, Miller Aron C. Mental health literacy in sport: The role of the social work profession. *Soc Work* 2022;67:298–300.
581 Social media and youth mental health: the US Surgeon General's Advisory. Washington DC: US Department of Health and Human Services; 2023. Available: https://hhs.gov/sites/default/files/sg-youth-mental-health-social-media-advisory.pdf [Accessed 20 Jun 2023].
582 Faustin M, Burton M, Callender S, *et al*. Effect of media on the mental health of elite athletes. *Br J Sports Med* 2022;56:123–4.
583 Putukian M, Blauwet C, Currie A, *et al*. Social media impact on athlete mental health: #RealityCheck. *Br J Sports Med* 2024;58:463–5.
584 Suchland H, Smith-Goodwin E, Dick E. Effects of TikTok on the mental health of men's and women's soccer teams at one institution. *JSMAHS* 2022;8:29–30.
585 Barry CT, Moran-Miller K, Levy HF, *et al*. Social media engagement, perceptions of social media costs and benefits, and well-being in college student-athletes. *Journal of American College Health* 2024;72:2938–47.
586 Zhang W, Jiang F, Zhu Y, *et al*. Risks of passive use of social network sites in youth athletes: A moderated mediation analysis. *Front Psychol* 2023;14:1219190.
587 Watkins RA, Sugimoto D, Hunt D, *et al*. Association of social media use on sleep quality and performance among collegiate athletes. *Clin J Sport Med* 2022;32:486–92.
588 Adjanin N, Beemer RC. Beyond the hashtags: The impacts of self-promotion and social media on sleep and mental health among student-athletes. *Journal of Intercollegiate Sport* 2024;17:1–25. Available: https://doi.org/10.17161/jis.v17i3.21474
589 Bjornsen CA, O'Connor J. Self-disclosure of body image, eating, and emotional concerns by female collegiate athletes: Associations with body image pressure, social media, and emotional distress. *Canadian Journal of Behavioural Science / Revue Canadienne Des Sciences Du Comportement* 2025;57:108–18.
590 Fiedler R, Heidari J, Birnkraut T, *et al*. Digital media and mental health in adolescent athletes. *Psychol Sport Exerc* 2023;67:102421.
591 Marco JH, Tormo-Irun MP, Galán-Escalante A, *et al*. Is cybervictimization associated with body dissatisfaction, depression, and eating disorder psychopathology. *Cyberpsychol Behav Soc Netw* 2018;21:611–7.
592 Wilson C, Sheridan L, Garratt-Reed D. What is cyberstalking? A review of measurements. *J Interpers Violence* 2022;37:NP9763–83.
593 Kavanagh E, Mountjoy M, Teunissen L, *et al*. "I'm outside, and I'm going to hurt you": The dark side of stalking in sport. *Br J Sports Med* 2025;59:1454–6.
594 World Athletics. World Athletics publishes online abuse study covering World Athletics Championships, Oregon. 2022. Available: https://worldathletics.org/news/press-releases/online-abuse-study-world-athletics-championships-oregon22 [Accessed 12 Nov 2023].
595 Phipps C. "You belong in the kitchen": Social media, virtual manhood acts, and women strength sport athletes' experiences of gender-based violence online. *Feminist Media Studies* 2023;23:4221–37.
596 Pocock M, Skey M. 'You feel a need to inspire and be active on these sites otherwise… people won't remember your name': Elite female athletes and the need to maintain 'appropriate distance' in navigating online gendered space. *New Media & Society* 2024;26:961–77.
597 Hayes M, Filo K, Riot C, *et al*. Athlete perceptions of social media benefits and challenges during major sport events. *Int J Sport Communication* 2019;12:449–81.
598 Vogels EA, Gelles-Watnick R, Massarat N. Teens, social media and technology 2022. Pew Research Center; 2022. Available: https://www.pewresearch.org/internet/2022/08/10/teens-social-media-and-technology-2022/
599 International Olympic Committee. Social media resources. 2023. Available: https://olympics.com/athlete365/courses/mindful-social-media/ [accessed Jul 2025]
600 Kavanagh EJ, Litchfield C, Osborne J. Social media and athlete welfare. *Int J Sport Communication* 2023;16:274–81.
601 Mitchell FR, van Wyk PM, Santarossa S. Curating a culture: The portrayal of disability stereotypes by Paralympians on instagram. *Int J Sport Communication* 2021;14:334–55.
602 American Psychological Association. Health advisory on social media use in adolescence. 2023. Available: https://www.apa.org/topics/social-media-internet/health-advisory-adolescent-social-media-use.pdf [Accessed 8 Aug 2025].
603 Parrott S, Billings AC, Hakim SD, *et al*. From #endthestigma to #realman: Stigma-challenging social media responses to NBA players' mental health disclosures. *Communication Reports* 2020;33:148–60.
604 Weber S, Olsen M, Martiny SE. Effects of instagram sports posts on the athletic motivation of female elite athletes: Do they inspire or backfire? *Psychol Sport Exerc* 2022;58:102080.
605 Brougham JK. The impact of social media on the mental health of student-athletes across NCAA divisions. *J Issues Intercoll Athl* 2021;713–39. Available: https://scholarcommons.sc.edu/jiia/vol14/iss1/33/
606 Encel K, Mesagno C, Brown H. Facebook use and its relationship with sport anxiety. *J Sports Sci* 2017;35:756–61.
607 International Olympic Committee. World Mental Health Day: How the IOC ran the largest online abuse prevention programme ever conducted in sport at Paris 2024. 2024. Available: https://www.olympics.com/ioc/news/world-mental-health-day-how-the-ioc-ran-the-largest-online-abuse-prevention-programme-ever-conducted-in-sport-at-paris-2024
608 Putukian M. How being injured affects mental health. In: Brown GT, Hainline B, Kroshus E, eds. *Mind, Body and Sport: Understanding and Supporting Student-Athlete Mental Wellness*. Indianapolis, 2014: 72–5.
609 Neal TL, Diamond AB, Goldman S, *et al*. Inter-association recommendations for developing a plan to recognize and refer student-athletes with psychological concerns at the collegiate level: An executive summary of a consensus statement. *J Athl Train* 2013;48:716–20.

Br J Sports Med: first published as 10.1136/bjsports-2025-111514 on 11 August 2026. Downloaded from http://bjsm.bmj.com/ on August 12, 2026 at Universitetsbiblioteket i Tromsø . Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies.

610 Tranaeus U, Gledhill A, Johnson U, *et al*. 50 years of research on the psychology of sport injury: A consensus statement. *Sports Med* 2024;54:1733–48.
611 Rogers DL, Tanaka MJ, Cosgarea AJ, *et al*. How mental health affects injury risk and outcomes in athletes. *Sports Health* 2024;16:222–9.
612 Ivarsson A, Johnson U, Podlog L. Psychological predictors of injury occurrence: A prospective investigation of professional Swedish soccer players. *J Sport Rehabil* 2013;22:19–26.
613 Ivarsson A, Johnson U. Psychological factors as predictors of injuries among senior soccer players. A prospective study. *J Sports Sci Med* 2010;9:347–52.
614 Ivarsson A, Johnson U, Andersen MB, *et al*. Psychosocial factors and sport injuries: Meta-analyses for prediction and prevention. *Sports Med* 2017;47:353–65.
615 Ardern CL, Taylor NF, Feller JA, *et al*. A systematic review of the psychological factors associated with returning to sport following injury. *Br J Sports Med* 2013;47:1120–6.
616 Wiese-Bjornstal DM. Psychology and socioculture affect injury risk, response, and recovery in high-intensity athletes: A consensus statement. *Scand J Med Sci Sports* 2010;20 Suppl 2:103–11.
617 Nippert AH, Smith AM. Psychologic stress related to injury and impact on sport performance. *Phys Med Rehabil Clin N Am* 2008;19:399–418.
618 Kiliç Ö, Aoki H, Goedhart E, *et al*. Severe musculoskeletal time-loss injuries and symptoms of common mental disorders in professional soccer: A longitudinal analysis of 12-month follow-up data. *Knee Surg Sports Traumatol Arthrosc* 2018;26:946–54.
619 Hainline B, Derman W, Vernec A, *et al*. International Olympic Committee consensus statement on pain management in elite athletes. *Br J Sports Med* 2017;51:1245–58.
620 Hainline B, Turner JA, Caneiro JP, *et al*. Pain in elite athletes—neurophysiological, biomechanical and psychosocial considerations: A narrative review. *Br J Sports Med* 2017;51:1259–64.
621 Ivarsson A, Johnson U, Lindwall M, *et al*. Psychosocial stress as a predictor of injury in elite junior soccer: A latent growth curve analysis. *J Sci Med Sport* 2014;17:366–70.
622 Chen AZ, Greaves KM, Fortney TA, *et al*. Psychological factors associated with return to play after ulnar collateral ligament reconstruction: A systematic review. *Orthop J Sports Med* 2022;10:23259671221125746.
623 Nwachukwu BU, Adjei J, Rauck RC, *et al*. How much do psychological factors affect lack of return to play after anterior cruciate ligament reconstruction? A systematic review. *Orthop J Sports Med* 2019;7:2325967119845313.
624 Wiese-Bjornstal DM, Smith AM, Shaffer SM, *et al*. An integrated model of response to sport injury: Psychological and sociological dynamics. *J Appl Sport Psychol* 1998;10:46–69.
625 Forsdyke D, Smith A, Jones M, *et al*. Psychosocial factors associated with outcomes of sports injury rehabilitation in competitive athletes: A mixed studies systematic review. *Br J Sports Med* 2016;50:537–44.
626 Li H, Moreland JJ, Peek-Asa C, *et al*. Preseason anxiety and depressive symptoms and prospective injury risk in collegiate athletes. *Am J Sports Med* 2017;45:2148–55.
627 Yang J, Cheng G, Zhang Y, *et al*. Influence of symptoms of depression and anxiety on injury hazard among collegiate American football players. *Res Sports Med* 2014;22:147–60.
628 Timpka T, Janson S, Jacobsson J, *et al*. Lifetime history of sexual and physical abuse among competitive athletics (track and field) athletes: Cross sectional study of associations with sports and non-sports injury. *Br J Sports Med* 2019;53:1412–7.
629 Fletcher D, Hanton S. Sources of organizational stress in elite sports performers. *Sport Psychol* 2003;17:175–95.
630 Johnson U, Ivarsson A. Psychological predictors of sport injuries among junior soccer players. *Scand J Med Sci Sports* 2011;21:129–36.
631 Pensgaard AM, Ivarsson A, Nilstad A, *et al*. Psychosocial stress factors, including the relationship with the coach, and their influence on acute and overuse injury risk in elite female football players. *BMJ Open Sport Exerc Med* 2018;4:e000317.
632 van der Does HTD, Brink MS, Otter RTA, *et al*. Injury risk is increased by changes in perceived recovery of team sport players. *Clin J Sport Med* 2017;27:46–51.
633 Steffen K, Pensgaard AM, Bahr R. Self-reported psychological characteristics as risk factors for injuries in female youth football. *Scand J Med Sci Sports* 2009;19:442–51.
634 Kellmann M. Preventing overtraining in athletes in high-intensity sports and stress/recovery monitoring. *Scand J Med Sci Sports* 2010;20 Suppl 2:95–102.
635 Staufenbiel SM, Penninx B, Spijker AT, *et al*. Hair cortisol, stress exposure, and mental health in humans: A systematic review. *Psychoneuroendocrinology* 2013;38:1220–35.
636 Chyi T, Lu FJH, Hsieh Y-C, *et al*. Relationship between athletes' history of stressors and sport injury: A systematic review and meta-analysis. *Percept Mot Skills* 2024;131:192–218.
637 Nicholls AR, Levy AR, Grice A, *et al*. Stress appraisals, coping, and coping effectiveness among international cross-country runners during training and competition. *Eur J Sport Sci* 2009;9:285–93.
638 Devantier C. Psychological predictors of injury among professional soccer players. *Sport Science Review* 2011;20:5–36.
639 Burland JP, Toonstra J, Werner JL, *et al*. Decision to return to sport after anterior cruciate ligament reconstruction, Part I: A qualitative investigation of psychosocial factors. *J Athl Train* 2018;53:452–63.
640 Coronado RA, Bley JA, Huston LJ, *et al*. Composite psychosocial risk based on the fear avoidance model in patients undergoing anterior cruciate ligament reconstruction: Cluster-based analysis. *Phys Ther Sport* 2021;50:217–25.
641 Presley J, Bailey L, Maloney K, *et al*. The Influence of mode-of-injury on psychological readiness for return-to-sport following anterior cruciate ligament reconstruction: A matched-controlled study. *Int J Sports Phys Ther* 2021;16:177–84.
642 Chase D, Slicer K, Schatz P. Relationship between standalone performance validity test failure and emotionality among youth/student athletes experiencing prolonged recovery following sports-related concussion. *Dev Neuropsychol* 2020;45:435–45.
643 Kerr ZY, Walton SR, Brett BL, *et al*. Measurement implications on the association between self-reported concussion history and depression: An NFL-LONG study. *Clin Neuropsychol* 2023;37:1136–53.
644 Proctor SL, Boan-Lenzo C. Prevalence of depressive symptoms in male intercollegiate student-athletes and nonathletes. *J Clin Sport Psychol* 2010;4:204–20.
645 Yang J, Peek-Asa C, Corlette JD, *et al*. Prevalence of and risk factors associated with symptoms of depression in competitive collegiate student athletes. *Clin J Sport Med* 2007;17:481–7.
646 Wolanin A, Gross M, Hong E. Depression in athletes: Prevalence and risk factors. *Curr Sports Med Rep* 2015;14:56–60.
647 Rice SM, Purcell R, De Silva S, *et al*. The mental health of elite athletes: A narrative systematic review. *Sports Med* 2016;46:1333–53.
648 Guskiewicz KM, Marshall SW, Bailes J, *et al*. Recurrent concussion and risk of depression in retired professional football players. *Med Sci Sports Exerc* 2007;39:903–9.
649 Huang J-H, Jacobs DF, Derevensky JL, *et al*. Gambling and health risk behaviors among U.S. college student-athletes: Findings from a national study. *J Adolescent Health* 2007;40:390–7.
650 Beals KA, Manore MM. Disorders of the female athlete triad among collegiate athletes. *Int J Sport Nutr Exerc Metab* 2002;12:281–93.
651 Weigand S, Cohen J, Merenstein D. Susceptibility for depression in current and retired student athletes. *Sports Health: A Multidisciplinary Approach* 2013;5:263–6.
652 Lindqvist A, Moberg T, Ehrnborg C, *et al*. Increased mortality rate and suicide in Swedish former elite male athletes in power sports . *Scand Med Sci Sports* 2014;24:1000–5.
653 Armstrong S, Oomen-Early J. Social connectedness, self-esteem, and depression symptomatology among collegiate athletes versus nonathletes. *J Am College Health* 2009;57:521–6.
654 Weinstock J, Whelan JP, Meyers AW, *et al*. Gambling behavior of student-athletes and a student cohort: What are the odds? *J Gambl Stud* 2007;23:13–24.
655 Kerr ZY, Marshall SW, Harding HP Jr, *et al*. Nine-year risk of depression diagnosis increases with increasing self-reported concussions in retired professional football players. *Am J Sports Med* 2012;40:2206–12.
656 Smith AM, Milliner EK. Injured athletes and the risk of suicide. *J Athl Train* 1994;29:337–41.
657 Ardern CL. Anterior cruciate ligament reconstruction-not exactly a one-way ticket back to the preinjury level: A review of contextual factors affecting return to sport after surgery. *Sports Health* 2015;7:224–30.
658 Czuppon S, Racette BA, Klein SE, *et al*. Variables associated with return to sport following anterior cruciate ligament reconstruction: A systematic review. *Br J Sports Med* 2014;48:356–64.
659 Glazer DD. Development and preliminary validation of the Injury-Psychological Readiness to Return to Sport (I-PRRS) Scale. *J Athl Train* 2009;44:185–9.
660 Tripp DA, Stanish W, Ebel-Lam A, *et al*. Fear of reinjury, negative affect, and catastrophizing predicting return to sport in recreational athletes with anterior cruciate ligament injuries at 1 year postsurgery. *Rehabil Psychol* 2007;52:74–81.
661 Gignac MA, Cao X, Ramanathan S, *et al*. Perceived personal importance of exercise and fears of re-injury: A longitudinal study of psychological factors related to activity after anterior cruciate ligament reconstruction. *BMC Sports Sci Med Rehabil* 2015;7:4.
662 Flanigan DC, Everhart JS, Glassman AH. Psychological factors affecting rehabilitation and outcomes following elective orthopaedic surgery. *Journal of the American Academy of Orthopaedic Surgeons* 2015;23:563–70.
663 Everhart JS, Best TM, Flanigan DC. Psychological predictors of anterior cruciate ligament reconstruction outcomes: A systematic review. *Knee Surg Sports Traumatol Arthrosc* 2015;23:752–62.
664 Podlog L, Banham SM, Wadey R, *et al*. Psychological readiness to return to competitive sport following injury: A qualitative study. *Sport Psychol* 2015;29:1–14.
665 Podlog L, Dimmock J, Miller J. A review of return to sport concerns following injury rehabilitation: Practitioner strategies for enhancing recovery outcomes. *Physical Therapy in Sport* 2011;12:36–42.
666 Ardern CL, Taylor NF, Feller JA, *et al*. Sports participation 2 years after anterior cruciate ligament reconstruction in athletes who had not returned to sport at 1 year: A prospective follow-up of physical function and psychological factors in 122 athletes. *Am J Sports Med* 2015;43:848–56.

Br J Sports Med: first published as 10.1136/bjsports-2025-111514 on 11 August 2026. Downloaded from http://bjsm.bmj.com/ on August 12, 2026 at Universitetsbiblioteket i Tromsoe . Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies.

667 Gajardo-Burgos R, Valdebenito-Tejos C, Gálvez-García G, *et al*. Pain and psychological readiness to return to sport in elite volleyball players: A cross-sectional study. *Int J Environ Res Public Health* 2023;20:2492.
668 Baez S, Jochimsen K. Current clinical concepts: Integration of psychologically informed practice for management of patients with sport-related injuries. *J Athl Train* 2023;58:687–96.
669 Gennarelli SM, Brown SM, Mulcahey MK. Psychosocial interventions help facilitate recovery following musculoskeletal sports injuries: A systematic review. *Phys Sportsmed* 2020;48:370–7.
670 Kortlever JTP, Karyampudi P, Ottenhoff JSE, *et al*. Using the Tampa Scale for Kinesiophobia Short Form in patients with upper extremity specific limitations. *Hand (N Y)* 2021;16:847–53.
671 Woby SR, Roach NK, Urmston M, *et al*. Psychometric properties of the TSK-11: A shortened version of the Tampa Scale for Kinesiophobia. *Pain* 2005;117:137–44.
672 Walker N, Thatcher J, Lavallee D. A preliminary development of the Re-Injury Anxiety Inventory (RIAI). *Phys Ther Sport* 2010;11:23–9.
673 Waddell G, Newton M, Henderson I, *et al*. A Fear-Avoidance Beliefs Questionnaire (FABQ) and the role of fear-avoidance beliefs in chronic low back pain and disability. *Pain* 1993;52:157–68.
674 Webster KE, Feller JA, Lambros C. Development and preliminary validation of a scale to measure the psychological impact of returning to sport following anterior cruciate ligament reconstruction surgery. *Phys Ther Sport* 2008;9:9–15.
675 Nicholas MK. The Pain Self-Efficacy Questionnaire: Taking pain into account. *Eur J Pain* 2007;11:153–63.
676 Sullivan MJL, Bishop SR, Pivik J. The Pain Catastrophizing Scale: Development and validation. *Psychol Assess* 1995;7:524–32.
677 Gómez-Piqueras P, Ardern C, Prieto-Ayuso A, *et al*. Psychometric analysis and effectiveness of the Psychological Readiness of Injured Athlete to Return to Sport (PRIA-RS) questionnaire on injured soccer players. *Int J Environ Res Public Health* 2020;17:1536.
678 Gómez-Piqueras P, Ruíz-Barquín R, Olmedilla A. Translation and adaptation to English of a questionnaire to determine the psychological readiness of the injured football player. *Revista de Psicologia Del Deporte* 2020;29:39–48.
679 Conti C, di Fronso S, Robazza C, *et al*. The Injury-Psychological Readiness to return to sport (I-PRRS) scale and the Sport Confidence Inventory (SCI): A cross-cultural validation. *Phys Ther Sport* 2019;40:218–24.
680 Aizawa J, Hirohata K, Ohji S, *et al*. Cross-sectional study on relationships between physical function and psychological readiness to return to sport after anterior cruciate ligament reconstruction. *BMC Sports Sci Med Rehabil* 2022;14:97.
681 Piussi R, Beischer S, Thomeé R, *et al*. Greater psychological readiness to return to sport, as well as greater present and future knee-related self-efficacy, can increase the risk for an anterior cruciate ligament re-rupture: A matched cohort study. *Arthroscopy* 2022;38:1267.
682 Sterling M, Smeets R, Keijzers G, *et al*. Physiotherapist-delivered stress inoculation training integrated with exercise versus physiotherapy exercise alone for acute whiplash-associated disorder (StressModex): A randomised controlled trial of a combined psychological/physical intervention. *Br J Sports Med* 2019;53:1240–7.
683 Newman M, Ravindranath D. Managing a psychiatric emergency. *Psychiatr Times* 2010;27:17–20.
684 Patel MX, Sethi FN, Barnes TR, *et al*. Joint BAP NAPICU evidence-based consensus guidelines for the clinical management of acute disturbance: De-escalation and rapid tranquillisation. *J Psychopharmacol* 2018;32:601–40.
685 Currie A, Johnston A. Mental health emergencies. In: Reardon CL, ed. *Mental Care for Elite Athletes*. Springer, Cham, 2022: 185–95. Available: https://doi.org/10.1007/978-3-031-08364-8_20
686 Currie A, McDuff D, Johnston A, *et al*. Management of mental health emergencies in elite athletes: A narrative review. *Br J Sports Med* 2019;53:772–8.
687 Purcell R, Currie A. Mental health crises in sport. In: Strauss B, ed. *The Psychology of Crises in Sport: Causes, Consequences and Solutions*. Springer, 2024: 165–79.
688 Mandelli L, Arminio A, Atti A-R, *et al*. Suicide attempts in eating disorder subtypes: A meta-analysis of the literature employing DSM-IV, DSM-5, or ICD-10 diagnostic criteria. *Psychol Med* 2019;49:1237–49.
689 Smith AR, Fink EL, Anestis MD, *et al*. Exercise caution: Over-exercise is associated with suicidality among individuals with disordered eating. *Psychiatry Res* 2013;206:246–55.
690 Hussain B, Khalily MT, Khalily MA. Suicidal tendencies and psychiatric symptoms as consequence of anabolic androgenic steroid usage among athletes in District Rawalpindi. *J Pak Med Assoc* 2022;72:616–9.
691 Avois L, Robinson N, Saudan C, *et al*. Central nervous system stimulants and sport practice. *Br J Sports Med* 2006;40:i16–20.
692 Knecht T. . Methamphetamine-related mental disorders. Swiss Archives of Neurology, Psychiatry and Psychotherapy; 2004.155. 16–23.
693 Di Forti M, Quattrone D, Freeman TP, *et al*. The contribution of cannabis use to variation in the incidence of psychotic disorder across Europe (EU-GEI): A multicentre case-control study. *Lancet Psychiatry* 2019;6:427–36.
694 Putukian M. The psychological response to injury in student athletes: A narrative review with a focus on mental health. *Br J Sports Med* 2016;50:145–8.
695 Green GA, Uryasz FD, Petr TA, *et al*. NCAA study of substance use and abuse habits of college student-athletes. *Clin J Sport Med* 2001;11:51–6.
696 Blakelock DJ, Chen MA, Prescott T. Psychological distress in elite adolescent soccer players following deselection. *J Clin Sport Psychol* 2016;10:59–77.
697 Edwards C, Berlin JT. The psychiatric and emergency care of vips and athletes in crisis. In: Glick R, Zeller S, Fishkind A, eds. *Emergency Psychiatry: Principles and Practice*. 2020: 459–69.
698 Helitzer M. Taming the beast: Riding out a sports crisis. *Sport Marketing Quarterly* 1992;1:33–42.
699 Edwards C. Mental health emergencies in athletes. In: Baron D, ed. *Sport and Mental Health*. Champaign, IL: Springer, 2023: 68–79.
700 Kroshus-Havril E, Wheelhouse C, Houle J, *et al*. Consensus recommendations to inform an update of the NCAA Mental Health Best Practices. *Br J Sports Med* 2025;59:968–80.
701 Mountjoy M, Sloan S, Ali-Zada M, *et al*. International Olympic Committee consensus-driven guidelines for athlete mental health support at sporting events. *Br J Sports Med* 2025;59:1459–70.
702 Young J, Neil ER, Granger K, *et al*. Preparedness, confidence, and best practices in preventing, recognizing, and managing mental health cases in National Collegiate Athletic Association institutions. *J Athl Train* 2023;58:156–62.
703 Richmond J, Berlin J, Fishkind A, *et al*. Verbal de-escalation of the agitated patient: Consensus statement of the American Association for Emergency Psychiatry Project BETA De-escalation Workgroup. *WestJEM* 2012;13:17–25.
704 Pilkington V, Rice S, Olive L, *et al*. Athlete mental health and wellbeing during the transition into elite sport: Strategies to prepare the system. *Sports Med - Open* 2024;10:24.
705 Reardon CL, Benoy R, Hitchcock M. Eating disorders and disordered eating in athletes during times of transition: A narrative systematic review of the literature. *Adv Psychiatry Behavioral Health* 2023;3:57–68.
706 Stambulova NB. Crisis-transitions in athletes: Current emphases on cognitive and contextual factors. *Curr Opin Psychol* 2017;16:62–6.
707 Storm LK, Larsen CH, Henriksen K. "When you Understand the environment, you can navigate the transition better": Supporting professional football players in transitions to clubs abroad. *J Sport Psychol Action* 2025;16:122–34.
708 Gentile PC, Arth ZW. "A beautiful experience": The cultural transitions of Latin American Minor League Baseball (MiLB) players. *Communication & Sport* 2024;12:684–707.
709 Voorheis P, Silver M, Consonni J. Adaptation to life after sport for retired athletes: A scoping review of existing reviews and programs. *PLoS One* 2023;18:e0291683.
710 Mannes ZL, Waxenberg LB, Cottler LB, *et al*. Prevalence and correlates of psychological distress among retired elite athletes: A systematic review. *Int Rev Sport Exerc Psychol* 2019;12:265–94.
711 Park S, Lavallee D, Tod D. Athletes' career transition out of sport: A systematic review. *Int Rev Sport Exerc Psychol* 2013;6:22–53.
712 Filbay S, Pandya T, Thomas B, *et al*. Quality of life and life satisfaction in former athletes: A systematic review and meta-analysis. *Sports Med* 2019;49:1723–38.
713 Bočková K, Anthony Procházka D, Lovásik J, *et al*. Retirement from elite sports: Key factors shaping athletes' transition to post-sport life in the Czech Republic. *Environ Soc Psychol* 2025;10:3800.
714 Simon JE, Lorence M, Docherty CL. Health-related quality of life in former National Collegiate Athletic Association Division I collegiate athletes compared with noncollegiate athletes: A 5-year follow-up. *J Athl Train* 2021;56:331–8.
715 Shander K, Petrie T, Moore EWG. Retired college student athletes' psychological well-being: A prospective analysis. *Journal of Loss and Trauma* 2024;29:828–43.
716 DeFreese JD, Walton SR, Kerr ZY, *et al*. Transition-related psychosocial factors and mental health outcomes in former National Football League players: An NFL-LONG study. *J Sport Exerc Psychol* 2022;44:169–76.
717 Esopenko C, Coury JR, Pieroth EM, *et al*. The psychological burden of retirement from sport. *Curr Sports Med Rep* 2020;19:430–7.
718 Zimet D, McDuff D, Iannone VN, *et al*. Elite athletes' subjective wellbeing trajectories throughout the retirement transition: An analysis using Ruthven's (2022) transition model. *Int Rev Psychiatry* 2024;36:272–83.
719 McDuff DR, Garvin M. Working with sports organizations and teams. *Int Rev Psychiatry* 2016;15:595–605.
720 DeFreese JD, Weight E, DeCicco J, *et al*. Transition experiences of former collegiate women's soccer athletes. *JIS* 2021;14.
721 Kiefer HR, Petrie TA, Walls R. The transition from collegiate sport: An analysis of the current retirement planning practices of NCAA institutions. *Journal for the Study of Sports and Athletes in Education* 2023;17:66–82.
722 Wilkinson RJ. A literature review exploring the mental health issues in academy football players following career termination due to deselection or injury and how counselling could support future players. *Couns and Psychother Res* 2021;21:859–68.
723 Australian Institute of Sports (AIS). Mental health referral network. 2022. Available: https://www.ais.gov.au/MHRN
724 Brockett CL, Stansen C, Bourke M, *et al*. Factors that influence mental health and well-being of high-performance athletes from Olympic or Paralympic sport who

Br J Sports Med: first published as 10.1136/bjsports-2025-111514 on 11 August 2026. Downloaded from http://bjsm.bmj.com/ on August 12, 2026 at Universitetsbiblioteket i Tromsoe . Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies.

have transitioned out of national-level or international-level sport: A mixed methods approach. *BMJ Open Sport Exerc Med* 2024;10:e001991.
725 Schmid MJ, De Crom L, Conzelmann A, *et al*. Patterns of reasons for retirement in Olympic athletes from Switzerland: A person-oriented study. *Int J Sport Exerc Psychol* 2025;1–20.
726 Rist B, Quartuccio C, Smith B, *et al*. End on a positive note: An exploration of flourishing, identity, and career confidence in professional Australian footballers transitioning out of football. *Sci Med Footb* 2025;9:76–82.
727 Torregrossa M, Reguela S, Mateos M. Career assistance programs. In: Hackfort D, Schinke RJ, eds. *The Routledge International Encyclopedia of Sport and Exercise Psychology*. London: Routledge, 2020: 73–88.
728 Harry M, Weight E. Post-collegiate athlete transitions and the influence of a coach. *Journal for the Study of Sports and Athletes in Education* 2021;15:219–44.
729 Lyons D. Engagement in sport career transition planning enhances performance: A commentary. *Journal of Loss and Trauma* 2019;24:9–11.
730 Lavallee D. Engagement in sport career transition planning enhances performance. *Journal of Loss and Trauma* 2019;24:1–8.
731 Hong HJ, Minikin B. An international analysis of career assistance programmes for high-performance athletes. *International Journal of Sport Policy and Politics* 2023;15:705–24.
732 Knights S, Sherry E, Ruddock-Hudson M. Investigating elite end-of-athletic-career transition: A systematic review. *J Appl Sport Psychol* 2016;28:291–308.
733 Gouttebarge V, Goedhart E, Kerkhoffs G. Empowering the health of retired professional footballers: The systematic development of an after career consultation and its feasibility. *BMJ Open Sport Exerc Med* 2018;4:e000466.
734 Hong HJ, Fraser I. High-performance athletes' transition out of sport: Developing corporate social responsibility. *International Journal of Sport Policy and Politics* 2023;15:725–41.
735 Fleischman D, Mulcahy R, English P, *et al*. Athlete career transitions: A systematic review and future directions for sport management research. *European Sport Management Quarterly* 2025;25:768–92.
736 International Olympic Committee. Athletic career programme. Available: https://stillmed.olympic.org/media/Document%20Library/OlympicOrg/IOC/What-We-Do/Protecting-Clean-Athletes/Athletes-Space/Athlete-Career-Programme/EN-ACP-IOC-Athlete-Career-Programme-ACP.pdf [Accessed 21 Aug 2024].
737 United States Olympic and Paralympic Committee (USOPC). Professional development program. Available: https://www.usopc.org/professionaldevelopment [Accessed 21 Aug 2024].
738 Brassard S, Dionne P, Bourdon S. Elite athletes' environment and their capability to plan and prepare a career transition out of sport. *Int J Educ Vocat Guidance* 2024;24:413–33.
739 Pettersen SD, Handegård BH, Koposov R, *et al*. Staying in the game? Predicting intention to quit in Norwegian elite and high-level female football players. *Soccer & Society* 2025;26:795–806.
740 Miller B, Barker JB, Hunton E, *et al*. Understanding experiences of and influences on well-being in elite para-sport: A qualitative exploration of U.K. para-athletes. *Sport Exerc Perform Psychol* 2024;13:126–43.
741 Arnold R, Wagstaff CRD, Steadman L, *et al*. The organisational stressors encountered by athletes with a disability. *J Sports Sci* 2017;35:1187–96.
742 Bundon A, Ashfield A, Smith B, *et al*. Struggling to stay and struggling to leave: The experiences of elite Para-athletes at the end of their sport careers. *Psychol Sport Exerc* 2018;37:296–305.
743 Ramey L, Hayano T, Blatz D, *et al*. A comparison of self-reported unmet healthcare needs among adaptive and able-bodied athletes. *PM&R* 2020;12:36–42.
744 Hoare E, Reyes J, Olive L, *et al*. Neurodiversity in elite sport: A systematic scoping review. *BMJ Open Sport Exerc Med* 2023;9:e001575.
745 Meidl V, Dallmann P, Leonhart R, *et al*. Validation of the Patient Health Questionnaire-4 for longitudinal mental health evaluation in elite Para athletes. *PM&R* 2024;16:141–9.
746 Macdougall H, O'Halloran P, Sherry E, *et al*. A pilot randomised controlled trial to enhance well-being and performance of athletes in Para sports. *EUJAPA* 2019;12:7.
747 World Health Organization. Global report on health equity for persons with disabilities. 2022. Available: https://www.who.int/publications/i/item/9789240063600
748 Solmi M, Radua J, Olivola M, *et al*. Age at onset of mental disorders worldwide: Large-scale meta-analysis of 192 epidemiological studies. *Mol Psychiatry* 2022;27:281–95.
749 Daley MM, Reardon CL. Mental health in the youth athlete. *Clin Sports Med* 2024;43:107–26.
750 McGorry PD, Mei C, Dalal N, *et al*. The Lancet Psychiatry Commission on youth mental health. *Lancet Psychiatry* 2024;11:731–74.
751 Arnett JJ, Hughes M. Adolescence and Emerging Adulthood. Boston, MA: Pearson, 2014.
752 Bell DR, Post EG, Biese K, *et al*. Sport specialization and risk of overuse injuries: A systematic review with meta-analysis. *Pediatrics* 2018;142:e20180657.
753 Clark VC, Ulman SM, Erdman AL, *et al*. Athletic identity, anxiety, and depression in moderate to highly specialized female adolescent volleyball players. *Front Psychol* 2025;16:1525074.
754 Walton CC, Rice S, Hutter RI (Vana), *et al*. Mental health in youth athletes. *Advances in Psychiatry and Behavioral Health* 2021;1:119–33.
755 Walton CC, Purcell R, Henderson JL, *et al*. Mental health among elite youth athletes: A narrative overview to advance research and practice. *Sports Health* 2024;16:166–76.
756 Thompson F, Rongen F, Cowburn I, *et al*. The impacts of sports schools on holistic athlete development: A mixed methods systematic review. *Sports Med* 2022;52:1879–917.
757 Gwyther K, Pilkington V, Bailey AP, *et al*. Mental health and well-being of elite youth athletes: A scoping review. *Br J Sports Med* 2024;58:1011–9.
758 Duquesne V, Richard R, Andrieu B, *et al*. Sports experiences of elite athletes with intellectual disabilities and/or autism spectrum disorders. *Sport Soc* 2023;26:854–68.
759 Donti A, Maraki MI, Psychountaki M, *et al*. Eating disorder symptoms and energy deficiency awareness in adolescent artistic gymnasts: Evidence of a knowledge gap. *Nutrients* 2025;17:1699.
760 Larsson M, Lindman I, Hörnell A, *et al*. Prospective study of food intake changes in adolescent elite athletes. *BMJNPH* 2024;7:366–74.
761 Mason L, Connolly J, Devenney LE, *et al*. The sleep, recovery, and nutrition characteristics of elite adolescent athletes. *Sports* 2025;13:50.
762 Xue W, Tao Y, Huang Y, *et al*. Emotional intelligence and burnout among adolescent basketball players: The mediating effect of emotional labor. *Sports* 2024;12:266.
763 Weber S, Puta C, Lesinski M, *et al*. Symptoms of anxiety and depression in young athletes using the Hospital Anxiety and Depression Scale. *Front Physiol* 2018;9:182.
764 Gerber M, Kellmann M, Brand S, *et al*. Differences in mental health outcomes between adolescent elite athletes and peers not engaged in elite sport. *CISS* 2022;7:013.
765 Rosenvinge JH, Sundgot-Borgen J, Pettersen G, *et al*. Are adolescent elite athletes less psychologically distressed than controls? A cross-sectional study of 966 Norwegian adolescents. *Open Access J Sports Med* 2018;9:115–23.
766 Brand R, Wolff W, Hoyer J. Psychological symptoms and chronic mood in representative samples of elite student-athletes, deselected student-athletes and comparison students. *School Ment Health* 2013;5:166–74.
767 Gustafsson H, Kenttä G, Hassmén P, *et al*. Prevalence of burnout in competitive adolescent athletes. *Sport Psychol* 2007;21:21–37.
768 Lima Y, Öz ND, Denerel N, *et al*. Mental health correlates of athletic shame and guilt among elite-level youth athletes in Turkey. *J Sci Med Sport* 2024;27:72–7.
769 Appleton PR, Hall HK, Hill AP. Relations between multidimensional perfectionism and burnout in junior-elite male athletes. *Psychol Sport Exerc* 2009;10:457–65.
770 Nixdorf I, Beckmann J, Nixdorf R. Psychological predictors for depression and burnout among German junior elite athletes. *Front Psychol* 2020;11:601.
771 Nixdorf I, Frank R, Beckmann J. Comparison of athletes' proneness to depressive symptoms in individual and team sports: Research on psychological mediators in junior elite athletes. *Front Psychol* 2016;7:893.
772 Xiang J, Gao J, Gao Y. The relationship between parenting style and the stigma of mental illness in adolescent athletes: The chain mediating role of self-esteem and mental toughness. *Curr Psychol* 2024;43:23130–43.
773 Koua AM, Djo Bi Djo F, Kouadio RN, *et al*. Survey of non-conventional mental health care facilities in Côte d'Ivoire: First stage. *Int J Ment Health Syst* 2021;15:83.
774 Nguendo-Yongsi HB. Access to health care in African cities: therapeutic pathways of city dwellers with mental health problems in Yaoundé – Cameroon. *Cities & Health* 2022;6:113–22.
775 Tachom Waffo B, Hauw D. Mental health of young talented football players in an African context: A cross-sectional observational study. *Compr Psychiatry* 2024;135:152519.
776 Wandik Y, Dina D, Guntoro TS, *et al*. Interrelation of mental toughness, religiosity, and happiness of elite adolescent athletes based on gender, type of sport, and level of education. *Retos* 2024;56:981–9.
777 Oja P, Memon AR, Titze S, *et al*. Health benefits of different sports: A systematic review and meta-analysis of longitudinal and intervention studies including 2.6 million adult participants. *Sports Med Open* 2024;10:46.
778 Pascoe M, Bailey AP, Craike M, *et al*. Physical activity and exercise in youth mental health promotion: A scoping review. *BMJ Open Sport Exerc Med* 2020;6:e000677.
779 Mahindru A, Patil P, Agrawal V. Role of physical activity on mental health and well-being: A review. *Cureus* 2023;15:e33475.
780 Eather N, Wade L, Pankowiak A, *et al*. The impact of sports participation on mental health and social outcomes in adults: A systematic review and the "Mental Health through Sport" conceptual model. *Syst Rev* 2023;12:102.
781 Vella SA, Aidman E, Teychenne M, *et al*. Optimising the effects of physical activity on mental health and wellbeing: A joint consensus statement from Sports Medicine Australia and the Australian Psychological Society. *J Sci Med Sport* 2023;26:132–9.
782 Kroshus-Havril E, Reardon CL, Gouttebarge V, *et al*. Research priorities for advancing mental health in elite sport: a companion to the IOC consensus statement on mental health in elite athletes. *Br J Sports Med* 2026;60:1187–90.
783 International Olympic Committee (IOC). Athlete 365. 2024. Available: https://olympics.com/athlete365

Br J Sports Med: first published as 10.1136/bjsports-2025-111514 on 11 August 2026. Downloaded from http://bjsm.bmj.com/ on August 12, 2026 at Universitetsbiblioteket i Tromsoe . Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies.

784 International Olympic Committee (IOC). IOC Diploma in Mental Health in Elite Sport. 2024. Available: https://www.sportsoracle.com/course/ioc-diploma-in-mental-health-in-elite-sport/
785 International Olympic Committee (IOC). IOC Certificate Program in Mental Health in Elite Sport. 2024. Available: https://www.sportsoracle.com/course/ioc-certificate-in-mental-health-in-elite-sport/
786 International Olympic Committee (IOC). Mental Health Guidelines for Sporting Events. 2023. Available: https://stillmed.olympics.com/media/Documents/Athletes/Safeguarding/Mental-Health-Guidelines-for-Events-EN.pdf
787 Dindo L, Van Liew JR, Arch JJ. Acceptance and commitment therapy: A transdiagnostic behavioral intervention for mental health and medical conditions. *Neurotherapeutics* 2017;14:546–53.
788 Donohue B, Gavrilova Y, Galante M, *et al*. Controlled evaluation of an optimization approach to mental health and sport performance. *J Clin Sport Psychol* 2018;12:234–67.
789 Peprah K, Argáez C. Dialectical Behavioral Therapy for Adults with Mental Illness: A Review of Clinical Effectiveness and Guidelines. Ottawa (ON): Canadian Agency for Drugs and Technologies in Health, 2017.
790 Ben-Porath DD, Koons CR. Telephone coaching in dialectical behavior therapy: A decision-tree model for managing inter-session contact with clients. *Cogn Behav Pract* 2005;12:448–60.
791 Dimeff LA, Linehan MM. Dialectical behavior therapy for substance abusers. *Addict Sci Clin Pract* 2008;4:39–47.
792 Simpson E, Carroll C, Sutton A, *et al*. Clinical and cost-effectiveness of eye movement desensitization and reprocessing for treatment and prevention of post-traumatic stress disorder in adults: A systematic review and meta-analysis. *Br J Psychol* 2025;116:1128–49.
793 Markowitz JC, Weissman MM. Interpersonal psychotherapy: Principles and applications. *World Psychiatry* 2004;3:136–9.
794 Goldberg SB, Tucker RP, Greene PA, *et al*. Mindfulness-based interventions for psychiatric disorders: A systematic review and meta-analysis. *Clin Psychol Rev* 2018;59:52–60.
795 Myall K, Montero-Marin J, Gorczynski P, *et al*. Effect of mindfulness-based programmes on elite athlete mental health: A systematic review and meta-analysis. *Br J Sports Med* 2023;57:99–108.
796 Boyd JE, Lanius RA, McKinnon MC. Mindfulness-based treatments for posttraumatic stress disorder: A review of the treatment literature and neurobiological evidence. *J Psychiatry Neurosci* 2018;43:7–25.
797 Agley J, Walker BB, Gassman RA. Adaptation of alcohol and drug screening, brief intervention and referral to treatment (SBIRT) to a department of intercollegiate athletics: The COMPASS project. *Health Educ J* 2013;72:647–59.
798 Opland C, Torrico TJ. Psychodynamic therapy. In: *Statpearls*. Treasure Island (FL): StatPearls Publishing, 2025.
799 Haynes PL, Gengler D, Kelly M. Social rhythm therapies for mood disorders: An update. *Curr Psychiatry Rep* 2016;18:75.
800 Vermeulen-Oskam E, Franklin C, Van't Hof LPM, *et al*. The current evidence of solution-focused brief therapy: A meta-analysis of psychosocial outcomes and moderating factors. *Clin Psychol Rev* 2024;114:102512.
801 Grover S, Avasthi A, Jagiwala M. Clinical practice guidelines for practice of supportive pychotherapy. *Indian J Psychiatry* 2020;62:173.
802 Longo LP, Johnson B. Addiction: Part I. Benzodiazepines--side effects, abuse risk and alternatives. *Am Fam Physician* 2000;61:2121–8.
803 Zandonai T, Lugoboni F, Zamboni L. A risk for athletes: when the desire to sleep becomes a nightmare. A brief case report on benzodiazepine addiction. *Psychopharmacology (Berl)* 2018;235:3359–60.
804 Zamboni L, Lugoboni F, Zandonai T. Benzodiazepine abuse among athletes: Pain relief or just a weapon against insomnia? A clinical case study. *Scand J Med Sci Sports* 2019;29:1937–40.
805 Zandonai T, Peiró AM, Fusina F, *et al*. Benzodiazepines in sport, an underestimated problem: Recommendations for sports medicine physicians' practice. *Front Psychiatry* 2022;13:1066330.
806 Holgado D, Manresa-Rocamora A, Zamboni L, *et al*. The effect of benzodiazepines on exercise in healthy adult participants: A systematic review. *J Addict Dis* 2022;40:336–44.
807 Atkinson G, Drust B, Reilly T, *et al*. The relevance of melatonin to sports medicine and science. *Sports Med* 2003;33:809–31.
808 Farjallah M, Graja A, Mahmoud L, *et al*. Effects of melatonin ingestion on physical performance and biochemical responses following exhaustive running exercise in soccer players. *Bs* 2022;39:473–9.
809 Watson P, Hasegawa H, Roelands B, *et al*. Acute dopamine/noradrenaline reuptake inhibition enhances human exercise performance in warm, but not temperate conditions. *J Physiol (Lond)* 2005;565:873–83.
810 Davidson J. Seizures and bupropion: a review. *J Clin Psychiatry* 1989;50:256–61.
811 Fowler NO, McCall D, Chou TC, *et al*. Electrocardiographic changes and cardiac arrhythmias in patients receiving psychotropic drugs. *Am J Cardiol* 1976;37:223–30.
812 Tanayan C, Reddy S, Shah AB, *et al*. A cyclist on a tricyclic: Exercise intolerance due to chronotropic incompetence. *JACC Case Rep* 2022;4:1335–40.
813 de Zwaan M. Exercise and antidepressant serum levels. *Biol Psychiatry* 1992;32:210–1.
814 Gorczynski P, Aron CM, Reardon CL. Loneliness and emotional blunting in elite sport. *Advances in Psychiatry and Behavioral Health* 2024;4:1–9.
815 Cordery P, Peirce N, Maughan RJ, *et al*. Dopamine/noradrenaline reuptake inhibition in women improves endurance exercise performance in the heat. *Scandinavian Med Sci Sports* 2017;27:1221–30.
816 Piacentini MF, Meeusen R, Buyse L, *et al*. Hormonal responses during prolonged exercise are influenced by a selective DA/NA reuptake inhibitor. *Br J Sports Med* 2004;38:129–33.
817 Kavanagh JJ, McFarland AJ, Taylor JL. Enhanced availability of serotonin increases activation of unfatigued muscle but exacerbates central fatigue during prolonged sustained contractions. *J Physiol* 2019;597:319–32.
818 Strachan AT, Leiper JB, Maughan RJ. Paroxetine administration failed [corrected] to influence human exercise capacity, perceived effort or hormone responses during prolonged exercise in a warm environment. *Exp Physiol* 2004;89:657–64.
819 Strüder HK, Hollmann W, Platen P, *et al*. Influence of paroxetine, branched-chain amino acids and tyrosine on neuroendocrine system responses and fatigue in humans. *Horm Metab Res* 1998;30:188–94.
820 Teixeira-Coelho F, Uendeles-Pinto JP, Serafim ACA, *et al*. The paroxetine effect on exercise performance depends on the aerobic capacity of exercising individuals. *J Sports Sci Med* 2014;13:232–43.
821 Thorstensen JR, Taylor JL, Tucker MG, *et al*. Enhanced serotonin availability amplifies fatigue perception and modulates the TMS-induced silent period during sustained low-intensity elbow flexions. *J Physiol (Lond)* 2020;598:2685–701.
822 Wilson WM, Maughan RJ. Evidence for a possible role of 5-hydroxytryptamine in the genesis of fatigue in man: administration of paroxetine, a 5-HT re-uptake inhibitor, reduces the capacity to perform prolonged exercise. *Exp Physiol* 1992;77:921–4.
823 Parise G, Bosman MJ, Boecker DR, *et al*. Selective serotonin reuptake inhibitors: Their effect on high-intensity exercise performance. *Arch Phys Med Rehabil* 2001;82:867–71.
824 WADA. WADA monitoring program. 2025. Available: https://www.wada-ama.org/en/resources/monitoring-program [Accessed 4 Jun 2025].
825 Du Y, Li Q, Dou Y, *et al*. Side effects and cognitive benefits of buspirone: A systematic review and meta-analysis. *Heliyon* 2024;10:e28918.
826 Forton K, Lamotte M, Gillet A, *et al*. Beta-adrenergic receptor blockade effects on cardio-pulmonary exercise testing in healthy young adults: A randomized, placebo-controlled trial. *Sports Med Open* 2022;8:150.
827 Marvin G, Sharma A, Aston W, *et al*. The effects of buspirone on perceived exertion and time to fatigue in man. *Exp Physiol* 1997;82:1057–60.
828 Cowan DA. Drug abuse. In: HarriesM, WilliamsC, StanishWD, eds. *Oxford textbook of sports medicine*. 1994: 314–29.
829 Roelands B, Hasegawa H, Watson P, *et al*. The effects of acute dopamine reuptake inhibition on performance. *Med Sci Sports Exerc* 2008;40:879–85.
830 Syed YY. Viloxazine extended-release: A review in attention-deficit/hyperactivity disorder in adults. *CNS Drugs* 2025;39:1355–64.
831 Hirota T, Schwartz S, Correll CU. Alpha-2 agonists for attention-deficit/hyperactivity disorder in youth: A systematic review and meta-analysis of monotherapy and add-on trials to stimulant therapy. *J Am Acad Child Adolesc Psychiatry* 2014;53:153–73.
832 Joo SW, Kim HW. Treatment of children and adolescents with attention deficit/hyperactivity disorder and/or Tourette's disorder with clonidine extended release. *Psychiatry Investig* 2018;15:90–3.
833 Smith GM, Beecher HK. Amphetamine sulfate and athletic performance. *JAMA* 1959;170:542.
834 Chandler JV, Blair SN. The effect of amphetamines on selected physiological components related to athletic success. *Med Sci Sports Exerc* 1980;12:65–9.
835 King M, Rauch LHG, Brooks SJ, *et al*. Methylphenidate enhances grip force and alters brain connectivity. *Med Sci Sports Exerc* 2017;49:1443–51.
836 National Collegiate Athletic Association (NCAA). NCAA banned substances. 2025. Available: https://www.ncaa.org/sports/2015/6/10/ncaa-banned-substances.aspx
837 Jefferson JW, Greist JH, Clagnaz PJ, *et al*. Effect of strenuous exercise on serum lithium level in man. *Am J Psychiatry* 1982;139:1593–5.
838 Miller EB, Pain RW, Skripal PJ. Sweat lithium in manic-depression. *Br J Psychiatry* 1978;133:477–8.
839 Baum AL. Psychopharmacology in athletes. In: Begel D, Burton RW, eds. *Sport Psychiatry*. New York: WW Norton & Company, 2000: 249–59.
840 Finley PR. Drug interactions with lithium: An update. *Clin Pharmacokinet* 2016;55:925–41.
841 Beach SR, Celano CM, Noseworthy PA, *et al*. QTc prolongation, torsades de pointes, and psychotropic medications. *Psychosomatics* 2013;54:1–13.
842 Gessel T, Lin CY. Exertional heatstroke in a marathon runner complicated by concurrent use of an antipsychotic medication affecting thermoregulation. *Clin J Sport Med* 2020;30:e166–8.
843 Gómez Ramos MJ, Miguel González Valverde F, Sánchez Álvarez C, *et al*. Fatal heat stroke in a schizophrenic patient. *Case Rep Crit Care* 2012;2012:924328.
844 Adamson GT, Finlay SE. A comparison of the effects of varying dose levels of oxypertine on mood and physical performance of trained athletes. *Br J Psychiatry* 1966;112:1177–80.

Br J Sports Med: first published as 10.1136/bjsports-2025-111514 on 11 August 2026. Downloaded from http://bjsm.bmj.com/ on August 12, 2026 at Universitetsbiblioteket i Tromsø . Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies.

845 Perez-Cruzado D, Cuesta-Vargas A, Vera-Garcia E, *et al*. Medication and physical activity and physical fitness in severe mental illness. *Psychiatry Res* 2018;267:19–24.
846 Kim DD, Lang DJ, Procyshyn RM, *et al*. Reduced cardiovascular fitness associated with exposure to clozapine in individuals with chronic schizophrenia. *Psychiatry Res* 2018;262:28–33.
847 Hays JT, Ebbert JO. Varenicline for tobacco dependence. *N Engl J Med* 2008;359:2018–24.
848 Gotts JE, Jordt S-E, McConnell R, *et al*. What are the respiratory effects of e-cigarettes? *BMJ* 2019;366:l5275.
849 Mountjoy M, Sundgot-Borgen JK, Burke LM, *et al*. IOC consensus statement on relative energy deficiency in sport (RED-S): 2018 update. *Br J Sports Med* 2018;52:687–97.
850 El Ghoch M, Soave F, Calugi S, *et al*. Eating disorders, physical fitness and sport performance: A systematic review. *Nutrients* 2013;5:5140–60.
851 Green MF. What are the functional consequences of neurocognitive deficits in schizophrenia? *AJP* 1996;153:321–30.
852 Rabinowitz J, Levine SZ, Garibaldi G, *et al*. Negative symptoms have greater impact on functioning than positive symptoms in schizophrenia: Analysis of CATIE data. *Schizophr Res* 2012;137:147–50.

Br J Sports Med: first published as 10.1136/bjsports-2025-111514 on 11 August 2026. Downloaded from http://bjsm.bmj.com/ on August 12, 2026 at Universitetsbiblioteket i Tromsoe . Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies.